

NOVEL ANTIFUNGAL AGENT CONTAINING HETEROCYCLIC COMPOUND

Technical Field

[0001]

5           The present invention relates to a novel antifungal agent containing a heterocyclic compound.

Background Art

[0002]

10           In recent years, managements of opportunistic infections have become more and more significant more than ever because of an increase in the number of elderly people and immunocompromised patients as a result of advanced chemotherapies or the like. As demonstrated by the fact that opportunistic infections are occurring one after another by different weakly avirulent bacteria, it is shown that the problem of infectious disease will not ends as long as there are underlying diseases that  
15           diminish the immune functions of patients. Consequently, new strategies for infectious diseases control, including the problem of resistant bacteria, will be one of the important issues in the soon-to-come aged society.

20           In the field of antifungal agents, heretofore, for instance, amphotericine B which is based on a polyene skeleton, fluconazole, itraconazole and voriconazole which are based on an azole skeleton, or the like, have been developed for the treatment of deep seated mycoses. Among pre-existing drugs already available commercially are many agents having similar mechanism of action, and currently, the appearance of azole-resistant fungi or the like has been problems.

25           In recent years, as a 1,3- $\beta$ -glucan synthetase inhibitor with a novel mechanism, naturally occurring compound-derived cyclic hexapeptides caspofungin

and micafungin or the like, have been developed; however, from the fact that these agents only exist in injectable form, they are not yet sufficient practically as antifungal agents.

Since there have been the situations that the pre-existing antifungal agents  
5 are insufficient for treatment of the deep seated mycoses, there is a demand and need for development of agents which are based on a novel mechanism and are of high safety.

As the prior art related to antifungal agents based on such a novel mechanism, Patent Document 1 describes heterocyclic compounds which  
10 demonstrates effects against the onset, progress, and persistence of infections by inhibiting the expression of cell wall proteins, inhibiting the cell wall assembly and also adhesion onto cells, and preventing pathogens from showing pathogenicity, with the process which transports GPI (Glycosylphosphatidylinositol)-anchored proteins to the cell wall being inhibited. However, groups of the compounds disclosed in Patent  
15 Document 1 have 2-benzyl pyridine moieties as the common structure, clearly differing structurally from compounds according to the present invention. In addition, the groups of the compounds disclosed in Patent Document 1 bear the problem that, although these compounds demonstrate activities *in vitro*, they are easily metabolized inside the body, or the like.

20 Meanwhile, there are Patent Documents 2 through 8 as the prior art that disclose structurally most similar compounds to the heterocyclic compound (I) according to the present invention. Patent Document 2 discloses N-(4-pyridyl)carboxamide derivatives having the effects as pesticides, in particular as insecticides, acaricides and nematicides. Patent Documents 3 through 6 disclose  
25 2-aryloxy nicotinamide derivatives having phosphodiesterase 4 (PDE4) inhibitory

action. Patent Document 7 discloses 6-(arylamino)nicotinamide derivatives having cannabinoid receptor modulation action and Patent Document 8 discloses 6-(aryloxy)nicotinamide derivatives having  $\text{Na}^+/\text{Ca}^{2+}$  exchanger inhibitory action. However, Patent Documents 2 through 8 do not disclose the compounds according to  
5 the present invention. In addition, Patent Documents 2 through 8 do not describe that the compounds disclosed in these documents exhibit antifungal and antimalarial actions for Candida, Aspergillus, Cryptococcus or the like which are general fungal strains in human mycosis at all.

[Patent Document 1] International Publication No. WO 02/04626 pamphlet;

10 [Patent Document 2] US. Patent No. 5852042 Specification;

[Patent Document 3] European Patent Application No. 1229034 specification;

[Patent Document 4] International Publication No. WO 02/060875 pamphlet;

[Patent Document 5] International Publication No. WO 02/060896 pamphlet;

[Patent Document 6] International Publication No. WO 03/068232 pamphlet;

15 [Patent Document 7] International Publication No. WO 2004/029027 pamphlet; and

[Patent Document 8] International Publication No. WO 2004/000813 pamphlet.

[Disclosure of the invention]


[0003]

20 It is an object of the present invention to 1) provide an antifungal agent which has excellent antifungal action not found in the antifungal agents in the prior art, and which is also excellent in terms of property, safety and metabolic stability, and 2) provide an excellent antimalarial agent.

In view of the aforementioned circumstances, the present inventors have  
25 intensively studied to attain the above object. As a result, the present inventions

have succeeded in synthesizing novel heterocycle-containing compounds represented by the following formula (I):

[0004]




5 [0005]

which has the chemical structural characteristics of a heterocyclic group A and a heterocyclic group or a phenyl group E being bonded through a linker X, and have discovered that these compounds have excellent antifungal action, so as to arrive at the present invention.

10 That is to say, the present invention provides:

[1]: an antifungal agent comprising a compound represented by the formula (I), or a salt or a hydrate thereof:

[0006]



15 [0007]

[wherein A represents a 5- to 10-membered heterocyclic group containing at least one nitrogen atom;

X represents a group represented by the formula  $\text{-NH-C(=Y)-(CH}_2\text{)}_n\text{-}$ , a group represented by the formula  $\text{-C(=Y)-NH-(CH}_2\text{)}_n\text{-}$ , a group represented by the formula  $\text{-C(=Z)-(CH}_2\text{)}_n\text{-}$ , a group represented by the formula  $\text{-CH}_2\text{-NH-(CH}_2\text{)}_n\text{-}$ , a group represented by the formula  $\text{-NH-CH}_2\text{-(CH}_2\text{)}_n\text{-}$  or a group represented by the formula  $\text{-Z-CH}_2\text{-(CH}_2\text{)}_n\text{-}$ ;

Y represents an oxygen atom, a sulfur atom or  $\text{NR}^{\text{Y}}$  (wherein  $\text{R}^{\text{Y}}$  represents a



C<sub>1-6</sub> alkoxy group or a cyano group);

Z represents an oxygen atom or a sulfur atom;

n represents an integer from 0 to 3;

E represents a furyl group, a thienyl group, a pyrrolyl group, a pyridyl group, a  
5 tetrazolyl group, a thiazolyl group, a pyrazolyl group or a phenyl group;

with the proviso that A may contain 1 to 3 substituents selected from the  
following substituent groups a-1 and a-2, and that E has one or two substituents  
selected from the following substituent groups a-1 and a-2;

<Substituent group a-1>

- 10 Substituent group a-1 represents the group consisting of: a halogen atom, a hydroxyl  
group, a mercapto group, a cyano group, a carboxyl group, an amino group, a  
carbamoyl group, a C<sub>1-6</sub> alkyl group, a C<sub>2-6</sub> alkenyl group, a C<sub>2-6</sub> alkynyl group, a C<sub>3-8</sub>  
cycloalkyl group, a C<sub>6-10</sub> aryl group, a 5- to 10-membered heterocyclic group, a C<sub>3-8</sub>  
cycloalkyl C<sub>1-6</sub> alkyl group, a C<sub>3-8</sub> cycloalkylidene C<sub>1-6</sub> alkyl group, a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkyl  
15 group, a 5- to 10-membered heterocyclic C<sub>1-6</sub> alkyl group, a C<sub>1-6</sub> alkoxy group, a C<sub>2-6</sub>  
alkenyloxy group, a C<sub>2-6</sub> alkynyloxy group, a C<sub>3-8</sub> cycloalkoxy group, a C<sub>6-10</sub> aryloxy  
group, a C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkoxy group, a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkoxy group, a 5- to  
10-membered heterocyclic C<sub>1-6</sub> alkoxy group, a C<sub>1-6</sub> alkylthio group, a C<sub>2-6</sub> alkenylthio  
group, a C<sub>2-6</sub> alkynylthio group, a C<sub>3-8</sub> cycloalkylthio group, a C<sub>6-10</sub> arylthio group, a  
20 C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkylthio group, a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkylthio group, a 5- to  
10-membered heterocyclic C<sub>1-6</sub> alkylthio group, a mono-C<sub>1-6</sub> alkylamino group, a  
mono-C<sub>2-6</sub> alkenylamino group, a mono-C<sub>2-6</sub> alkynylamino group, a mono-C<sub>3-8</sub>  
cycloalkylamino group, a mono-C<sub>6-10</sub> arylamino group, a mono-C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub>  
alkylamino group, a mono-C<sub>6-10</sub> aryl C<sub>1-6</sub> alkylamino group, a mono-5- to  
25 10-membered heterocyclic C<sub>1-6</sub> alkylamino group, a di-C<sub>1-6</sub> alkylamino group, a N-C<sub>2-6</sub>

alkenyl-N-C<sub>1-6</sub> alkylamino group, a N-C<sub>2-6</sub> alkynyl-N-C<sub>1-6</sub> alkylamino group, a N-C<sub>3-8</sub> cycloalkyl-N-C<sub>1-6</sub> alkylamino group, a N-C<sub>6-10</sub> aryl-N-C<sub>1-6</sub> alkylamino group, a N-C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkyl-N-C<sub>1-6</sub> alkylamino group, a N-C<sub>6-10</sub> aryl C<sub>1-6</sub> alkyl-N-C<sub>1-6</sub> alkylamino group, a N-5- to 10-membered heterocyclic C<sub>1-6</sub> alkyl-N-C<sub>1-6</sub> alkylamino group, a C<sub>1-6</sub> alkylcarbonyl group, a C<sub>1-6</sub> alkoxy carbonyl group, a C<sub>1-6</sub> alkylsulfonyl group, a group represented by the formula -C(=N-R<sup>a1</sup>)R<sup>a2</sup> (wherein R<sup>a1</sup> represents a hydroxyl group or a C<sub>1-6</sub> alkoxy group; R<sup>a2</sup> represents a C<sub>1-6</sub> alkyl group), a C<sub>6-10</sub> aryloxy C<sub>1-6</sub> alkyl group and a 5- to 10-membered heterocycle oxy C<sub>1-6</sub> alkyl group;

<Substituent group a-2>

10 Substituent Group a-2 represents the group consisting of: a C<sub>1-6</sub> alkyl group, a C<sub>2-6</sub> alkenyl group, a C<sub>2-6</sub> alkynyl group, a C<sub>3-8</sub> cycloalkyl group, a C<sub>6-10</sub> aryl group, a 5- to 10-membered heterocyclic group, a C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkyl group, a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkyl group, a 5- to 10-membered heterocyclic C<sub>1-6</sub> alkyl group, a C<sub>1-6</sub> alkoxy group, a C<sub>2-6</sub> alkenyloxy group, a C<sub>2-6</sub> alkynyloxy group, a C<sub>3-8</sub> cycloalkoxy group, a C<sub>6-10</sub> aryloxy group, a C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkoxy group, a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkoxy group, a 5- to 10-membered heterocyclic C<sub>1-6</sub> alkoxy group, a C<sub>1-6</sub> alkylthio group, a C<sub>2-6</sub> alkenylthio group, a C<sub>2-6</sub> alkynylthio group, a C<sub>3-8</sub> cycloalkylthio group, a C<sub>6-10</sub> arylthio group, a C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkylthio group, a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkylthio group, a 5- to 10-membered heterocyclic C<sub>1-6</sub> alkylthio group, a mono-C<sub>1-6</sub> alkylamino group, a  
 20 mono-C<sub>2-6</sub> alkenylamino group, a mono-C<sub>2-6</sub> alkynylamino group, a mono-C<sub>3-8</sub> cycloalkylamino group, a mono-C<sub>6-10</sub> arylamino group, a mono-C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkylamino group, a mono-C<sub>6-10</sub> aryl C<sub>1-6</sub> alkylamino group, a mono-5- to 10-membered heterocyclic C<sub>1-6</sub> alkylamino group, a di-C<sub>1-6</sub> alkylamino group, a N-C<sub>2-6</sub> alkenyl-N-C<sub>1-6</sub> alkylamino group, a N-C<sub>2-6</sub> alkynyl-N-C<sub>1-6</sub> alkylamino group, a N-C<sub>3-8</sub>  
 25 cycloalkyl-N-C<sub>1-6</sub> alkylamino group, a N-C<sub>6-10</sub> aryl-N-C<sub>1-6</sub> alkylamino group, a N-C<sub>3-8</sub>

cycloalkyl C<sub>1-6</sub> alkyl-N-C<sub>1-6</sub> alkylamino group, a N-C<sub>6-10</sub> aryl C<sub>1-6</sub> alkyl-N-C<sub>1-6</sub> alkylamino group, a N-5- to 10-membered heterocyclic C<sub>1-6</sub> alkyl-N-C<sub>1-6</sub> alkylamino group, a C<sub>6-10</sub> aryloxy-C<sub>1-6</sub> alkyl group and a 5- to 10-membered heterocycle oxy C<sub>1-6</sub> alkyl group;

- 5           with the proviso that each group described in the substituent group a-2 has 1 to 3 substituents selected from the following substituent group b;


<Substituent group b>

- Substituent group b represents the group consisting of: a halogen atom, a hydroxyl group, a mercapto group, a cyano group, a carboxyl group, an amino group, a carbamoyl group, a nitro group, a C<sub>1-6</sub> alkyl group, a C<sub>3-8</sub> cycloalkyl group, a C<sub>6-10</sub> aryl group, a 5- to 10-membered heterocyclic group, a C<sub>1-6</sub> alkoxy group, a C<sub>6-10</sub> aryloxy group, a 5- to 10-membered heterocycle oxy group, a C<sub>1-6</sub> alkylcarbonyl group, a C<sub>1-6</sub> alkoxy carbonyl group, a C<sub>1-6</sub> alkylsulfonyl group, a trifluoromethyl group, a trifluoromethoxy group, a mono-C<sub>1-6</sub> alkylamino group, a di-C<sub>1-6</sub> alkylamino group, a mono-C<sub>6-10</sub> arylamino group which may have one amino group or aminosulfonyl group and a N-C<sub>6-10</sub> aryl C<sub>1-6</sub> alkyl-N-C<sub>1-6</sub> alkylamino group which may have one amino group];
- 10
- 15

- [2]: the antifungal agent according to item [1], wherein X represents a group represented by the formula -NH-C(=Y)-CH<sub>2</sub>-, a group represented by the formula -C(=Y)-NH-CH<sub>2</sub>-, a group represented by the formula -CH<sub>2</sub>-NH- or a group represented by the formula -NH-CH<sub>2</sub>- (wherein Y has the same meaning as defined above);
- 20

[3]: a compound represented by the formula (I-a), or a salt or a hydrate thereof:

[0008]



[0009]

[wherein A<sup>1</sup> represents a 3-pyridyl group, a pyrazinyl group, a pyrimidinyl group, a pyrazolyl group, a quinolyl group, an isoquinolyl group, a naphthyldinyl group, a quinoxaliny group, a cinnoliny group, a quinazoliny group, an imidazopyridyl group, a benzothiazolyl group, a benzoxazolyl group, a benzimidazolyl group, an indolyl group, a pyrrolopyridyl group, a thienopyridyl group, a furopyridyl group, a 2,3-dihydro-1*H*-pyrrolo[2,3-*b*]-pyridin-5-yl group or a benzothiadiazolyl group;

X<sup>1</sup> represents a group represented by the formula -NH-C(=Y<sup>1</sup>)- or a group represented by the formula -C(=Y<sup>1</sup>)-NH-;


Y<sup>1</sup> represents an oxygen atom, a sulfur atom or NR<sup>Y1</sup> (wherein R<sup>Y1</sup> represents a C<sub>1-6</sub> alkoxy group or a cyano group);

E represents a furyl group, a thienyl group, a pyrrolyl group, a pyridyl group, a tetrazolyl group, a thiazolyl group, a pyrazolyl group or a phenyl group;

with the proviso that A<sup>1</sup> may contain 1 to 3 substituents selected from the substituent groups a-1 and a-2 as defined above, and that E has 1 or 2 substituents selected from the substituent groups a-1 and a-2 defined above]

[with the proviso that (1) a compound in which E represents a group represented by the formula:

[0010]




[0011]

(wherein R<sup>A1</sup> represents a phenyl group having a halogen atom, a methoxy group, an

ethoxy group, a C<sub>1-6</sub> alkoxy carbonyl group or a carboxyl group),

(2) a compound in which E represents a group represented by the formula:

[0012]



5


[0013]

(wherein R<sup>A2</sup> represents a halogen atom or a methoxy group; R<sup>A3</sup> represents a C<sub>1-6</sub> alkyl group having a carboxyl group, a C<sub>3-8</sub> cycloalkyl group having a carboxyl group or a phenyl group having a carboxyl group),

(3) a compound in which A<sup>1</sup> represents a group represented by the formula:

10

[0014]




[0015]

(wherein R<sup>A4</sup> represents a hydrogen atom or a halogen atom; Ar represents a phenyl group which may have a substituent) and X<sup>1</sup> represents a group represented by the  
15 formula -C(=O)-NH-,

(4) a compound in which A<sup>1</sup> represents a group represented by the formula:

[0016]



[0017]


20 (wherein R<sup>A5</sup> represents a hydrogen atom, a C<sub>1-6</sub> alkyl group or a trifluoromethyl

group;  $R^{A6}$  represents a hydrogen atom or a trifluoromethyl group;  $Ar^2$  represents a phenyl group that may have a substituent) and  $X^1$  represents a group represented by the formula  $-C(=O)-NH-$  and

(5) a compound in which  $A^1$  represents a group represented by the formula:

5

[0018]



[0019]

(wherein  $R^{A7}$  represents a hydrogen atom, a halogen atom or a  $C_{1-6}$  alkyl group;  $Ar^3$  represents a phenyl group which may have a substituent) and  $X^1$  represents a group represented by the formula  $-C(=O)-NH-$  or a group represented by the formula  $-NH-C(=O)-$  are excluded];

10

[4]: the compound according to item [3], or the salt or the hydrate thereof, wherein  $A^1$  represents a 3-pyridyl group, a quinolyl group, a naphthyldinyl group, a quinoxaliny group, an imidazopyridyl group, a benzothiazolyl group, a pyrrolopyridyl group, a thienopyridyl group or a furopyridyl group (with the proviso that  $A^1$  may have 1 to 3 substituents selected from the substituent groups a-1 and a-2 defined above);

15

[5]: the compound according to item [3], or the salt or the hydrate thereof, wherein  $A^1$  represents a 3-pyridyl group (with the proviso that  $A^1$  may have 1 to 3 substituents selected from the following substituent groups c-1 and c-2);

20

<Substituent group c-1>

Substituent group c-1 represents the group consisting of: a halogen atom, an amino group, a  $C_{1-6}$  alkyl group, a  $C_{2-6}$  alkenyl group, a  $C_{2-6}$  alkynyl group, a  $C_{3-8}$  cycloalkyl group, a  $C_{6-10}$  aryl group, a 5- to 10-membered heterocyclic group, a  $C_{3-8}$  cycloalkyl  $C_{1-6}$  alkyl group, a  $C_{6-10}$  aryl  $C_{1-6}$  alkyl group, a 5- to 10-membered heterocyclic  $C_{1-6}$

alkyl group, a C<sub>1-6</sub> alkoxy group, a C<sub>2-6</sub> alkenyloxy group, a C<sub>2-6</sub> alkynyloxy group, a C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkoxy group, a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkoxy group, a 5- to 10-membered heterocyclic C<sub>1-6</sub> alkoxy group, a mono-C<sub>1-6</sub> alkylamino group, a mono-C<sub>2-6</sub> alkenylamino group, a mono-C<sub>2-6</sub> alkynylamino group, a mono-C<sub>3-8</sub> cycloalkylamino group, a mono-C<sub>6-10</sub> arylamino group, a mono-C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkylamino group, a mono-C<sub>6-10</sub> aryl C<sub>1-6</sub> alkylamino group, a mono-5- to 10-membered heterocyclic C<sub>1-6</sub> alkylamino group, a C<sub>1-6</sub> alkylcarbonyl group and a group represented by the formula -C(=N-OH)R<sup>a2</sup> (wherein R<sup>a2</sup> has the same meaning as defined above);

<Substituent group c-2>

10 Substituent group c-2 represents the group consisting of: a C<sub>1-6</sub> alkyl group, a C<sub>2-6</sub> alkenyl group, a C<sub>2-6</sub> alkynyl group, a C<sub>3-8</sub> cycloalkyl group, a C<sub>6-10</sub> aryl group, a 5- to 10-membered heterocyclic group, a C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkyl group, a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkyl group, a 5- to 10-membered heterocyclic C<sub>1-6</sub> alkyl group, a C<sub>1-6</sub> alkoxy group, a C<sub>2-6</sub> alkenyloxy group, a C<sub>2-6</sub> alkynyloxy group, a C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkoxy group, C<sub>6-10</sub> aryl C<sub>1-6</sub> alkoxy group, a 5- to 10-membered heterocyclic C<sub>1-6</sub> alkoxy group, a mono-C<sub>1-6</sub> alkylamino group, a mono-C<sub>2-6</sub> alkenylamino group, a mono-C<sub>2-6</sub> alkynylamino group, a mono-C<sub>3-8</sub> cycloalkylamino group, a mono-C<sub>6-10</sub> arylamino group, a mono-C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkylamino group, a mono-C<sub>6-10</sub> aryl C<sub>1-6</sub> alkylamino group and a mono-5- to 10-membered heterocyclic C<sub>1-6</sub> alkylamino group;

20 with the proviso that each group described in substituent group c-2 has 1 to 3 substituents selected from the following substituent group d;


<Substituent group d>

Substituent group d represents the group consisting of: a halogen atom, a hydroxyl group, a carboxyl group, an amino group, a carbamoyl group, a C<sub>1-6</sub> alkoxy group, a mono-C<sub>1-6</sub> alkylamino group, a di-C<sub>1-6</sub> alkylamino group, a mono-C<sub>6-10</sub> arylamino

group that may have one amino group or aminosulfonyl group, a N-C<sub>6-10</sub> aryl C<sub>1-6</sub> alkyl-N-C<sub>1-6</sub> alkylamino group which may have one amino group, a cyano group, a C<sub>6-10</sub> aryl group, a 5- to 10-membered heterocyclic group and a C<sub>1-6</sub> alkoxy carbonyl group.

- 5 [6]: the compound according to item [3], or the salt or the hydrate thereof, wherein A<sup>1</sup> represents a group represented by the formula:

[0020]




[0021]

- 10 [wherein R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> may be the same as or different from each other and represent a substituent selected from the substituent groups c-1 and c-2 defined above];

[7]: the compound according to item [3], or the salt or the hydrate thereof, wherein A<sup>1</sup> represents a group represented by the formula:

- 15 [0022]



[0023]


- [wherein R<sup>1</sup> and R<sup>2</sup> have the same meanings as defined above, respectively; R<sup>6</sup> and R<sup>7</sup> may be the same or different from each other and represent a hydrogen atom, a C<sub>1-6</sub> alkyl group, a C<sub>3-8</sub> cycloalkyl group or a group represented by the formula -CHR<sup>8</sup>-(CH<sub>2</sub>)<sub>n1</sub>-R<sup>9</sup> (wherein R<sup>8</sup> represents a hydrogen atom, a carboxyl group or a C<sub>1-6</sub> alkoxy carbonyl group; R<sup>9</sup> represents a hydroxyl group, a carboxyl group, a
- 20



carbamoyl group, a C<sub>3-8</sub> cycloalkyl group, a furyl group, a thienyl group, a pyrrolyl group, a pyridyl group, a triazolyl group, a tetrahydrofuryl group, a C<sub>1-6</sub> alkoxy group, a C<sub>1-6</sub> alkoxy carbonyl group, a mono-C<sub>1-6</sub> alkylamino group, a di-C<sub>1-6</sub> alkylamino group, a phenyl group which may have 1 to 3 substituents selected from the substituent group d defined above, a mono-C<sub>6-10</sub> arylamino group which may have one amino group or an N-C<sub>6-10</sub> aryl C<sub>1-6</sub> alkyl-N-C<sub>1-6</sub> alkylamino group which may have one amino group; n<sub>1</sub> represents an integer from 0 to 3));

[8]: the compound according to item [3], or the salt or the hydrate thereof, wherein A<sup>1</sup> represents a group represented by the formula:

10 [0024]




[0025]

(wherein R<sup>11</sup> represents a hydrogen atom or a group represented by the formula -CHR<sup>12</sup>-(CH<sub>2</sub>)<sub>n2</sub>-R<sup>13</sup> (wherein R<sup>12</sup> represents a hydrogen atom or a carboxyl group; R<sup>13</sup> represents a carboxyl group or a phenyl group which may have 1 to 3 substituents selected from the substituent group d defined above; n<sub>2</sub> represents an integer from 0 to 3));

[9]: the compound according to item [3], or the salt or the hydrate thereof, wherein A<sup>1</sup> represents a group represented by the formula:

20 [0026]



[0027]

(wherein R<sup>14</sup> represents a C<sub>1-6</sub> alkyl group having one C<sub>1-6</sub> alkoxy group);

[10]: the compound according to item [3], or the salt or the hydrate thereof, wherein A<sup>1</sup> represents a 6-quinolyl group, a [1,5]naphthylidin-2-yl group, a 6-quinoxaliny group, an imidazo[1,2-a]pyridin-6-yl group, a benzothiazol-6-yl group, a  
5 1*H*-pyrrolo[2,3-b]pyridin-5-yl group, a pyrrolo[3,2-b]pyridin-1-yl group, a thieno[2,3-b]pyridin-5-yl group, a thieno[3,2-b]pyridin-6-yl group or a furo[3,2-b]pyridin-6-yl group (with the proviso that A<sup>1</sup> may have 1 to 3 substituents selected from the substituent groups c-1 and c-2 defined above);

[11]: the compound according to item [3], or the salt or the hydrate thereof, wherein  
10 A<sup>1</sup> represents a 6-quinolyl group, a [1,5]naphthylidin-2-yl group, a 6-quinoxaliny group, an imidazo[1,2-a]pyridin-6-yl group, a benzothiazol-6-yl group, a pyrrolo[3,2-b]pyridin-1-yl group, a 1*H*-pyrrolo[2,3-b]pyridin-5-yl group which may have one amino group, a thieno[2,3-b]pyridin-5-yl group which may have one amino group, a thieno[3,2-b]pyridin-6-yl group which may have one amino group or  
15 furo[3,2-b]pyridin-6-yl group which may have one amino group.

[12]: the compound according to item [3], or the salt or the hydrate thereof, wherein A<sup>1</sup> represents a 6-quinolyl group;

[13]: the compound according to item [3], or the salt or the hydrate thereof, wherein A<sup>1</sup> represents a [1,5]naphthylidin-2-yl group;

20 [14]: the compound according to item [3], or the salt or the hydrate thereof, wherein A<sup>1</sup> represents an imidazo[1,2-a]pyridin-6-yl group;

[15]: the compound according to item [3], or the salt or the hydrate thereof, wherein A<sup>1</sup> represents a benzothiazol-6-yl group;

[16]: the compound according to item [3], or the salt or the hydrate thereof, wherein  
25 A<sup>1</sup> represents a 3-pyridyl group, a pyrazinyl group, a pyrimidinyl group, a quinolyl

group, an isoquinolyl group, a naphthyldinyl group, a quinoxaliny group, a cinnoliny group, a quinazoliny group, an imidazopyridyl group, a benzothiazolyl group, a benzoxazolyl group, a benzimidazolyl group, an indolyl group, a pyrrolopyridyl group, a thienopyridyl group, a furopyridyl group, a 2,3-dihydro-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl group or a benzothiadiazolyl group (with the proviso that A<sup>1</sup> may have 1 to 3 substituents selected from the substituent groups a-1 and a-2 defined above);

[17]: the compound according to item [3], or the salt or the hydrate thereof, wherein A<sup>1</sup> represents a 3-pyridyl group, a pyrazinyl group, a pyrimidinyl group, a quinolyl group, an isoquinolyl group, a naphthyldinyl group, a quinoxaliny group, a cinnoliny group, a quinazoliny group, an imidazopyridyl group, a benzothiazolyl group, a benzoxazolyl group, a benzimidazolyl group, an indolyl group, a pyrrolopyridyl group, a thienopyridyl group, a furopyridyl group, a 2,3-dihydro-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl group or a benzothiadiazolyl group (with the proviso that A<sup>1</sup> may have 1 to 3 substituents selected from the substituent groups c-1 and c-2 defined above);

[18]: the compound according to item [3], or the salt or the hydrate thereof, wherein A<sup>1</sup> represents a 3-pyridyl group, a pyrazinyl group, a pyrimidinyl group, a quinolyl group, an isoquinolyl group, a naphthyldinyl group, a quinoxaliny group, a cinnoliny group, a quinazoliny group, an imidazopyridyl group, a benzothiazolyl group, a benzoxazolyl group, a benzimidazolyl group, an indolyl group, a pyrrolopyridyl group, a thienopyridyl group, a furopyridyl group, a 2,3-dihydro-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl group or a benzothiadiazolyl group (with the proviso that A<sup>1</sup> may have 1 to 3 substituents selected from the following substituent groups c'-1 and c'-2);

<Substituent group c'-1>

Substituent group c'-1 represents the group consisting of: an amino group, a C<sub>1-6</sub> alkyl group and a mono-C<sub>1-6</sub> alkylamino group; and

<Substituent group c'-2>

Substituent group c'-2 represents the group consisting of: a C<sub>1-6</sub> alkyl group and a mono-C<sub>1-6</sub> alkylamino group;

with the proviso that each group described in substituent group c'-2 has 1 to  
5 3 substituents selected from the following substituent group d';

<Substituent group d'>

Substituent group d' represents the group consisting of: a halogen atom, a hydroxyl group, a cyano group, a carboxyl group and a C<sub>1-6</sub> alkoxy group.

[19]: the compound according to any one of items [3] to [18], or the salt or the hydrate  
10 thereof, wherein X<sup>1</sup> represents a group represented by the formula -C(=O)-NH- or a group represented by the formula -NH-C(=O)-;

[20]: the compound according to any one of items [3] to [18], or the salt or the hydrate thereof, wherein X<sup>1</sup> represents a group represented by the formula -C(=O)-NH-;

[21]: the compound according to any one of items [3] to [20], or the salt or the hydrate  
15 thereof, wherein E represents a furyl group, a thienyl group, a pyrrolyl group, a phenyl group or pyridyl group (with the proviso that E has 1 or 2 substituents selected from the substituent groups a-1 and a-2 defined above);

[22]: the compound according to any one of items [3] to [20], or the salt or the hydrate thereof, wherein E represents a furyl group, a thienyl group, a pyrrolyl group, a  
20 phenyl group or pyridyl group (with the proviso that E has 1 or 2 substituents selected from the following substituent groups e-1 and e-2);

<Substituent group e-1>

Substituent group e-1 represents the group consisting of: a halogen atom, a hydroxyl group, a C<sub>1-6</sub> alkyl group, a C<sub>2-6</sub> alkenyl group, a C<sub>2-6</sub> alkynyl group, a C<sub>6-10</sub> aryl group,  
25 a C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkyl group, a C<sub>3-8</sub> cycloalkylidene C<sub>1-6</sub> alkyl group, a C<sub>6-10</sub> aryl

C<sub>1-6</sub> alkyl group, 5- to 10-membered heterocyclic C<sub>1-6</sub> alkyl group, a C<sub>1-6</sub> alkoxy group, a C<sub>2-6</sub> alkenyloxy group, a C<sub>2-6</sub> alkynyloxy group, a C<sub>6-10</sub> aryloxy group, a C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkoxy group, a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkoxy group, a 5- to 10-membered heterocyclic C<sub>1-6</sub> alkoxy group, a C<sub>6-10</sub> arylthio group, a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkylthio group, a mono-C<sub>6-10</sub> arylamino group, a mono-C<sub>6-10</sub> aryl C<sub>1-6</sub> alkylamino group, a N-C<sub>6-10</sub> aryl-N-C<sub>1-6</sub> alkylamino group, a N-C<sub>6-10</sub> aryl C<sub>1-6</sub> alkyl-N-C<sub>1-6</sub> alkylamino group, a C<sub>6-10</sub> aryloxy C<sub>1-6</sub> alkyl group and a 5- to 10-membered heterocycle oxy C<sub>1-6</sub> alkyl group;

<Substituent group e-2>

Substituent group e-2 represents the group consisting of: a C<sub>1-6</sub> alkyl group, a C<sub>2-6</sub> alkenyl group, a C<sub>2-6</sub> alkynyl group, a C<sub>6-10</sub> aryl group, a C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkyl group, a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkyl group, a 5- to 10-membered heterocyclic C<sub>1-6</sub> alkyl group, a C<sub>1-6</sub> alkoxy group, a C<sub>2-6</sub> alkenyloxy group, a C<sub>2-6</sub> alkynyloxy group, a C<sub>6-10</sub> aryloxy group, a C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkoxy group, a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkoxy group, 5- to 10-membered heterocycle-C<sub>1-6</sub> alkoxy group, a C<sub>6-10</sub> arylthio group, a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkylthio group, a mono-C<sub>6-10</sub> arylamino group, a mono-C<sub>6-10</sub> aryl C<sub>1-6</sub> alkylamino group, a N-C<sub>6-10</sub> aryl-N-C<sub>1-6</sub> alkylamino group, a N-C<sub>6-10</sub> aryl C<sub>1-6</sub> alkyl-N-C<sub>1-6</sub> alkylamino group, a C<sub>6-10</sub> aryloxy C<sub>1-6</sub> alkyl group and a 5- to 10-membered heterocycle oxy C<sub>1-6</sub> alkyl group;

with the proviso that each group described in substituent group e-2 has 1 to 3 substituents selected from the following substituent group f;

<Substituent group f>

Substituent group f represents the group consisting of: a halogen atom, a hydroxyl group, a cyano group, an amino group, a nitro group, a C<sub>3-8</sub> cycloalkyl group, a C<sub>1-6</sub> alkoxy group, a C<sub>6-10</sub> aryloxy group, a 5- to 10-membered heterocycle oxy group, a C<sub>1-6</sub> alkylcarbonyl group, a C<sub>1-6</sub> alkoxy carbonyl group, a C<sub>1-6</sub> alkylsulfonyl group, a

mono-C<sub>6-10</sub> arylamino group, a trifluoromethyl group, a trifluoromethoxy group and a C<sub>1-6</sub> alkyl group;

[23]: the compound according to any one of items [3] to [20], or the salt or the hydrate thereof, wherein E represents a furyl group, a thienyl group, a pyrrolyl group, a phenyl group or a pyridyl group (with the proviso that E has one substituent selected from the following substituent groups g-1 and g-2);

<Substituent group g-1>

Substituent group g-1 represents the group consisting of: a C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkyl group, a phenyl C<sub>1-6</sub> alkyl group, a furyl C<sub>1-6</sub> alkyl group, a thienyl C<sub>1-6</sub> alkyl group, a benzofuryl C<sub>1-6</sub> alkyl group, a benzothienyl C<sub>1-6</sub> alkyl group, a C<sub>1-6</sub> alkoxy group, a phenoxy group, a C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkoxy group, a phenyl C<sub>1-6</sub> alkoxy group, a furyl C<sub>1-6</sub> alkoxy group, a thienyl C<sub>1-6</sub> alkoxy group, a pyridyl C<sub>1-6</sub> alkoxy group, a phenoxy C<sub>1-6</sub> alkyl group and a pyridyloxy C<sub>1-6</sub> alkyl group;

<Substituent group g-2>

Substituent group g-2 represents the group consisting of: a C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkyl group, a phenyl C<sub>1-6</sub> alkyl group, a furyl C<sub>1-6</sub> alkyl group, a thienyl C<sub>1-6</sub> alkyl group, a benzofuryl C<sub>1-6</sub> alkyl group, a benzothienyl C<sub>1-6</sub> alkyl group, a C<sub>1-6</sub> alkoxy group, a phenoxy group, a C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkoxy group, a phenyl C<sub>1-6</sub> alkoxy group, a furyl C<sub>1-6</sub> alkoxy group, a thienyl C<sub>1-6</sub> alkoxy group, a pyridyl C<sub>1-6</sub> alkoxy group, a phenoxy C<sub>1-6</sub> alkyl group and a pyridyloxy C<sub>1-6</sub> alkyl group;

with the proviso that each group described in substituent group g-2 has 1 to 3 substituents selected from the following substituent group h;


<Substituent group h>

Substituent group h represents the group consisting of: a halogen atom, a hydroxyl group, a cyano group and a C<sub>1-6</sub> alkyl group;

[24]: the compound according to any one of items [3] to [20], or the salt or the hydrate thereof, wherein E represents a 2-furyl group, a 2-thienyl group, a 3-pyrrolyl group, a phenyl group, a 2-pyridyl group or 3-pyridyl group (with the proviso that E has one substituent selected from the substituent groups g-1 and g-2 defined above);

- 5 [25]: the compound according to item [3], or the salt or the hydrate thereof, wherein  $X^1$  represents a group represented by the formula  $-C(=O)-NH-$ , and  $A^1$  represents a group represented by the formula:

[0028]




10

[0029]


(wherein  $R^1$ ,  $R^2$  and  $R^3$  have the same meanings as defined above, respectively), and E represents a 2-furyl group, a 2-thienyl group, a 3-pyrrolyl group, a phenyl group, a 2-pyridyl group or a 3-pyridyl group (with the proviso that E has one substituent selected from the substituent group g-1 or g-2 defined above).

- 15 [26]: the compound according to item [25], or the salt or the hydrate thereof, wherein  $A^1$  represents a group represented by the formula:

[0030]



or




[0031]

- 20 (wherein  $R^1$ ,  $R^2$ ,  $R^6$  and  $R^7$  have the same meanings as defined above, respectively);

[27]: the compound according to item [25], or the salt or the hydrate thereof, wherein  $A^1$  represents a group represented by the formula:

[0032]




[0033]

(wherein R<sup>11</sup> has the same meaning as defined above);

- 5 [28]: the compound according to item [25], or the salt or the hydrate thereof, wherein A<sup>1</sup> represents a group represented by the formula:

[0034]



[0035]

- 10 (wherein R<sup>14</sup> has the same meaning as defined above);

- [29]: the compound according to item [3], or the salt or the hydrate thereof, wherein X<sup>1</sup> represents a group represented by the formula -C(=O)-NH-, A<sup>1</sup> represents a 6-quinolyl group, a [1,5]naphthylidin-2-yl group, a 6-quinoxaliny group, an imidazo[1,2-a]pyridin-6-yl group, a benzothiazol-6-yl group, a pyrrolo[3,2-b]pyridin-1-yl group, a 1*H*-pyrrolo[2,3-b]pyridin-5-yl group which may have one amino group, a thieno[2,3-b]pyridin-5-yl group which may have one amino group, a thieno[3,2-b]pyridin-6-yl group which may have one amino group or a furo[3,2-b]pyridin-6-yl group which may have one amino group, and E represents a 2-furyl group, a 2-thienyl group, a 3-pyrrolyl group, a phenyl group or a 2-pyridyl group (with the proviso that E has a substituent selected from the substituent group g-1 or g-2 defined above);
- 15
- 20

[30]: the compound according to item [29], or the salt or the hydrate thereof, wherein



A<sup>1</sup> represents a 6-quinolyl group;

[31]: the compound according to item [29], or the salt or the hydrate thereof, wherein

A<sup>1</sup> represents a [1,5]naphthylidin-2-yl group;

[32]: the compound according to item [29], or the salt or the hydrate thereof, wherein


5 A<sup>1</sup> represents an imidazo[1,2-a]pyridin-6-yl group;

[33]: the compound according to item [29], or the salt or the hydrate thereof, wherein

A<sup>1</sup> represents a benzothiazol-6-yl group;

[34]: A compound represented by the formula (I-b), or a salt or a hydrate thereof:

[0036]



[0037]

[wherein A<sup>2</sup> represents a 6-quinolyl group, a 4-quinazolinyl group or a pyrido[2,3-d]pyrimidin-4-yl group which may have an amino group;

15 X<sup>2</sup> represents a group represented by the formula -O-CH<sub>2</sub>-, a group represented by the formula -S-CH<sub>2</sub>-, a group represented by the formula -C(=O)-CH<sub>2</sub>-, a group represented by the formula -NH-CH<sub>2</sub>- or a group represented by the formula -CH<sub>2</sub>-NH-;

R<sup>10</sup> represents a C<sub>1-6</sub> alkyl group, a C<sub>6-10</sub> aryloxy group or a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkoxy group];

20 [35]: the compound according to item [34], or the salt or the hydrate thereof, wherein X<sup>2</sup> represents a group represented by the formula -NH-CH<sub>2</sub>- or a group represented by the formula -CH<sub>2</sub>-NH-;

[36]: a pharmaceutical composition comprising the compound according to item [3] or [34], or the salt or the hydrate thereof;

[37]: an antifungal agent comprising, as an active ingredient, the compound according to item [3] or [34], or the salt or the hydrate thereof;

[38]: a method for prevention or treatment of fungal infection comprising administering a pharmacologically effective amount of the compound according to  
5 item [3] or [34], or the salt or the hydrate thereof;

[39]: a use of the compound according to item [3] or [34], or the salt or the hydrate thereof, for manufacture of an antifungal agent;

[40]: a method for prevention or treatment of fungal infection comprising administering a pharmacologically effective amount of the antifungal agent according  
10 to item [1].

[Best Mode for Carrying Out the Invention]

[0038]

The present invention is explained below in more detail by reference to the symbols and the terms used herein being defined and the following examples.

15        Herein, a structural formula of a compound sometimes represents a certain isomer for convenience of description. However, compounds according to the present invention may include all possible isomers, such as structurally possible geometric isomers, optical isomers generated due to the presence of asymmetric carbons, stereoisomers, tautomers, and mixtures of isomers, and are not limited to  
20 formulae being used for the convenience of description, and may be either of two isomers of a mixture of both isomers. Thus, the compounds according to the present invention may be either optically active compounds having an asymmetric carbon atom in their molecules or their racemates, and are not restricted to either of them but include both. Furthermore, the compounds according to the present  
25 invention may exhibit crystalline polymorphism, but likewise are not restricted to any

one of these, but may be in any one of these crystal forms or exist as a mixture of two or more crystal forms. The compounds according to the present invention also include both anhydrous and hydrated forms.

In addition, compounds resulting from compounds according to the present invention that undergo *in vivo* metabolism such as oxidation, reduction, hydrolysis and conjugation (so-called metabolites), and compounds that undergo *in vivo* metabolism such as oxidation, reduction, hydrolysis and conjugation and generates compounds according to the present invention (so-called prodrugs) are also encompassed by the scope of the present invention.

The term "C<sub>1-6</sub> alkyl group" used in the present specification means a linear or branched chain alkyl group containing 1 to 6 carbon atoms, which is a monovalent group derived by removal of any one of the hydrogen atoms from an aliphatic hydrocarbon containing 1 to 6 carbons. Specifically, examples of "C<sub>1-6</sub> alkyl group" include a methyl group, an ethyl group, a *n*-propyl group, an *iso*-propyl group, a *n*-butyl group, an *iso*-butyl group, a *sec*-butyl group, a *tert*-butyl group, a *n*-pentyl group, an *iso*-pentyl group, a *sec*-pentyl group, a neopentyl group, a 1-methylbutyl group, a 2-methylbutyl group, a 1,1-dimethylpropyl group, a 1,2-dimethylpropyl group, a *n*-hexyl group, an *iso*-hexyl group, a 1-methylpentyl group, a 2-methylpentyl group, a 3-methylpentyl group, a 1,1-dimethylbutyl group, a 1,2-dimethylbutyl group, a 2,2-dimethylbutyl group, a 1,3-dimethylbutyl group, a 2,3-dimethylbutyl group, a 3,3-dimethylbutyl group, a 1-ethylbutyl group, a 2-ethylbutyl group, a 1,1,2-trimethylpropyl group, a 1,2,2-trimethylpropyl group, a 1-ethyl-1-methylpropyl group, a 1-ethyl-2-methylpropyl group or the like, and preferably, a methyl group, an ethyl group, a *n*-propyl group, an *iso*-propyl group, a *n*-butyl group, an *iso*-butyl group, a *sec*-butyl group, a *tert*-butyl group or the like.

The term "C<sub>2-6</sub> alkenyl group" used in the present specification means a linear or branched chain alkenyl group containing 2 to 6 carbon atoms, which may contain 1 to 2 double-bonds. Specifically, examples of "C<sub>2-6</sub> alkenyl group" include an ethenyl group, a 1-propenyl group, a 2-propenyl group, a 1-butenyl group, a 2-butenyl group, a 3-butenyl group, a 2-methyl-1-propenyl group, a pentenyl group, a 3-methyl-2-butenyl group, a hexenyl group, a hexane dienyl group or the like, and preferably an ethenyl group, a 1-propenyl group, a 2-propenyl group, a 1-butenyl group, a 2-butenyl group, a 3-butenyl group, a 2-methyl-1-propenyl group, a 3-methyl-2-butenyl group or the like.

The term "C<sub>2-6</sub> alkynyl group" used in the present specification means a linear or branched chain alkynyl group containing 2 to 6 carbon atoms, which may contain 1 to 2 triple-bonds. Specifically, examples of "C<sub>2-6</sub> alkynyl group" include an ethynyl group, a 1-propynyl group, a 2-propynyl group, a 1-butynyl group, a 2-butynyl group, a 3-butynyl group, a pentynyl group, a hexynyl group, a hexane diynyl group or the like, and preferably, an ethynyl group, a 1-propynyl group, a 2-propynyl group, a 1-butynyl group, a 2-butynyl group, a 3-butynyl group or the like.

The term "C<sub>3-8</sub> cycloalkyl group" used in the present specification means a cyclic aliphatic hydrocarbon group containing 3 to 8 carbon atoms. Specifically, examples of "C<sub>3-8</sub> cycloalkyl group" include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group or the like, and preferably, a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group or the like.

The term "C<sub>1-6</sub> alkoxy group" used in the present specification means a group in which an oxygen atom is bonded to terminus of the "C<sub>1-6</sub> alkyl group" defined above. Specifically, examples of "C<sub>1-6</sub> alkoxy group" include a methoxy group, an

ethoxy group, a *n*-propoxy group, an *iso*-propoxy group, a *n*-butoxy group, an *iso*-butoxy group, a *sec*-butoxy group, a *tert*-butoxy group, a *n*-pentyloxy group, an *iso*-pentyloxy group, a *sec*-pentyloxy group, a neopentyloxy group, a 1-methylbutoxy group, a 2-methylbutoxy group, a 1,1-dimethylpropoxy group, a 1,2-dimethylpropoxy group, a *n*-hexyloxy group, an *iso*-hexyloxy group, a 1-methylpentyloxy group, a 2-methylpentyloxy group, a 3-methylpentyloxy group, a 1,1-dimethylbutoxy group, a 1,2-dimethylbutoxy group, a 2,2-dimethylbutoxy group, a 1,3-dimethylbutoxy group, a 2,3-dimethylbutoxy group, a 3,3-dimethylbutoxy group, a 1-ethylbutoxy group, a 2-ethylbutoxy group, a 1,1,2-trimethylpropoxy group, a 1,2,2-trimethylpropoxy group, a 1-ethyl-1-methylpropoxy group, a 1-ethyl-2-methylpropoxy group or the like and preferably, a methoxy group, an ethoxy group, a *n*-propoxy group, an *iso*-propoxy group, a *n*-butoxy group, an *iso*-butoxy group, a *sec*-butoxy group, a *tert*-butoxy group and the like.

The term "C<sub>1-6</sub> alkylthio group" used in the present specification means a group in which a sulfur atom is bonded to terminus of the "C<sub>1-6</sub> alkyl group" defined above. Specifically, examples of "C<sub>1-6</sub> alkylthio group" include a methylthio group, an ethylthio group, a *n*-propylthio group, an *iso*-propylthio group, a *n*-butylthio group, an *iso*-butylthio group, a *sec*-butylthio group, a *tert*-butylthio group, a *n*-pentylthio group, an *iso*-pentylthio group, a *sec*-pentylthio group, a neopentylthio group, a 1-methylbutylthio group, a 2-methylbutylthio group, a 1,1-dimethylpropylthio group, a 1,2-dimethylpropylthio group, a *n*-hexylthio group, an *iso*-hexylthio group, a 1-methylpentylthio group, a 2-methylpentylthio group, a 3-methylpentylthio group, a 1,1-dimethylbutylthio group, a 1,2-dimethylbutylthio group, a 2,2-dimethylbutylthio group, a 1,3-dimethylbutylthio group, a 2,3-dimethylbutylthio group, a 3,3-dimethylbutylthio group, a 1-ethylbutylthio group, a 2-ethylbutylthio group, a

1,1,2-trimethylpropylthio group, a 1,2,2-trimethylpropylthio group, a 1-ethyl-1-methylpropylthio group, a 1-ethyl-2-methylpropylthio group or the like, and preferably, a methylthio group, an ethylthio group, a *n*-propylthio group, an *iso*-propylthio group, a *n*-butylthio group, an *iso*-butylthio group, a *sec*-butylthio group, a *tert*-butylthio group or the like.

The term “C<sub>1-6</sub> alkylcarbonyl group” used in the present specification means a group in which a carbonyl group is bonded to terminus of the “C<sub>1-6</sub> alkyl group” defined above. Specifically, examples of “C<sub>1-6</sub> alkylcarbonyl group” include a methylcarbonyl group, an ethylcarbonyl group, a *n*-propylcarbonyl group, an *iso*-propylcarbonyl group or the like.

The term “C<sub>1-6</sub> alkoxy carbonyl group” used in the present specification means a group in which a carbonyl group is bonded to terminus of the “C<sub>1-6</sub> alkoxy group” defined above. Specifically, examples of “C<sub>1-6</sub> alkoxy carbonyl group” include a methoxycarbonyl group, an ethoxycarbonyl group, a *n*-propoxycarbonyl group, an *iso*-propoxycarbonyl group or the like.

The term “C<sub>1-6</sub> alkylsulfonyl group” used in the present specification means a group in which a sulfonyl group is bonded to terminus of the “C<sub>1-6</sub> alkyl group” defined above. Specifically, examples of “C<sub>1-6</sub> alkylsulfonyl group” include a methylsulfonyl group, an ethylsulfonyl group, a *n*-propylsulfonyl group, an *iso*-propylsulfonyl group or the like.

The term “C<sub>2-6</sub> alkenyloxy group” used in the present specification means a group in which an oxygen atom is bonded to terminus of the “C<sub>2-6</sub> alkenyl group” defined above. Specifically, examples of “C<sub>2-6</sub> alkenyloxy group” include an ethenyloxy group, a 1-propenyloxy group, a 2-propenyloxy group, a 1-butenyloxy group, a 2-butenyloxy group, a 3-butenyloxy group, a 2-methyl-1-propenyloxy group,

a pentenyloxy group, a 3-methyl-2-butenyloxy group, a hexenyloxy group, a hexane dienyloxy group or the like, and preferably, an ethenyloxy group, a 1-propenyloxy group, a 2-propenyloxy group, a 1-butenyloxy group, a 2-butenyloxy group, a 3-butenyloxy group, a 2-methyl-1-propenyloxy group, a 3-methyl-2-butenyloxy group or the like.

The term "C<sub>2-6</sub> alkenylthio group" used in the present specification means a group in which a sulfur atom is bonded to terminus of the "C<sub>2-6</sub> alkenyl group" defined above. Specifically, examples of "C<sub>2-6</sub> alkenylthio group" include an ethenylthio group, a 1-propenylthio group, a 2-propenylthio group, a 1-butenylthio group, a 2-butenylthio group, a 3-butenylthio group, a 2-methyl-1-propenylthio group, a pentenylthio group, a 3-methyl-2-butenylthio group, a hexenylthio group, a hexane dienylthio group or the like, and preferably, an ethenylthio group, a 1-propenylthio group, a 2-propenylthio group, a 1-butenylthio group, a 2-butenylthio group, a 3-butenylthio group, a 2-methyl-1-propenylthio group, a 3-methyl-2-butenylthio group or the like.

The term "C<sub>2-6</sub> alkynyloxy group" used in the present specification means a group in which an oxygen atom is bonded to terminus of the "C<sub>2-6</sub> alkynyl group" defined above. Specifically, examples of "C<sub>2-6</sub> alkynyloxy group" include an ethynyloxy group, a 1-propynyloxy group, a 2-propynyloxy group, a 1-butyloxy group, a 2-butyloxy group, a 3-butyloxy group, a pentynyloxy group, a hexynyloxy group, a hexane diynyloxy group or the like, and preferably, an ethynyloxy group, a 1-propynyloxy group, a 2-propynyloxy group, a 1-butyloxy group, a 2-butyloxy group, a 3-butyloxy group or the like.

The term "C<sub>2-6</sub> alkynylthio group" used in the present specification means a group in which a sulfur atom is bonded to terminus of the "C<sub>2-6</sub> alkynyl group" defined

above. Specifically, examples of "C<sub>2-6</sub> alkynylthio group" include an ethynylthio group, a 1-propynylthio group, a 2-propynylthio group, a 1-butynylthio group, a 2-butynylthio group, a 3-butynylthio group, a pentynylthio group, a hexynylthio group, a hexane diynylthio group or the like, and preferably, an ethynylthio group, a 1-propynylthio group, a 2-propynylthio group, a 1-butynylthio group, a 2-butynylthio group, a 3-butynylthio group or the like.

The term "C<sub>3-8</sub> cycloalkoxy group" used in the present specification means a group in which an oxygen atom is bonded to terminus of the "C<sub>3-8</sub> cycloalkyl group" defined above. Specifically, examples of "C<sub>3-8</sub> cycloalkoxy group" include a cyclopropoxy group, a cyclobutoxy group, a cyclopentyloxy group, a cyclohexyloxy group, a cycloheptyloxy group, a cyclooctyloxy group or the like, and preferably, a cyclopropoxy group, a cyclobutoxy group, a cyclopentyloxy group, a cyclohexyloxy group or the like.

The term "C<sub>3-8</sub> cycloalkylthio group" used in the present specification means a group in which a sulfur atom is bonded to terminus of the "C<sub>3-8</sub> cycloalkyl group" defined above. Specifically, examples of "C<sub>3-8</sub> cycloalkylthio group" include a cyclopropylthio group, a cyclobutylthio group, a cyclopentylthio group, a cyclohexylthio group, a cycloheptylthio group, a cyclooctylthio group or the like, and preferably, a cyclopropylthio group, a cyclobutylthio group, a cyclopentylthio group, a cyclohexylthio group or the like.

The term "C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkyl group" used in the present specification means a group in which any hydrogen atom within the "C<sub>1-6</sub> alkyl group" defined above is substituted by the "C<sub>3-8</sub> cycloalkyl group" defined above. Specifically, examples of "C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkyl group" include a cyclopropylmethyl group, a cyclobutylmethyl group, a cyclopentylmethyl group, a cyclohexylmethyl group, a



cyclopropylethyl group, a cyclobutylethyl group, a cyclopentylethyl group, a cyclohexylethyl group or the like.

The term "C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkoxy group" used in the present specification means a group in which any hydrogen atom within the "C<sub>1-6</sub> alkoxy group" defined above is substituted by the "C<sub>3-8</sub> cycloalkyl group" defined above. Specifically, examples of "C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkoxy group" include a cyclopropylmethoxy group, a cyclobutylmethoxy group, a cyclopentylmethoxy group, a cyclohexylmethoxy group, a cyclopropylethoxy group, a cyclobutylethoxy group, a cyclopentylethoxy group, a cyclohexylethoxy group or the like.

The term "C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkylthio group" used in the present specification means a group in which any hydrogen atom within the "C<sub>1-6</sub> alkylthio group" defined above is substituted by the "C<sub>3-8</sub> cycloalkyl group" defined above. Specifically, examples of "C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkylthio group" include a cyclopropylmethylthio group, a cyclobutylmethylthio group, a cyclopentylmethylthio group, a cyclohexylmethylthio group, a cyclopropylethylthio group, a cyclobutylethylthio group, a cyclopentylethylthio group, a cyclohexylethylthio group or the like.

The term "C<sub>3-8</sub> cycloalkylidene C<sub>1-6</sub> alkyl group" used in the present specification means a group in which any carbon atom of a saturated cyclic aliphatic hydrocarbon containing 3 to 8 carbon atoms and the "C<sub>1-6</sub> alkyl group" defined above are bonded through a double-bond. Specifically, examples of "C<sub>3-8</sub> cycloalkylidene C<sub>1-6</sub> alkyl group" include a cyclopropylidene methyl group, a cyclobutylidene methyl group, a cyclopentylidene methyl group, a cyclohexylidene methyl group, a cyclopropylidene ethyl group, a cyclobutylidene ethyl group, a cyclopentylidene ethyl group, a cyclohexylidene ethyl group or the like.

The term "C<sub>6-10</sub> aryl group" used in the present specification means an aromatic cyclic hydrocarbon group containing 6 to 10 carbon atoms. Specifically, examples of "C<sub>6-10</sub> aryl group" include a phenyl group, a 1-naphthyl group, a 2-naphthyl group, an indenyl group, an azulenyl group, a heptalenyl group or the like, and preferably, a phenyl group, a 1-naphthyl group, a 2-naphthyl group or the like.

The term "C<sub>6-10</sub> aryloxy group" used in the present specification means a group in which an oxygen atom is bonded to terminus of the "C<sub>6-10</sub> aryl group" defined above. Specifically, examples of "C<sub>6-10</sub> aryloxy group" include a phenoxy group, a 1-naphthyloxy group, a 2-naphthyloxy group, an indenyloxy group, an azulenyloxy group, a heptalenyloxy group or the like, and preferably, a phenoxy group, a 1-naphthyloxy group, a 2-naphthyloxy group or the like.

The term "C<sub>6-10</sub> arylthio group" used in the present specification means a group in which a sulphur atom is bonded to terminus of the "C<sub>6-10</sub> aryl group" defined above. Specifically, examples of "C<sub>6-10</sub> arylthio group" include a phenylthio group, a 1-naphthylthio group, a 2-naphthylthio group, an indenylthio group, an azulenylthio group, a heptalenylthio group or the like, and preferably, a phenylthio group, a 1-naphthylthio group, a 2-naphthylthio group or the like.

The term "C<sub>6-10</sub> aryl C<sub>1-6</sub> alkyl group" used in the present specification means a group in which any hydrogen atom within the "C<sub>1-6</sub> alkyl group" defined above is substituted by the "C<sub>6-10</sub> aryl group" defined above. Specifically, examples of "C<sub>6-10</sub> aryl C<sub>1-6</sub> alkyl group" include a benzyl group, a 1-naphthylmethyl group, a 2-naphthylmethyl group, a phenethyl group, a 1-naphthylethyl group, a 2-naphthylethyl group, a 3-phenyl-1-propyl group or the like.

The term "phenyl C<sub>1-6</sub> alkyl group" used in the present specification means a group in which any hydrogen atom within the "C<sub>1-6</sub> alkyl group" defined above is

substituted by a phenyl group. Specifically, examples of "phenyl C<sub>1-6</sub> alkyl group" include a benzyl group, a phenethyl group, a 3-phenyl-1-propyl group or the like.

The term "C<sub>6-10</sub> aryl C<sub>1-6</sub> alkoxy group" used in the present specification means a group in which any hydrogen atom within the "C<sub>1-6</sub> alkoxy group" defined  
5 above is substituted by the "C<sub>6-10</sub> aryl group" defined above. Specifically, examples of "C<sub>6-10</sub> aryl C<sub>1-6</sub> alkoxy group" include a benzyloxy group, a 1-naphthylmethoxy group, a 2-naphthylmethoxy group, a phenethyloxy group, a 1-naphthylethoxy group, a 2-naphthylethoxy group, a 3-phenyl-1-propoxy group or the like.

The term "phenyl C<sub>1-6</sub> alkoxy group" used in the present specification means  
10 a group in which any hydrogen atom within the "C<sub>1-6</sub> alkoxy group" defined above is substituted by a phenyl group. Specifically, examples of "phenyl C<sub>1-6</sub> alkoxy group" include a benzyloxy group, a phenethyloxy group, a 3-phenyl-1-propoxy group or the like.

The term "C<sub>6-10</sub> aryl C<sub>1-6</sub> alkylthio group" used in the present specification  
15 means a group in which any hydrogen atom within the "C<sub>1-6</sub> alkylthio group" defined above is substituted by the "C<sub>6-10</sub> aryl group" defined above. Specifically, examples of "C<sub>6-10</sub> aryl C<sub>1-6</sub> alkylthio group" include a benzylthio group, a phenethylthio group, a 3-phenyl-1-propylthio group or the like.

The term "mono-C<sub>1-6</sub> alkylamino group" used in the present specification  
20 means a group in which one hydrogen atom within an amino group is substituted by the "C<sub>1-6</sub> alkyl group" defined above. Specifically, examples of "mono-C<sub>1-6</sub> alkylamino group" include a methylamino group, an ethylamino group, a *n*-propylamino group, an *iso*-propylamino group, a *n*-butylamino group, an *iso*-butylamino group, a *sec*-butylamino group, a *tert*-butylamino group, a  
25 *n*-pentylamino group, an *iso*-pentylamino group, a *sec*-pentylamino group, a

neopentylamino group, a 1-methylbutylamino group, a 2-methylbutylamino group, a 1,1-dimethylpropylamino group, a 1,2-dimethylpropylamino group, a *n*-hexylamino group, an *iso*-hexylamino group, a 1-methylpentylamino group, a 2-methylpentylamino group, a 3-methylpentylamino group, a 1,1-dimethylbutylamino group, a 1,2-dimethylbutylamino group, a 2,2-dimethylbutylamino group, a 1,3-dimethylbutylamino group, a 2,3-dimethylbutylamino group, a 3,3-dimethylbutylamino group, a 1-ethylbutylamino group, a 2-ethylbutylamino group, a 1,1,2-trimethylpropylamino group, a 1,2,2-trimethylpropylamino group, a 1-ethyl-1-methylpropylamino group, a 1-ethyl-2-methylpropylamino group or the like, and preferably, a methylamino group, an ethylamino group, a *n*-propylamino group, an *iso*-propylamino group, a *n*-butylamino group, an *iso*-butylamino group, a *sec*-butylamino group, a *tert*-butylamino group or the like.

The term "mono-C<sub>2-6</sub> alkenylamino group" used in the present specification means a group in which one hydrogen atom within an amino group is substituted by the "C<sub>2-6</sub> alkenyl group" defined above. Specifically, examples of "mono-C<sub>2-6</sub> alkenylamino group" include an ethenylamino group, a 1-propenylamino group, a 2-propenylamino group, a 1-butenylamino group, a 2-butenylamino group, a 3-butenylamino group, a 2-methyl-1-propenylamino group, a pentenylamino group, a 3-methyl-2-butenylamino group, a hexenylamino group, a hexane dienylamino group or the like, and preferably, an ethenylamino group, a 1-propenylamino group, a 2-propenylamino group, a 1-butenylamino group, a 2-butenylamino group, a 3-butenylamino group, a 2-methyl-1-propenylamino group, a 3-methyl-2-butenylamino group or the like.

The term "mono-C<sub>2-6</sub> alkynylamino group" used in the present specification means a group in which one hydrogen atom within an amino group is substituted by

the "C<sub>2-6</sub> alkynyl group" defined above. Specifically, examples of "mono-C<sub>2-6</sub> alkynylamino group" include an ethynylamino group, a 1-propynylamino group, a 2-propynylamino group, a 1-butynylamino group, a 2-butynylamino group, a 3-butynylamino group, a pentynylamino group, a hexynylamino group, a hexane  
5 diynylamino group or the like, and preferably, an ethynylamino group, a 1-propynylamino group, a 2-propynylamino group, a 1-butynylamino group, a 2-butynylamino group, a 3-butynylamino group or the like.

The term "mono-C<sub>3-8</sub> cycloalkylamino group" used in the present specification means a group in which one hydrogen atom within an amino group is substituted by  
10 the "C<sub>3-8</sub> cycloalkyl group" defined above. Specifically, examples of "mono-C<sub>3-8</sub> cycloalkylamino group" include a cyclopropylamino group, a cyclobutylamino group, a cyclopentylamino group, a cyclohexylamino group, a cycloheptylamino group, a cyclooctylamino group or the like, and preferably, a cyclopropylamino group, a cyclobutylamino group, a cyclopentylamino group, a cyclohexylamino group or the  
15 like.

The term "mono-C<sub>6-10</sub> arylamino group" used in the present specification means a group in which one hydrogen atom within an amino group is substituted by the "C<sub>6-10</sub> aryl group" defined above. Specifically, examples of "mono-C<sub>6-10</sub> arylamino group" include a phenylamino group, a 1-naphthylamino group, a  
20 2-naphthylamino group, an indenylamino group, an azulenylamino group, a heptalenylamino group or the like, and preferably, a phenylamino group, a 1-naphthylamino group, a 2-naphthylamino group or the like.

The term "mono-C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkylamino group" used in the present specification means a group in which one hydrogen atom within an amino group is  
25 substituted by the "C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkyl group" defined above. Specifically,

examples of "mono-C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkylamino group" include a cyclopropylmethylamino group, a cyclobutylmethylamino group, a cyclopentylmethylamino group, a cyclohexylmethylamino group, a cyclopropylethylamino group, a cyclobutylethylamino group, a cyclopentylethylamino group, a cyclohexylethylamino group or the like.

The term "mono-C<sub>6-10</sub> aryl C<sub>1-6</sub> alkylamino group" used in the present specification means a group in which one hydrogen atom within an amino group is substituted by the "C<sub>6-10</sub> aryl C<sub>1-6</sub> alkyl group" defined above. Specifically, examples of "mono-C<sub>6-10</sub> aryl C<sub>1-6</sub> alkylamino group" include a benzyl amino group, a 1-naphthylmethylamino group, a 2-naphthylmethylamino group, a phenethylamino group, a 1-naphthylethylamino group, a 2-naphthylethylamino group or the like.

The term "di-C<sub>1-6</sub> alkylamino group" used in the present specification means a group in which two hydrogen atoms within an amino group are substituted by the identical to or different from the "C<sub>1-6</sub> alkyl groups" defined above. Specifically, examples of "di-C<sub>1-6</sub> alkylamino group" include a N,N-dimethylamino group, a N,N-diethylamino group, a N,N-di-*n*-propylamino group, a N,N-di-*iso*-propylamino group, a N,N-di-*n*-butylamino group, a N,N-di-*iso*-butylamino group, a N,N-di-*sec*-butylamino group, a N,N-di-*tert*-butylamino group, a N-ethyl-N-methylamino group, a N-*n*-propyl-N-methylamino group, a N-*iso*-propyl-N-methylamino group, a N-*n*-butyl-N-methylamino group, a N-*iso*-butyl-N-methylamino group, a N-*sec*-butyl-N-methylamino group, a N-*tert*-butyl-N-methylamino group or the like, and preferably, a N,N-dimethylamino group, a N,N-diethylamino group, a N-ethyl-N-methylamino group or the like.

The term "N-C<sub>2-6</sub> alkenyl-N-C<sub>1-6</sub> alkylamino group" used in the present specification means a group in which one of the hydrogen atoms within an amino

group is substituted by the "C<sub>2-6</sub> alkenyl group" defined above, and the other hydrogen atom is substituted by the "C<sub>1-6</sub> alkyl group" defined above. Specifically, examples of "N-C<sub>2-6</sub> alkenyl-N-C<sub>1-6</sub> alkylamino group" include a N-ethenyl-N-methylamino group, a N-1-propenyl-N-methylamino group, a  
 5 N-2-propenyl-N-methylamino group, a N-1-butenyl-N-methylamino group, a N-2-butenyl-N-methylamino group, a N-3-butenyl-N-methylamino group, a N-2-methyl-1-propenyl-N-methylamino group, a N-pentenyl-N-methylamino group, a N-3-methyl-2-butenyl-N-methylamino group, a N-hexenyl-N-methylamino group, a N-hexanedi-enyl-N-methylamino group or the like, and preferably, a  
 10 N-ethenyl-N-methylamino group, a N-1-propenyl-N-methylamino group, a N-2-propenyl-N-methylamino group, a N-1-butenyl-N-methylamino group, a N-2-butenyl-N-methylamino group, a N-3-butenyl-N-methylamino group, a N-2-methyl-1-propenyl-N-methylamino group, a N-3-methyl-2-butenyl-N-methylamino a group or the like.

15 The term "N-C<sub>2-6</sub> alkynyl-N-C<sub>1-6</sub> alkylamino group" used in the present specification means a group in which one of the hydrogen atoms within an amino group is substituted by the "C<sub>2-6</sub> alkynyl group" defined above, and the other hydrogen atom is substituted by the "C<sub>1-6</sub> alkyl group" defined above. Specifically, examples of "N-C<sub>2-6</sub> alkynyl-N-C<sub>1-6</sub> alkylamino group" include a  
 20 N-ethynyl-N-methylamino group, a N-1-propynyl-N-methylamino group, a N-2-propynyl-N-methylamino group, a N-1-butynyl-N-methylamino group, a N-2-butynyl-N-methylamino group, a N-3-butynyl-N-methylamino group, a N-pentynyl-N-methylamino group, a N-hexynyl-N-methylamino group, a N-hexanediynyl-N-methylamino group or the like, and preferably, a  
 25 N-ethynyl-N-methylamino group, a N-1-propynyl-N-methylamino group, a

N-2-propynyl-N-methylamino group, a N-1-butyryl-N-methylamino group, a N-2-butyryl-N-methylamino group, a N-3-butyryl-N-methylamino group or the like.

The term "N-C<sub>3-8</sub> cycloalkyl-N-C<sub>1-6</sub> alkylamino group" used in the present specification means a group in which one of the hydrogen atoms within an amino group is substituted by the "C<sub>3-8</sub> cycloalkyl group" defined above, and the other hydrogen atom is substituted by the "C<sub>1-6</sub> alkyl group" defined above. Specifically, examples of "N-C<sub>3-8</sub> cycloalkyl-N-C<sub>1-6</sub> alkylamino group" include a N-cyclopropyl-N-methylamino group, a N-cyclobutyl-N-methylamino group, a N-cyclopentyl-N-methylamino group, a N-cyclohexyl-N-methylamino group, a N-cycloheptyl-N-methylamino group, a N-cyclooctyl-N-methylamino group or the like, and preferably, a N-cyclopropyl-N-methylamino group, a N-cyclobutyl-N-methylamino group, a N-cyclopentyl-N-methylamino group, a N-cyclohexyl-N-methylamino group or the like.

The term "N-C<sub>6-10</sub> aryl-N-C<sub>1-6</sub> alkylamino group" used in the present specification means a group in which one of the hydrogen atoms within an amino group is substituted by the "C<sub>6-10</sub> aryl group" defined above, and the other hydrogen atom is substituted by the "C<sub>1-6</sub> alkyl group" defined above. Specifically, examples of "N-C<sub>6-10</sub> aryl-N-C<sub>1-6</sub> alkylamino group" include a N-phenyl-N-methylamino group, a N-1-naphthyl-N-methylamino group, a N-2-naphthyl-N-methylamino group, a N-indenyl-N-methylamino group, a N-azulenyl-N-methylamino group, a N-heptalenyl-N-methylamino group or the like, and preferably, a N-phenyl-N-methylamino group, a N-1-naphthyl-N-methylamino group, a N-2-naphthyl-N-methylamino group or the like.

The term "N-C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkyl-N-C<sub>1-6</sub> alkylamino group" used in the present specification means a group in which one of the hydrogen atoms within an



amino group is substituted by the "C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkyl group" defined above, and the other hydrogen atom is substituted by the "C<sub>1-6</sub> alkyl group" defined above. Specifically, examples of "N-C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkyl-N-C<sub>1-6</sub> alkylamino group" include a N-cyclopropylmethyl-N-methylamino group, a N-cyclobutylmethyl-N-methylamino group, a N-cyclopentylmethyl-N-methylamino group, a N-cyclohexylmethyl-N-methylamino group, a N-cyclopropylethyl-N-amino group, a N-cyclobutylethyl-N-methylamino group, a N-cyclopentylethyl-N-methylamino group, a N-cyclohexylethyl-N-methylamino group or the like.

The term "N-C<sub>6-10</sub> aryl C<sub>1-6</sub> alkyl-N-C<sub>1-6</sub> alkylamino group" used in the present specification means a group in which one of the hydrogen atoms within an amino group is substituted by the "C<sub>6-10</sub> aryl C<sub>1-6</sub> alkyl group" defined above, and the other hydrogen atom is substituted by the "C<sub>1-6</sub> alkyl group" defined above. Specifically, examples of "N-C<sub>6-10</sub> aryl C<sub>1-6</sub> alkyl-N-C<sub>1-6</sub> alkylamino group" include a N-benzyl-N-methylamino group, a N-1-naphthylmethyl-N-methylamino group, a N-2-naphthylmethyl-N-methylamino group, a N-phenethyl-N-methylamino group, a N-1-naphthylethyl-N-methylamino group, a N-2-naphthylethyl-N-methylamino group or the like.

The term "halogen atom" used in the present specification refers to a fluorine atom, a chlorine atom, a bromine atom or an iodine atom, and preferably, a fluorine atom, a chlorine atom or a bromine atom.

The term "heteroatom" used in the present specification refers to a nitrogen atom, a sulfur atom or an oxygen atom.

The term "5- to 10-membered heterocyclic group" used in the present specification means a monovalent group derived by removal of any one hydrogen atom from the ring of an aromatic or a non-aromatic that has one or a plurality of

heteroatoms among the atoms constituting the ring, the number of atoms constituting the ring being 5 to 10. Specifically, examples of aromatic "5- to 10-membered heterocyclic group" include a furyl group (for instance, a 2-furyl group, a 3-furyl group or the like), a thienyl group (for instance, a 2-thienyl group, a 3-thienyl group or the like), a pyrrolyl group (for instance, a 1-pyrrolyl group, a 2-pyrrolyl group, a 3-pyrrolyl group or the like), a pyridyl group (for instance, a 2-pyridyl group, a 3-pyridyl group, a 4-pyridyl group or the like), a pyrazinyl group, a pyridazinyl group (for instance, a 3-pyridazinyl group, a 4-pyridazinyl group or the like), a pyrimidinyl group (for instance, a 2-pyrimidinyl group, a 4-pyrimidinyl group, a 5-pyrimidinyl group or the like), a triazolyl group (for instance, a 1,2,3-triazolyl group, a 1,2,4-triazolyl group or the like), a tetrazolyl group (for instance, a 1*H*-tetrazolyl group, a 2*H*-tetrazolyl group or the like), a thiazolyl group (for instance, a 2-thiazolyl group, a 4-thiazolyl group, a 5-thiazolyl group or the like), a pyrazolyl group (for instance, a 3-pyrazolyl group, a 4-pyrazolyl group or the like), an oxazolyl group (for instance, a 2-oxazolyl group, a 4-oxazolyl group, a 5-oxazolyl group or the like), an isooxazolyl group (for instance, a 3-isooxazolyl group, a 4-isooxazolyl group, a 5-isooxazolyl group or the like), an isothiazolyl group (for instance, a 3-isothiazolyl group, a 4-isothiazolyl group, a 5-isothiazolyl group or the like), a quinolyl group (for instance, a 5-quinolyl group, a 6-quinolyl group, a 7-quinolyl group, a 8-quinolyl group or the like), an isoquinolyl group (for instance, a 5-isoquinolyl group, a 6-isoquinolyl group, a 7-isoquinolyl group, a 8-isoquinolyl group or the like), a naphthylidinyl group (for instance, a [1,5]naphthylidin-2-yl group, a [1,5]naphthylidin-3-yl group, a [1,8]naphthylidin-2-yl group, a [1,8]naphthylidin-3-yl group or the like), a quinoxaliny group (for instance, a 5-quinoxaliny group, a 6-quinoxaliny group, a 7-quinoxaliny group, a 8-quinoxaliny group or the like), a cinnoliny group (for instance, a 5-cinnoliny group, a 6-cinnoliny

group, a 7-cinnolinyl group, a 8-cinnolinyl group or the like), a quinazolinyl group (for instance, a 4-quinazolinyl group, a 5-quinazolinyl group, a 6-quinazolinyl group, a 7-quinazolinyl group, a 8-quinazolinyl group or the like), an imidazopyridyl group (for instance, an imidazo[1,2-a]pyridin-6-yl group or the like), a benzothiazolyl group (for instance, a benzothiazol-4-yl group, a benzothiazol-5-yl group, a benzothiazol-6-yl group, a benzothiazol-7-yl group or the like), a benzoxazolyl group (for instance, a benzoxazol-4-yl group, a benzoxazol-5-yl group, a benzoxazol-6-yl group, a benzoxazol-7-yl group or the like), a benzimidazolyl group (for instance, a benzimidazol-4-yl group, a benzimidazol-5-yl group, a benzimidazol-6-yl group, a benzimidazol-7-yl group or the like), an indolyl group (for instance, an indol-4-yl group, an indol-5-yl group, an indol-6-yl group, an indol-7-yl group or the like), a pyrrolopyridyl group (for instance, a 1*H*-pyrrolo[2,3-*b*]pyridin-5-yl group, a pyrrolo[3,2-*b*]pyridin-1-yl group or the like), a thienopyridyl group (for instance, a thieno[2,3-*b*]pyridin-5-yl group, a thieno[3,2-*b*]pyridin-6-yl group or the like), a furopyridyl group (for instance, a furo[2,3-*b*]pyridin-5-yl group, a furo[3,2-*b*]pyridin-6-yl group or the like), a 2,3-dihydro-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl group, a benzothiadiazolyl group (for instance, a benzo[1, 2, 5]thiadiazol-5-yl group or the like), a benzoxadiazolyl group (for instance, a benzo[1, 2, 5]oxadiazol-5-yl group or the like), a pyridopyrimidinyl group (for instance, a pyrido[2,3-*d*]pyrimidin-4-yl group or the like), a benzofuryl group (for instance, a benzofuran-4-yl group, a benzofuran-5-yl group, a benzofuran-6-yl group, a benzofuran-7-yl group or the like), a benzothieryl group (for instance, a benzothiophen-4-yl group, a benzothiophen-5-yl group, a benzothiophen-6-yl group, a benzothiophen-7-yl group or the like), a benzo[1,3]dioxol group (for instance, a benzo[1,3]dioxol-5-yl group or the like) or the like. Specifically, examples of non-aromatic “5-10 membered heterocyclic group” include a pyrrolidinyl

group, a piperidinyl group, a homopiperidinyl group, a piperazinyl group, a homopiperazinyl group, a morpholinyl group, a thiomorpholinyl group, a tetrahydrofuryl group, a tetrahydropyranyl group or the like.

The term "5- to 10-membered heterocyclic group containing at least one nitrogen atom" means a monovalent group derived by removal of one hydrogen atom from any position on the ring of an aromatic or a non-aromatic having from 1 to a plurality of heteroatoms among the atoms constituting the ring, and having at least one nitrogen atom, the number of atoms constituting the ring being 5 to 10. Specifically, examples of "5- to 10-membered heterocyclic group containing at least one nitrogen atom" include a pyrrolyl group, a pyridyl group, a pyrazinyl group, a pyridazinyl group, a pyrimidinyl group, a tetrazolyl group, a thiazolyl group, a pyrazolyl group, an oxazolyl group, an isooxazolyl group, an isothiazolyl group, a quinolyl group, an isoquinolyl group, a naphthyldinyl group, a quinoxalinyl group, a cinnolinyl group, a quinazolinyl group, an imidazopyridyl group, a benzothiazolyl group, a benzoxazolyl group, a benzimidazolyl group, an indolyl group, a pyrrolopyridyl group, a thienopyridyl group, a furopyridyl group, a 2,3-dihydro-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl group, a benzothiadiazolyl group, a benzoxadiazolyl group, a pyridopyrimidinyl group or the like.

The term "5- to 10-membered heterocyclic C<sub>1-6</sub> alkyl group" used in the present specification means a group in which any hydrogen atom within the "C<sub>1-6</sub> alkyl group" defined above is substituted by the "5- to 10-membered heterocyclic group" defined above. Specifically, examples of "5- to 10-membered heterocyclic C<sub>1-6</sub> alkyl group" include a furylmethyl group, a thienylmethyl group, a pyrrolylmethyl group, a pyridylmethyl group, a triazolylmethyl group, a tetrazolylmethyl group, a thiazolylmethyl group, a pyrazolylmethyl group, an oxazolylmethyl group, a

benzo[1,3]dioxol methyl group, a tetrahydrofurylmethyl group, a furylethyl group, a thienylethyl group, a pyrrolylethyl group, a pyridylethyl group, a triazolylethyl group, a tetrazolylethyl group, a thiazolylethyl group, a pyrazolylethyl group, an oxazolylethyl group, a benzo[1,3]dioxol ethyl group, a tetrahydrofurylethyl group or the like.

5           The term "furyl C<sub>1-6</sub> alkyl group" used in the present specification means a group in which any hydrogen atom within the "C<sub>1-6</sub> alkyl group" defined above is substituted by a furyl group. Specifically, examples of "furyl C<sub>1-6</sub> alkyl group" include a furylmethyl group, a furylethyl group or the like.

10           The term "thienyl C<sub>1-6</sub> alkyl group" used in the present specification means a group in which any hydrogen atom within the "C<sub>1-6</sub> alkyl group" defined above is substituted by a thienyl group. Specifically, examples of "thienyl C<sub>1-6</sub> alkyl group" include a thienylmethyl group, a thienylethyl group or the like.

15           The term "benzofuryl C<sub>1-6</sub> alkyl group" used in the present specification means a group in which any hydrogen atom within the "C<sub>1-6</sub> alkyl group" defined above is substituted by a benzofuryl group. Specifically, examples of "benzofuryl C<sub>1-6</sub> alkyl group" include a benzofurylmethyl group, a benzofurylethyl group or the like.

20           The term "benzothienyl C<sub>1-6</sub> alkyl group" used in the present specification means a group in which any hydrogen atom within the "C<sub>1-6</sub> alkyl group" defined above is substituted by a benzothienyl group. Specifically, examples of "benzothienyl C<sub>1-6</sub> alkyl group" include a benzothienylmethyl group, a benzothienylethyl group or the like.

25           The term "5- to 10-membered heterocycle oxy group" used in the present specification means a group in which an oxygen atom is bonded to terminus of the "5-10 membered heterocyclic group" defined above. Specifically, examples of "5-

and 10-membered heterocycle oxy group” include a furyloxy group, a thienyloxy group, a pyrrolyloxy group, a pyridyloxy group, a triazolyloxy group, a tetrazolyloxy group, a thiazolyloxy group, a pyrazolyloxy group, an oxazolyloxy group, a benzo[1,3]dioxol oxy group, a tetrahydrofuryloxy group or the like.

5           The term “5- to 10-membered heterocyclic C<sub>1-6</sub> alkoxy group” used in the present specification means a group in which any hydrogen atom within the “C<sub>1-6</sub> alkoxy group” defined above is substituted by the “5- to 10-membered heterocyclic group” defined above. Specifically, examples of “5- to 10-membered heterocyclic C<sub>1-6</sub> alkoxy group” include a furylmethoxy group, a thienylmethoxy group, a  
10   pyrrolylmethoxy group, a pyridylmethoxy group, a triazolylmethoxy group, a tetrazolylmethoxy group, a thiazolylmethoxy group, a pyrazolylmethoxy group, an oxazolylmethoxy group, a benzo[1,3]dioxol methoxy group, a tetrahydrofurylmethoxy group, a furylethoxy group, a thienylethoxy group, a pyrrolylethoxy group, a pyridylethoxy group, a triazolylethoxy group, a tetrazolylethoxy group, a  
15   thiazolylethoxy group, a pyrazolylethoxy group, an oxazolylethoxy group, a benzo[1,3]dioxol ethoxy group, a tetrahydrofurylethoxy group or the like.

          The term “furyl C<sub>1-6</sub> alkoxy group” used in the present specification means a group in which any hydrogen atom within the “C<sub>1-6</sub> alkoxy group” defined above is substituted by a furyl group. Specifically, examples of “furyl C<sub>1-6</sub> alkoxy group”  
20   include a furylmethoxy group, a furylethoxy group or the like.

          The term “thienyl C<sub>1-6</sub> alkoxy group” used in the present specification means a group in which any hydrogen atom within the “C<sub>1-6</sub> alkoxy group” defined above is substituted by a thienyl group. Specifically, examples of “thienyl C<sub>1-6</sub> alkoxy group” include a thienylmethoxy group, a thienylethoxy group or the like.

25           The term “pyridyl C<sub>1-6</sub> alkoxy group” used in the present specification means

a group in which any hydrogen atom within the "C<sub>1-6</sub> alkoxy group" defined above is substituted by a pyridyl group. Specifically, examples of "pyridyl C<sub>1-6</sub> alkoxy group" include a pyridylmethoxy group, a pyridylethoxy group or the like.

The term "5- to 10-membered heterocyclic C<sub>1-6</sub> alkylthio group" used in the present specification means a group in which any hydrogen atom within the "C<sub>1-6</sub> alkylthio group" defined above is substituted by the "5- to 10-membered heterocyclic group" defined above. Specifically, examples of "5- to 10-membered heterocyclic C<sub>1-6</sub> alkylthio group" include a furylmethylthio group, a thienylmethylthio group, a pyrrolylmethylthio group, a pyridylmethylthio group, a triazolylmethylthio group, a tetrazolylmethylthio group, a thiazolylmethylthio group, a pyrazolylmethylthio group, an oxazolylmethylthio group, a benzo[1,3]dioxol methylthio group, a tetrahydrofurylmethylthio group, a furylethylthio group, a thienylethylthio group, a pyrrolylethylthio group, a pyridylethylthio group, a triazolylethylthio group, a tetrazolylethylthio group, a thiazolylethylthio group, a pyrazolylethylthio group, an oxazolylethylthio group, a benzo[1,3]dioxol ethylthio group, a tetrahydrofurylethylthio group or the like.

The term "mono-5- to 10-membered heterocyclic C<sub>1-6</sub> alkylamino group" used in the present specification means a group in which one hydrogen atom within an amino group is substituted by the "5- to 10-membered heterocyclic C<sub>1-6</sub> alkyl group" defined above. Specifically, examples of "mono-5- to 10-membered heterocyclic C<sub>1-6</sub> alkylamino group" include a furylmethylamino group, a thienylmethylamino group, a pyrrolylmethylamino group, a pyridylmethylamino group, a triazolylmethylamino group, a tetrazolylmethylamino group, a thiazolylmethylamino group, a pyrazolylmethylamino group, an oxazolylmethylamino group, a tetrahydrofurylmethylamino group, a furylethylamino group, a thienylethylamino

group, a pyrrolylethylamino group, a pyridylethylamino group, a triazolylethylamino group, a tetrazolylethylamino group, a thiazolylethylamino group, a pyrazolylethylamino group, an oxazolylethylamino group, a tetrahydrofurylethylamino group, a triazolyl-1-propylamino group or the like.

5           The term "N-5- to 10-membered heterocyclic C<sub>1-6</sub> alkyl-N-C<sub>1-6</sub> alkylamino group" used in the present specification means a group in which one of the hydrogen atoms within an amino group is substituted by the "5- to 10-membered heterocyclic C<sub>1-6</sub> alkyl group" defined above and the other hydrogen atom is substituted by the "C<sub>1-6</sub> alkyl group" defined above. Specifically, examples of "N-5- to 10-membered

10 heterocyclic C<sub>1-6</sub> alkyl-N-C<sub>1-6</sub> alkylamino group" include a N-furylmethyl-N-methylamino group, a N-thienylmethyl-N-methylamino group, a N-pyrrolylmethyl-N-methylamino group, a N-pyridylmethyl-N-methylamino group, a N-triazolylmethyl-N-methylamino group, a N-tetrazolylmethyl-N-methylamino group, a N-thiazolylmethyl-N-methylamino group, a N-pyrazolylmethyl-N-methylamino group,

15 a N-oxazolylmethyl-N-methylamino group, a N-tetrahydrofurylmethyl-N-methylamino group, a N-furylethyl-N-methylamino group, a N-thienylethyl-N-methylamino group, a N-pyrrolylethyl-N-methylamino group, a N-pyridylethyl-N-methylamino group, a N-triazolylethyl-N-methylamino group, a N-tetrazolylethyl-N-methylamino group, a N-thiazolylethyl-N-methylamino group, a N-pyrazolylethyl-N-methylamino group, a

20 N-oxazolylethyl-N-methylamino group, a N-tetrahydrofurylethyl-N-methylamino group or the like.

          The term "C<sub>6-10</sub> aryloxy C<sub>1-6</sub> alkyl group" used in the present specification means a group in which any hydrogen atom within the "C<sub>1-6</sub> alkyl group" defined above is substituted by the "C<sub>6-10</sub> aryloxy group" defined above. Specifically,

25 examples of "C<sub>6-10</sub> aryloxy C<sub>1-6</sub> alkyl group" include a phenoxymethyl group, a



1-naphthyloxymethyl group, a 2-naphthyloxymethyl group, an indenylloxymethyl group, an azulenyloxymethyl group, a heptalenylloxymethyl group or the like, and preferably, a phenoxymethyl group, a 1-naphthyloxymethyl group, a 2-naphthyloxymethyl group or the like.

5           The term "phenoxy C<sub>1-6</sub> alkyl group" used in the present specification means a group in which any hydrogen atom within the "C<sub>1-6</sub> alkyl group" defined above is substituted by a phenoxy group. Specifically, examples of "phenoxy C<sub>1-6</sub> alkyl group" include a phenoxymethyl group or the like.

10           The term "5- to 10-membered heterocycle oxy C<sub>1-6</sub> alkyl group" used in the present specification means a group in which any hydrogen atom within the "C<sub>1-6</sub> alkyl group" defined above is substituted by the "5- to 10-membered heterocycle oxy group". Specifically, examples of "5- to 10-membered heterocycle oxy C<sub>1-6</sub> alkyl group" include a furyloxymethyl group, a thienylloxymethyl group, a pyrrolyloxymethyl group, a pyridylloxymethyl group, a triazolyloxymethyl group, a tetrazolyloxymethyl group, a thiazolyloxymethyl group, a pyrazolyloxymethyl group, an oxazolyloxymethyl group, a benzo[1,3]dioxol oxymethyl group, a tetrahydrofuryloxymethyl group or the like, and preferably, a furyloxymethyl group, a thienylloxymethyl group, a pyrrolyloxymethyl group, a pyridylloxymethyl group or the like.

20           The term "pyridyloxy C<sub>1-6</sub> alkyl group" used in the present specification means a group in which any hydrogen atom within the "C<sub>1-6</sub> alkyl group" defined above is substituted by a pyridyl group. Specifically, examples of "pyridyloxy C<sub>1-6</sub> alkyl group" include a pyridylloxymethyl group or the like.

25           The term "which may have a substituent" used in the present specification means that a group may have from one to a plurality of substituents combined at replaceable positions arbitrarily.

The term "having a substituent" used in the present specification refers to having from one to a plurality of substituents combined at replaceable positions arbitrarily.


The term "A" used in the present specification means a 5- to 10-membered heterocyclic group containing at least one nitrogen atom (with the proviso that A may have 1 to 3 substituents selected from the substituent group a-1 or a-2 defined above). Preferable examples of "A" include an aromatic 5- to 10-membered heterocyclic group containing at least one nitrogen atom, such as, a 3-pyridyl group, a pyrazinyl group, a pyrimidinyl group, a pyrazolyl group, a quinolyl group, an isoquinolyl group, a naphthylidinyl group, a quinoxalinyl group, a cinnolinyl group, a quinazolinyl group, an imidazopyridyl group, a benzothiazolyl group, a benzoxazolyl group, a benzimidazolyl group, an indolyl group, a 1*H*-pyrrolopyridyl group, a thienopyridyl group, a furopyridyl group, a 2,3-dihydro-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl group, a benzothiadiazolyl group or a pyrido[2,3-*d*]pyrimidinyl group (with the proviso that each above-mentioned group may have 1 to 3 substituents selected from the substituent groups a-1 and a-2 defined above).

The term "X" used in the present specification means a group represented by the formula  $\text{-NH-C(=Y)-(CH}_2\text{)}_n\text{-}$ , a group represented by the formula  $\text{-C(=Y)-NH-(CH}_2\text{)}_n\text{-}$ , a group represented by the formula  $\text{-C(=Z)-(CH}_2\text{)}_n\text{-}$ , a group represented by the formula  $\text{-CH}_2\text{-NH-(CH}_2\text{)}_n\text{-}$ , a group represented by the formula  $\text{-NH-CH}_2\text{-(CH}_2\text{)}_n\text{-}$  or a group represented by the formula  $\text{-Z-CH}_2\text{-(CH}_2\text{)}_n\text{-}$  (wherein Y, Z and n have the same meanings as defined above, respectively). Preferable examples of "X" include a group represented by the formula  $\text{-NH-C(=Y)-CH}_2\text{-}$ , a group represented by the formula  $\text{-C(=Y)-NH-CH}_2\text{-}$ , a group represented by the formula  $\text{-CH}_2\text{-NH-}$  or a group represented by the formula  $\text{-NH-CH}_2\text{-}$  (wherein Y has

the same meaning as defined above).

The term "A<sup>1</sup>" used in the present specification means a 3-pyridyl group, a pyrazinyl group, a pyrimidinyl group, a pyrazolyl group, a quinolyl group, an isoquinolyl group, a naphthyldinyl group, a quinoxalinyl group, a cinnolinyl group, a quinazolinyl group, an imidazopyridyl group, a benzothiazolyl group, a benzoxazolyl group, a benzimidazolyl group, an indolyl group, a pyrrolopyridyl group, a thienopyridyl group, a furopyridyl group, a 2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-5-yl group or a benzothiadiazolyl group (with the proviso that each above-mentioned group may have 1 to 3 substituents selected from the substituent groups a-1 and a-2 defined above). Preferable examples of "A<sup>1</sup>" include a 3-pyridyl group, a quinolyl group, a naphthyldinyl group, a quinoxalinyl group, an imidazopyridyl group, a benzothiazolyl group, a pyrrolopyridyl group, a thienopyridyl group or a furopyridyl group (with the proviso that each above-mentioned group may have 1 to 3 substituents selected from the substituent groups a-1 and a-2 defined above), more preferably, a 3-pyridyl group, a 6-quinolyl group, a [1,5]naphthylidin-2-yl group, a 6-quinoxalinyl group, an imidazo[1,2-a]pyridin-6-yl group, a benzothiazol-6-yl group, a 1H-pyrrolo[2,3-b]pyridin-5-yl group, a pyrrolo[3,2-b]pyridin-1-yl group, a thieno[2,3-b]pyridin-5-yl group, a thieno[3,2-b]pyridin-6-yl group or a furo[3,2-b]pyridin-6-yl group (with the proviso that each above-mentioned group may have 1 to 3 substituents selected from the substituent groups c-1 and c-2 defined above), still more preferably, group represented by the formula:


[0039]



[0040]

(wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$  and  $R^5$  have the same meanings as defined above, respectively), a 6-quinolyl group, a [1,5]naphthylidin-2-yl group, a 6-quinoxaliny group, an imidazo[1,2-a]pyridin-6-yl group, a benzothiazol-6-yl group, a pyrrolo[3,2-b]pyridin-1-yl group, a 1*H*-pyrrolo[2,3-b]pyridin-5-yl group which may have  
 5 one amino group, a thieno[2,3-b]pyridin-5-yl group which may have one amino group, a thieno[3,2-b]pyridin-6-yl group which may have one amino group or a furo[3,2-b]pyridin-6-yl group which may have one amino group, and particularly preferably, a group represented by the formula:


[0041]



[0042]

(wherein  $R^1$ ,  $R^2$ ,  $R^6$  and  $R^7$  have the same meanings as defined above, respectively), a 6-quinolyl group, a [1,5]naphthylidin-2-yl group, an imidazo[1,2-a]pyridin-6-yl group or a benzothiazol-6-yl group, most preferably 1) a group represented by the formula:


15 [0043]



[0044]

(wherein  $R^{11}$  has the same meaning as defined above) or 2) group represented by the formula:

20 [0045]



[0046]

(wherein  $R^{14}$  has the same meaning as defined above), a 6-quinolyl group, a [1,5]naphthylidin-2-yl group or an imidazo[1,2-a]pyridin-6-yl group.

In addition, preferable examples of "A<sup>1</sup>" include a 3-pyridyl group, a pyrazinyl group, a pyrimidinyl group, a quinolyl group, an isoquinolyl group, a naphthyldinyl group, a quinoxaliny group, a cinnoliny group, a quinazoliny group, an imidazopyridyl group, a benzothiazolyl group, a benzoxazolyl group, a benzimidazolyl group, an indolyl group, a pyrrolopyridyl group, a thienopyridyl group, a furopyridyl group, a 2,3-dihydro-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl group or a benzothiadiazolyl group (with the proviso that A<sup>1</sup> may have 1 to 3 substituents selected from the substituent groups a-1 and a-2 defined above), more preferably, a 3-pyridyl group, a pyrazinyl group, a pyrimidinyl group, a quinolyl group, an isoquinolyl group, a naphthyldinyl group, a quinoxaliny group, a cinnoliny group, a quinazoliny group, an imidazopyridyl group, a benzothiazolyl group, a benzoxazolyl group, a benzimidazolyl group, an indolyl group, a pyrrolopyridyl group, a thienopyridyl group, a furopyridyl group, a 2,3-dihydro-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl group or a benzothiadiazolyl group (with the proviso that A<sup>1</sup> may have 1 to 3 substituents selected from the substituent groups c-1 and c-2 defined above), still more preferably, a 3-pyridyl group, a pyrazinyl group, a pyrimidinyl group, a quinolyl group, an isoquinolyl group, a naphthyldinyl group, a quinoxaliny group, a cinnoliny group, a quinazoliny group, an imidazopyridyl group, a benzothiazolyl group, a benzoxazolyl group, a benzimidazolyl group, an indolyl group, a pyrrolopyridyl group, a thienopyridyl group, a furopyridyl group, a 2,3-dihydro-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl group or a benzothiadiazolyl group (with the proviso that A<sup>1</sup> may have 1 to 3 substituents selected from the substituent groups c'-1 and c'-2 defined above).

Preferable examples of "a group represented by the formula  $-C(=N-R^{a1}) R^{a2}$ " (wherein  $R^{a1}$  and  $R^{a2}$  have the same meanings as defined above, respectively) used in the present specification includes a group represented by the formula  $-C(=N-OH)R^{a2}$  (wherein  $R^{a2}$  has the same meaning as defined above), and more preferably, a group represented by the formula  $-C(=N-OH)CH_3$ .

The term "X<sup>1</sup>" used in the present specification means a group represented by the formula  $-NH-C(=O)-$ , a group represented by the formula  $-C(=O)-NH-$ , a group represented by the formula  $-NH-C(=S)-$ , a group represented by the formula  $-C(=S)-NH-$ , a group represented by the formula  $-NH-C(=NR^{Y1})-$  or a group represented by the formula  $-C(=R^{Y1})-NH-$  (wherein  $R^{Y1}$  has the same meaning as defined above). Preferable examples of "X<sup>1</sup>" include a group represented by  $-C(=O)-NH-$  or a group represented by the formula  $-NH-C(=O)-$ , more preferably a group represented by  $-C(=O)-NH-$ .

The term "E" used in the present specification refers to a furyl group, a thienyl group, a pyrrolyl group, a phenyl group, a pyridyl group, a tetrazolyl group, a thiazolyl group or a pyrazolyl group (with the proviso that each above-mentioned group has 1 or 2 substituents selected from the substituent groups a-1 and a-2 defined above). Preferable examples of "E" include a furyl group, a thienyl group, a pyrrolyl group, a phenyl group or a pyridyl group (with the proviso that each above-mentioned group has 1 or 2 substituents selected from the substituent groups a-1 and a-2 defined above), more preferably, a furyl group, a thienyl group, a pyrrolyl group, a phenyl group or a pyridyl group (with the proviso that each above-mentioned group has 1 or 2 substituents selected from the substituent groups e-1 and e-2 defined above), still more preferably, a furyl group, a thienyl group, a pyrrolyl group, a phenyl group or a pyridyl group (with the proviso that each above-mentioned group has one substituent

selected from the substituent groups g-1 and g-2 defined above), particularly preferably, a 2-furyl group, a 2-thienyl group, a 3-pyrrolyl group, a phenyl group, a 2-pyridyl group or a 3-pyridyl group (with the proviso that each above-mentioned group has one substituent selected from the substituent groups g-1 and g-2 defined above).

Examples of compound (I-a) include compounds in which the "X<sup>1</sup>" defined above, the "A<sup>1</sup>" defined above and the "E" defined above are combined arbitrarily; it is preferable that the compound (I-a) be the compounds in which X<sup>1</sup> represents a group represented by the formula -C(=O)-NH- or a group represented by the formula

-NH-C(=O)-,

(1) A<sup>1</sup> represents a 3-pyridyl group and E represents a furyl group;

(2) A<sup>1</sup> represents a 3-pyridyl group and E represents a thienyl group;

(3) A<sup>1</sup> represents a 3-pyridyl group and E represents a pyrrolyl group;

(4) A<sup>1</sup> represents a 3-pyridyl group and E represents a phenyl group;

(5) A<sup>1</sup> represents a 3-pyridyl group and E represents a pyridyl group;

(6) A<sup>1</sup> represents a quinolyl group and E represents a furyl group;

(7) A<sup>1</sup> represents a quinolyl group and E represents a thienyl group;

(8) A<sup>1</sup> represents a quinolyl group and E represents a pyrrolyl group;

(9) A<sup>1</sup> represents a quinolyl group and E represents a phenyl group;

(10) A<sup>1</sup> represents a quinolyl group and E represents a pyridyl group;

(11) A<sup>1</sup> represents a naphthyldinyl group and E represents a furyl group;

(12) A<sup>1</sup> represents a naphthyldinyl group and E represents a thienyl group;

(13) A<sup>1</sup> represents a naphthyldinyl group and E represents a pyrrolyl group;

(14) A<sup>1</sup> represents a naphthyldinyl group and E represents a phenyl group;

(15) A<sup>1</sup> represents a naphthyldinyl group and E represents a pyridyl group;

- (16) A<sup>1</sup> represents a quinoxaliny group and E represents a furyl group;
- (17) A<sup>1</sup> represents a quinoxaliny group and E represents a thienyl group;
- (18) A<sup>1</sup> represents a quinoxaliny group and E represents a pyrrolyl group;
- (19) A<sup>1</sup> represents a quinoxaliny group and E represents a phenyl group;
- 5 (20) A<sup>1</sup> represents a quinoxaliny group and E represents a pyridyl group;
- (21) A<sup>1</sup> represents an imidazopyridyl group and E represents a furyl group;
- (22) A<sup>1</sup> represents an imidazopyridyl group and E represents a thienyl group;
- (23) A<sup>1</sup> represents an imidazopyridyl group and E represents a pyrrolyl group;
- (24) A<sup>1</sup> represents an imidazopyridyl group and E represents a phenyl group;
- 10 (25) A<sup>1</sup> represents an imidazopyridyl group and E represents a pyridyl group;
- (26) A<sup>1</sup> represents a benzothiazolyl group and E represents a furyl group;
- (27) A<sup>1</sup> represents a benzothiazolyl group and E represents a thienyl group;
- (28) A<sup>1</sup> represents a benzothiazolyl group and E represents a pyrrolyl group;
- (29) A<sup>1</sup> represents a benzothiazolyl group and E represents a phenyl group;
- 15 (30) A<sup>1</sup> represents a benzothiazolyl group and E represents a pyridyl group;
- (31) A<sup>1</sup> represents a 1*H*-pyrrolopyridyl group and E represents a furyl group;
- (32) A<sup>1</sup> represents a 1*H*-pyrrolopyridyl group and E represents a thienyl  
group;
- (33) A<sup>1</sup> represents a 1*H*-pyrrolopyridyl group and E represents a pyrrolyl  
20 group;
- (34) A<sup>1</sup> represents a 1*H*-pyrrolopyridyl group and E represents a phenyl  
group;
- (35) A<sup>1</sup> represents a 1*H*-pyrrolopyridyl group and E represents a pyridyl  
group;
- 25 (36) A<sup>1</sup> represents a thienopyridyl group and E represents a furyl group;



(37) A<sup>1</sup> represents a thienopyridyl group and E represents a thienyl group;

(38) A<sup>1</sup> represents a thienopyridyl group and E represents a pyrrolyl group;

(39) A<sup>1</sup> represents a thienopyridyl group and E represents a phenyl group;

(40) A<sup>1</sup> represents a thienopyridyl group and E represents a pyridyl group;

5 (41) A<sup>1</sup> represents a furopyridyl group and E represents a furyl group;

(42) A<sup>1</sup> represents a furopyridyl group and E represents a thienyl group;

(43) A<sup>1</sup> represents a furopyridyl group and E represents a pyrrolyl group;

(44) A<sup>1</sup> represents a furopyridyl group and E represents a phenyl group or

(45) A<sup>1</sup> represents a furopyridyl group and E represents a pyridyl group (with

10 the proviso that A<sup>1</sup> may contain 1 to 3 substituents selected from the substituent groups a-1 and a-2 defined above, and E has 1 or 2 substituents selected from the substituent group a-1 and a-2 defined above), more preferably, compounds in which X<sup>1</sup> represents a group represented by the formula -C(=O)-NH-,

(1) A<sup>1</sup> represents a 3-pyridyl group and E represents a furyl group;

15 (2) A<sup>1</sup> represents a 3-pyridyl group and E represents a thienyl group;

(3) A<sup>1</sup> represents a 3-pyridyl group and E represents a pyrrolyl group;

(4) A<sup>1</sup> represents a 3-pyridyl group and E represents a phenyl group;

(5) A<sup>1</sup> represents a 3-pyridyl group and E represents a pyridyl group;

(6) A<sup>1</sup> represents a 6-quinolyl group and E represents a furyl group;

20 (7) A<sup>1</sup> represents a 6-quinolyl group and E represents a thienyl group;

(8) A<sup>1</sup> represents a 6-quinolyl group and E represents a pyrrolyl group;

(9) A<sup>1</sup> represents a 6-quinolyl group and E represents a phenyl group;

(10) A<sup>1</sup> represents a 6-quinolyl group and E represents a pyridyl group;

(11) A<sup>1</sup> represents a [1,5]naphthylidin-2-yl group and E represents a furyl

25 group;

(12) A<sup>1</sup> represents a [1,5]naphthylidin-2-yl group and E represents a thienyl group;

(13) A<sup>1</sup> represents a [1,5]naphthylidin-2-yl group and E represents a pyrrolyl group;

5 (14) A<sup>1</sup> represents a [1,5]naphthylidin-2-yl group and E represents a phenyl group;

(15) A<sup>1</sup> represents a [1,5]naphthylidin-2-yl group and E represents a pyridyl group;

10 (16) A<sup>1</sup> represents an imidazo[1,2-a]pyridin-6-yl group and E represents a furyl group;

(17) A<sup>1</sup> represents an imidazo[1,2-a]pyridin-6-yl group and E represents a thienyl group;

(18) A<sup>1</sup> represents an imidazo[1,2-a]pyridine-6-yl group and E represents a pyrrolyl group;

15 (19) A<sup>1</sup> represents an imidazo[1,2-a]pyridin-6-yl group and E represents a phenyl group;

(20) A<sup>1</sup> represents an imidazo[1,2-a]pyridin-6-yl group and E represents a pyridyl group;

(21) A<sup>1</sup> represents a benzothiazol-6-yl group and E represents a furyl group;

20 (22) A<sup>1</sup> represents a benzothiazol-6-yl group and E represents a thienyl group;

(23) A<sup>1</sup> represents a benzothiazol-6-yl group and E represents a pyrrolyl group;

25 (24) A<sup>1</sup> represents a benzothiazol-6-yl group and E represents a phenyl group or

(25) A<sup>1</sup> represents a benzothiazol-6-yl group and E represents a pyridyl group (with the proviso that A<sup>1</sup> may contain 1 to 3 substituents selected from the substituent groups c-1 and c-2 defined above, and E has 1 or 2 substituents selected from the substituent groups e-1 and e-2 defined above), still more preferably, compounds in

5 which X<sup>1</sup> represents a group represented by the formula -C(=O)-NH-,

(1) A<sup>1</sup> represents a 3-pyridyl group and E represents a 2-furyl group;

(2) A<sup>1</sup> represents a 3-pyridyl group and E represents a 2-thienyl group;

(3) A<sup>1</sup> represents a 3-pyridyl group and E represents a 3-pyrrolyl group;

(4) A<sup>1</sup> represents a 3-pyridyl group and E represents a phenyl group;

10 (5) A<sup>1</sup> represents a 3-pyridyl group and E represents a 2-pyridyl group;

(6) A<sup>1</sup> represents a 3-pyridyl group and E represents a 3-pyridyl group;

(7) A<sup>1</sup> represents a 6-quinolyl group and E represents a 2-furyl group;

(8) A<sup>1</sup> represents a 6-quinolyl group and E represents a 2-thienyl group;

(9) A<sup>1</sup> represents a 6-quinolyl group and E represents a 3-pyrrolyl group;

15 (10) A<sup>1</sup> represents a 6-quinolyl group and E represents a phenyl group;

(11) A<sup>1</sup> represents a 6-quinolyl group and E represents a 2-pyridyl group;

(12) A<sup>1</sup> represents a 6-quinolyl group and E represents a 3-pyridyl group;

(13) A<sup>1</sup> represents a [1,5]naphthylidin-2-yl group and E represents a 2-furyl group;

20 (14) A<sup>1</sup> represents a [1,5]naphthylidin-2-yl group and E represents a 2-thienyl group;

(15) A<sup>1</sup> represents a [1,5]naphthylidin-2-yl group and E represents a 3-pyrrolyl group;

25 (16) A<sup>1</sup> represents a [1,5]naphthylidin-2-yl group and E represents a phenyl group;

(17) A<sup>1</sup> represents a [1,5]naphthylidin-2-yl group and E represents a 2-pyridyl group;

(18) A<sup>1</sup> represents a [1,5]naphthylidin-2-yl group and E represents a 3-pyridyl group;

5 (19) A<sup>1</sup> represents an imidazo[1,2-a]pyridin-6-yl group and E represents a 2-furyl group;

(20) A<sup>1</sup> represents an imidazo[1,2-a]pyridin-6-yl group and E represents a 2-thienyl group;

10 (21) A<sup>1</sup> represents an imidazo[1,2-a]pyridin-6-yl group and E represents a 3-pyrrolyl group;

(22) A<sup>1</sup> represents an imidazo[1,2-a]pyridin-6-yl group and E represents a phenyl group;

(23) A<sup>1</sup> represents an imidazo[1,2-a]pyridin-6-yl group and E represents a 2-pyridyl group;

15 (24) A<sup>1</sup> represents an imidazo[1,2-a]pyridin-6-yl group and E represents a 3-pyridyl group;

(25) A<sup>1</sup> represents a benzothiazol-6-yl group and E represents a 2-furyl group;

20 (26) A<sup>1</sup> represents a benzothiazol-6-yl group and E represents a 2-thienyl group;

(27) A<sup>1</sup> represents a benzothiazol-6-yl group and E represents a 3-pyrrolyl group;

(28) A<sup>1</sup> represents a benzothiazol-6-yl group and E represents a phenyl group;

25 (29) A<sup>1</sup> represents a benzothiazol-6-yl group and E represents a 2-pyridyl

group or

(29) A<sup>1</sup> represents a benzothiazol-6-yl group and E represents a 3-pyridyl group (with the proviso that A<sup>1</sup> may contain 1 to 3 substituents selected from the substituent groups c-1 and c-2 defined above, and E has one substituent selected  
5 from the substituent group g-1 and g-2 defined above).

Examples of compound (I-b) include compounds in which the "X<sup>2</sup>" defined above, the "A<sup>2</sup>" defined above and "R<sup>10</sup>" defined above are combined arbitrarily, preferably,

(1) compounds in which X<sup>2</sup> represents a group represented by the formula  
10 -O-CH<sub>2</sub>-, A<sup>2</sup> represents a 6-quinolyl group and R<sup>10</sup> represents a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkoxy group;

(2) compounds in which X<sup>2</sup> represents a group represented by the formula  
-S-CH<sub>2</sub>-, A<sup>2</sup> represents a 6-quinolyl group and R<sup>10</sup> represents a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkoxy group;

15 (3) compounds in which X<sup>2</sup> represents a group represented by the formula  
-C(=O)-CH<sub>2</sub>-, A<sup>2</sup> represents a 6-quinolyl group and R<sup>10</sup> represents a C<sub>6-10</sub> aryloxy group;

(4) compounds in which X<sup>2</sup> represents a group represented by the formula  
-C(=O)-CH<sub>2</sub>-, A<sup>2</sup> represents a 6-quinolyl group and R<sup>10</sup> represents a C<sub>6-10</sub> aryl C<sub>1-6</sub>  
20 alkoxy group;

(5) compounds in which X<sup>2</sup> represents a group represented by the formula  
-NH-CH<sub>2</sub>-, A<sup>2</sup> represents a 6-quinolyl group and R<sup>10</sup> represents a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkoxy group;

(6) compounds in which X<sup>2</sup> represents a group represented by the formula  
25 -CH<sub>2</sub>-NH-, A<sup>2</sup> represents a 6-quinolyl group and R<sup>10</sup> represents a C<sub>6-10</sub> aryl C<sub>1-6</sub>

alkoxy group;

(7) compounds in which  $X^2$  represents a group represented by the formula -NH-CH<sub>2</sub>-,  $A^2$  represents a 4-quinazoliny group and  $R^{10}$  represents a C<sub>1-6</sub> alkyl group;

(8) compounds in which  $X^2$  represents a group represented by the formula  
5 -NH-CH<sub>2</sub>-,  $A^2$  represents a 4-quinazoliny group and  $R^{10}$  represents a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkoxy group or

(9) compounds in which  $X^2$  represents a group represented by the formula -NH-CH<sub>2</sub>-,  $A^2$  represents a pyrido[2,3-d]pyrimidin-4-yl group which may have an amino group and  $R^{10}$  represents a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkoxy group, more preferably,

10 (1) compounds in which  $X^2$  represents a group represented by the formula -NH-CH<sub>2</sub>-,  $A^2$  represents a 6-quinolyl group and  $R^{10}$  represents a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkoxy group;

(2) compounds in which  $X^2$  represents a group represented by the formula -CH<sub>2</sub>-NH-,  $A^2$  represents a 6-quinolyl group and  $R^{10}$  represents a C<sub>6-10</sub> aryl C<sub>1-6</sub>

15 alkoxy group;

(3) compounds in which  $X^2$  represents a group represented by the formula -NH-CH<sub>2</sub>-,  $A^2$  represents a 4-quinazoliny group and  $R^{10}$  represents a C<sub>1-6</sub> alkyl group;

(4) compounds in which  $X^2$  represents a group represented by the formula -NH-CH<sub>2</sub>-,  $A^2$  represents a 4-quinazoliny group and  $R^{10}$  represents a C<sub>6-10</sub> aryl C<sub>1-6</sub>

20 alkoxy group or

(5) compounds in which  $X^2$  represents a group represented by the formula -NH-CH<sub>2</sub>-,  $A^2$  represents a pyrido[2,3-d]pyrimidin-4-yl group which may have an amino group and  $R^{10}$  represents a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkoxy group.

Examples of the term "salt" used in the present specification include a salt  
25 with an inorganic acid, a salt with an organic acid, a salt with an inorganic base, a

salt with an organic base, a salt with an acidic or basic amino acid or the like. Among these salts, it is preferable that a salt used herein be a pharmaceutically acceptable.

5       Preferable examples of the salt with the inorganic acid include salts with hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid or the like. Preferable examples of the salt with the organic acid include salts with acetic acid, succinic acid, fumaric acid, maleic acid, tartaric acid, citric acid, lactic acid, stearic acid, benzoic acid, methanesulfonic acid, ethanesulfonic acid, *p*-toluenesulfonic acid or the like.

10       Preferable examples of the salt with the inorganic base include alkaline metal salts such as sodium salt and potassium salt, alkaline earth metal salts such as calcium salt and magnesium salt, aluminum salt, ammonium salt or the like. Preferable examples of the salt with the organic base include salts with diethyl amine, diethanolamine, meglumine, N,N-dibenzyl ethylenediamine or the like.

15       Preferable examples of the salt with the acidic amino acid include salts with aspartic acid, glutamic acid or the like. Preferable examples of the salt with the basic amino acid include salts with arginine, lysine, ornithine or the like.

The term "antifungal agent" used in the present specification refers to a preventive agent or a therapeutic agent for fungal infection.

20       The compounds according to the present invention, or salts or hydrates thereof, can be formulated into tablets, powders, fine granules, granules, coated tablets, capsules, syrups, troches, inhalants, suppositories, injections, ointments, eye ointments, tapes, eye drops, nose drops, ear drops, cataplasms, lotions or the like, by the conventional methods. Such formulation can be achieved by using  
25       typical diluents, binders, lubricants, colorants, flavorants, and, as necessary,

stabilizers, emulsifiers, absorbefaciants, surfactants, pH modulators, preservatives, antioxidants or the like, and materials commonly used as ingredients of pharmaceutical preparations according to the conventional methods. For example, an oral preparation can be produced by combining a compound according to the present invention or a pharmaceutically acceptable salt thereof with a diluent, and if required, a binder, a disintegrating agent, a lubricant, a colorant, a flavorant or the like, and formulating the mixture into powders, fine granules, granules, tablets, coated tablets, capsules or the like according to the conventional methods. Examples of the materials include animal and vegetable oils such as soya bean oil, beef tallow, and synthetic glyceride; hydrocarbons such as liquid paraffin, squalane, and solid paraffin; ester oils such as octyldodecyl myristate and *iso*-propyl myristate; higher alcohols such as cetostearyl alcohol and behenyl alcohol; silicone resins; silicone oils; surfactants such as polyoxyethylene fatty acids ester, sorbitan fatty acid ester, polyoxyethylene sorbitan fatty acid ester, polyoxyethylene hydrogenated castor oil, and polyoxyethylene polyoxypropylene block co-polymer; water-soluble polymers such as hydroxyethyl cellulose, polyacrylic acid, carboxyvinyl polymer, polyethylene glycol, polyvinylpyrrolidone, and methyl cellulose; lower alcohols such as ethanol and isopropanol; polyhydric alcohols such as glycerol, propylene glycol, dipropylene glycol, and sorbitol; sugars such as glucose and sucrose; inorganic powder such as anhydrous silicic acid, magnesium aluminum silicate, and aluminum silicate; and pure water. Examples of the diluents include lactose, corn starch, white sugar, glucose, mannitol, sorbitol, crystalline cellulose, silicon dioxide or the like. Examples of the binders include polyvinyl alcohol, polyvinyl ether, methylcellulose, ethylcellulose, gum Arabic, tragacanth, gelatin, shellac, hydroxypropyl methylcellulose, hydroxypropyl cellulose, polyvinylpyrrolidone, polypropylene glycol-polyoxyethylene block



co-polymer, and meglumine or the like. Examples of disintegrating agents include starch, agar, gelatin powder, crystalline cellulose, calcium carbonate, sodium hydrogencarbonate, calcium citrate, dextrin, pectin, calcium carboxymethyl cellulose or the like. Examples of lubricants include magnesium stearate, talc, polyethylene glycol, silica, hydrogenated vegetable oil or the like. Examples of colorants include those pharmaceutically acceptable. Examples of flavorants include cocoa powder, peppermint camphor, aromatic powder peppermint oil, Borneo camphor, cinnamon powder or the like. Tablets and granules may be coated with sugar, or if required, other appropriate coatings can be made. Solutions, such as syrups or injectable preparations, to be administered can be formulated by combining a compound according to the present invention or a pharmaceutically acceptable salt thereof with a pH modulator, a solubilizing agent, an isotonicizing agent or the like, and if required, with an auxiliary solubilizing agent, a stabilizer or the like, according to the conventional methods. Methods for manufacturing an external preparations are not limited and such preparations can be manufactured by the conventional methods. Specifically, various materials typically used for manufacturing pharmaceuticals, quasi drugs, cosmetics or the like can be used as base materials for the external formulation. More specifically, examples of base materials to be used include animal and vegetable oils, mineral oils, ester oils, wax, higher alcohols, fatty acids, silicone oil, surfactants, phospholipids, alcohols, polyhydric alcohols, water-soluble polymers, clay minerals, pure water or the like. Furthermore, external preparations of the present invention can contain, as required, pH modulators, antioxidants, chelating agents, antibacterial/antifungal agents, colorants, odoriferous substances or the like. But this does not limit the type of base materials that are to be used in the external preparations of the present invention. If required, the preparation may

contain differentiation inducers, blood flow improving agents, antimicrobial agents, antiphlogistics, cell activators, vitamins, amino acids, humectants, keratolytic agents or the like. The amount of the base materials listed above is adjusted within a concentration range used for producing typical external preparations.

5           When administering a compound according to the present invention or a salt thereof, or a hydrate thereof, the forms of the compounds are not limited in particular, and the compound can be given orally or parenterally by the conventional method. For instance, the compound can be administered as a dosage form such as tablets, powders, granules, capsules, syrups, troches, inhalants, suppositories, injections,  
10 ointments, eye ointments, tapes, eye drops, nasal drops, ear drops, cataplasms and lotions.

Dose of a pharmaceutical of the present invention can be selected appropriately according to symptom severity, age, sex, body weight, forms of administration, type of salts, specific type of disease or the like.

15           The dose varies remarkably depending on the patient's disease, symptom severity, age and sex, drug susceptibility or the like. An oral preparation of the present invention can be generally administered once or several times at a dose of from 1 to 10000 mg/adult/day, preferably from 10 to 2000 mg/adult/day. An injection of the present invention can be generally administered at a dose of from 0.1 to 10000  
20 mg/adult/day, preferably from 1 to 2000 mg/adult/day.


The method for preparing compounds according to the present invention represented by Formula (I) (hereinafter referred to as compound (I)) will be described. The compounds according to the present invention can be prepared by the  
25 conventional organic synthesis methods. For instance, among the compounds (I),

the compounds represented by the following Formula (1a), Formula (2a), Formula (3a), Formula (3b), Formula (3c), Formula (3d) and Formula (3e) (hereinafter referred to as compound (1a), compound (2a), compound (3a), compound (3b), compound (3c), compound (3d) and compound (3e), respectively) can be prepared by the methods described in the following [Preparation Method 1] to [Preparation Method 3] or the like. In addition, conversion of substituents on A and E of compound (I) according to the present invention can be accomplished by the methods described in the following [Preparation Method 4-1] to [Preparation Method 4-5] or the like.

# 10 [General synthesis method]

[Preparation Method 1] Representative preparation method for compound (1a)

[0047]




15 [0048]

[wherein each symbol has the same meaning as defined above.]

[Preparation Method 1-1] Amidation

[0049]



20

[0050]

[wherein each symbol has the same meaning as defined above.]

Compound (1b) which is a commercially available product can be used, or compound (1b) can also be prepared from a commercially available product by the well known method. In addition, compound (1b) can also be prepared by the methods described in the preparation examples among the following examples.

5       Compound (1c) which is a commercially available product can be used, or compound (1c) can also be prepared from a commercially available product by the well known method. In addition, compound (1c) can also be prepared by the method described in the preparation examples among the following examples or [Preparation Method 1-2-1], or the like.

10

[Step 1]

This step is a step wherein compound (1b) and compound (1c) are condensed in a solvent using a condensing agent to obtain compound (1a). The solvent used is not limited in particular. Examples of the solvent in this step include  
15   halogenated hydrocarbons such as dichloromethane and chloroform; sulfoxides such as dimethylsulfoxide; esters such as ethyl acetate; ethers such as tetrahydrofuran and 1,4-dioxane; amides such as N,N-dimethylformamide, and N,N-dimethylacetamide. Examples of the condensing agent include Bop (1*H*-1,2,3-benzotriazole-1-yloxy(tri(dimethylamino))phosphonium  
20   hexafluorophosphate), WSC (1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride), DCC (N,N-dicyclohexylcarbodiimide), CDI (carbonyldiimidazole), diethylphosphoryl cyanide or the like. Compound (1c) can be used in amounts of 1 equivalent to 1.5 equivalents based on compound (1b). The condensing agent can be used in amounts of 1 equivalent to 1.5 equivalents based on compound (1b). In  
25   addition, it is adequate to add from 1 equivalent to excess amount of an organic base,

for instance, triethylamine or the like, as necessary. The reaction temperature is from room temperature to 80°C, and the reaction time is from 10 minutes to 30 hours.

Furthermore, compound (1a) can also be prepared from compound (1b) and compound (1c) by the method described in the other method (1), (2) or (3) below.

5           Other method (1): Compound (1a) can be obtained by converting compound (1b) into an acid chloride, then, reacting the acid chloride and compound (1c). The step for obtaining the acid chloride is carried out by reacting from 1 equivalent to excess amount of acid chloride synthesis reagent based on compound (1b), without solvent or in the presence of a solvent such as, dichloromethane,  
10 benzene, toluene or the like. Catalytic amounts of N,N-dimethylformamide may be added in the reaction system. Examples of the acid chloride synthesis reagent include, for instance, thionyl chloride, oxalyl chloride, phosphorus trichloride, phosphorus pentachloride or the like. The reaction temperature is from room temperature to reflux temperature, and the reaction time is from 10 minutes to 24  
15 hours.

The step for condensing the acid chloride and compound (1c) is carried out by reacting the acid chloride and compound (1c) in a solvent such as, for instance, dichloromethane, tetrahydrofuran, N,N-dimethylformamide or the like, in the presence of 1 equivalent to 3 equivalents of base based on the acid chloride. Examples of the  
20 base include, for instance, an organic base such as triethylamine, pyridine and the like or an inorganic base such as potassium carbonate, cesium carbonate or the like. Compound (1c) is used in the amounts of 1 equivalent to 1.5 equivalents based on the acid chloride. The reaction time is from 10 minutes to 24 hours, and the reaction temperature is from 0°C to reflux temperature.

25           Other method (2): Compound (1a) can be obtained by converting

compound (1b) into mixed acid anhydride, then, reacting the mixed acid anhydride with compound (1c). The step for obtaining the mixed acid anhydride is carried out by reacting compound (1b) and, for instance, chloroformates such as ethyl chloroformate or the like, in the presence of a base such as, for instance, triethylamine or the like. Chloroformates and base are used in the amounts of 1  
5 equivalent to 2 equivalents based on compound (1b). The reaction time is from 10 minutes to 5 hours. The reaction temperature is from 0°C to room temperature.

The step for condensing the mixed acid anhydride and compound (1c) is carried out by reacting the mixed acid anhydride and compound (1c) in a solvent  
10 such as, for instance, dichloromethane, tetrahydrofuran, N,N-dimethylformamide or the like. Compound (1c) is used in the amounts of 1 equivalent to 1.5 equivalents based on the mixed acid anhydride. The reaction time is from 10 minutes to 24 hours, and the reaction temperature is from 0°C to 50°C.

Other method (3): Compound (1a) can be obtained by converting  
15 compound (1b) into an active ester, then, reacting the active ester and compound (1c). The step for obtaining the active ester is carried out by reacting compound (1b) and an active ester synthesis reagent in a solvent such as, for instance, 1,4-dioxane, tetrahydrofuran or the like, in the presence of a condensing agent such as, for instance, DCC or the like. Examples of the active ester synthesis reagent  
20 include, for instance, N-hydroxysuccinimide or the like. The active ester synthesis reagent and the condensing agent are used in the amounts of 1 equivalent to 1.5 equivalents based on compound (1b). The reaction temperature is from 0°C to room temperature. The reaction time is from 2 hours to 24 hours.


The step for condensing the active ester and compound (1c) is carried out by  
25 reacting the active ester and compound (1c) in a solvent such as, for instance,

dichloromethane, tetrahydrofuran, N,N-dimethylformamide or the like. Compound (1c) is used in the amounts of 1 equivalent to 1.5 equivalents based on the active ester. The reaction temperature is from 0°C to 50°C, and the reaction time is from 10 minutes to 24 hours.

- 5           Note that, after [Step 1], a substituent on A and E of compound (1a) can be converted using a well known method. Furthermore, a substituent on A of compound (1a) can also be converted using a method described in [Preparation Method 4-1] or [Preparation Method 4-4], and a substituent on E of compound (1a) can also be converted using a method described in [Preparation Method 4-2],
- 10   [Preparation Method 4-3], [Preparation Method 4-5] or the like.

[Preparation Method 1-2-1] Preparation method for compound (1c)

[0051]



[0052]

[wherein E has the same meaning as defined above; L represents a leaving group such as a halogen atom, a methanesulfonyloxy group or a *p*-toluenesulfonyloxy group]

For each compound in the above step figure, a commercially available product can be used as is, or it can also be prepared from a commercially available product by a well known method. In addition, it can be prepared using the method described in the preparation examples among the examples and in [Preparation



Method 1-2-2] to [Preparation Method 1-2-6]. Further, each compound in the above step figure can also be prepared by converting a substituent on E using the method described in [Preparation Method 4-2] to [Preparation Method 4-5] or the like.

5 [Step 1-1]

The present step is a step wherein compound (1c-1) is reduced to obtain compound (1c-2). Examples of the reducing agent include, for instance, sodium borohydride, lithium borohydride, lithium aluminum hydride or the like. Examples of the solvent include alcohols such as, for instance, methanol, ethanol or the like, and  
10 ethers such as, for instance, tetrahydrofuran, diethyl ether or the like. The reducing agent is used in the amounts of 1 equivalent to 10 equivalents based on compound (1c-1). The reaction temperature is from 0°C to reflux temperature, and the reaction time is from 5 minutes to 24 hours.

15 [Step 1-2]

The present step is a step wherein a hydroxyl group of compound (1c-2) is converted into a leaving group to obtain compound (1c-3).

When L represents a methanesulfonyloxy group or *p*-toluenesulfonyloxy group, compound (1c-3) can be obtained by reacting compound (1c-2) and  
20 methanesulfonyl chloride or *p*-toluenesulfonyl chloride in a solvent such as, for instance, dichloromethane or the like, in the presence of an organic base such as, for instance, triethylamine or the like. The organic base is used in the amounts of 2 equivalents to 6 equivalents based on compound (1c-2). Methanesulfonyl chloride or *p*-toluenesulfonyl chloride is used in the amounts of 1 equivalent to 3 equivalents  
25 based on compound (1c-2). The reaction temperature is from 0°C to room

temperature, and the reaction time is from 10 minutes to 24 hours.

When L represents a chlorine atom, compound (1c-3) can be obtained by action of a chlorination reagent such as, for instance, thionyl chloride, oxalyl chloride or the like on compound (1c-2). The chlorination reagent is used in the amounts of 1 equivalent to excess amount based on compound (1c-2). The reaction temperature is from 0°C to room temperature, and the reaction time is from 10 minutes to 24 hours.

[Step 1-3]

The present step is a step wherein compound (1c-3) and phthalimide potassium salt are reacted to obtain compound (1c-4). Compound (1c-4) can be obtained by reacting compound (1c-3) and phthalimide potassium salt in a solvent such as, for instance, N,N-dimethylformamide or the like. Phthalimide is used in the amounts of 1 equivalent to 2 equivalents based on compound (1c-3). The reaction temperature is from room temperature to 160°C. The reaction time is from 10 minutes to 48 hours.

[Step 1-4]

The present step is a step for obtaining compound (1c-8) from compound (1c-4). Compound (1c-8) can be obtained by adding from 1 equivalent to excess amount of hydrazine hydrate based on compound (1c-4) in a solvent such as, for instance, ethanol or the like. The reaction temperature is from 80°C to reflux temperature, and the reaction time is from 10 minutes to 24 hours.

[Step 1-5]

The present step is a step wherein compound (1c-2) and phthalimide are reacted to obtain compound (1c-4). Compound (1c-4) can be obtained by reacting compound (1c-2), phthalimide, triphenylphosphine and diethyl azodicarboxylate or diisopropyl azodicarboxylate, in a solvent such as, for instance, dichloromethane, tetrahydrofuran or the like. Phthalimide, triphenylphosphine and diethyl azodicarboxylate or diisopropyl azodicarboxylate are used in the amounts of 1 equivalent to 2 equivalents based on compound (1c-2). The reaction temperature is from -20°C to 80°C, and the reaction time is from 5 minutes to 48 hours.

## 10 [Step 1-6]

The present step is a step wherein compound (1c-3) and an amine protected with a *tert*-butoxycarbonyl group are reacted to obtain compound (1c-5). Compound (1c-5) can be obtained by reacting compound (1c-3) and the amine protected with the *tert*-butoxycarbonyl group in a solvent such as, for instance, N,N-dimethylformamide or the like, in the presence of a base such as, for instance, sodium hydride or the like. The base is used in the amounts of 1 equivalent to 2 equivalents based on compound (1c-3). The amine protected with the *tert*-butoxycarbonyl group is used in the amounts of 1 equivalent to 2 equivalents based on compound (1c-3). The reaction temperature is from room temperature to 80°C, and the reaction time is from 1 hour to 24 hours.

## [Step 1-7]

The present step is a step wherein the *tert*-butoxycarbonyl group of compound (1c-5) is deprotected to obtain compound (1c-8). Compound (1c-8) can be obtained by deprotecting the *tert*-butoxycarbonyl group of compound (1c-5)

without a solvent or in a solvent such as, for instance, dichloromethane or the like, in the presence of an acid such as trifluoroacetic acid or the like in the amounts of 2 equivalents to excess amount based on compound (1c-5). The reaction temperature is from 0°C to 60°C, and the reaction time is from 10 minutes to 24  
5 hours.

[Step 1-8]

The present step is a step wherein the leaving group of compound (1c-3) is converted into an azide group to obtain compound (1c-6). Compound (1c-6) can be  
10 obtained by reacting compound (1c-3) and an azidation reagent such as, for instance, sodium azide, potassium azide or the like in a solvent such as, for instance, N,N-dimethylformamide or the like. The azidation reagent is used in the amounts of 1 equivalent to 5 equivalents based on compound (1c-3). The reaction temperature is from room temperature to 80°C, and the reaction time is from 10 minutes to 48  
15 hours.

[Step 1-9]

The present step is a step wherein the azide group of compound (1c-6) is reduced to obtain compound (1c-8). Compound (1c-8) can be obtained by carrying  
20 out catalytic hydrogenation using Lindlar catalyst in a solvent such as, for instance, ethanol or the like. Lindlar catalyst is used in catalytic amount to excess amount based on compound (1c-6). The reaction temperature is from room temperature to 80°C, the reaction time is from 30 minutes to 36 hours, and the reaction pressure is from 1 atm to 4 atm.

25 As other method, compound (1c-8) can be obtained by action of

triphenylphosphine in a solvent such as, for instance, dichloromethane, tetrahydrofuran or the like. Triphenylphosphine is used in the amounts of 1.0 equivalent to 2.0 equivalents based on compound (1c-6).

5 [Step 1-10]

The present step is a step wherein a formyl group of compound (1c-1) is converted into a cyano group to obtain compound (1c-7). Compound (1c-7) can be obtained by reacting 1 equivalent to 3 equivalent of hydroxyamine hydrochloride based on compound (1c-1) in a solvent such as, for instance, ethanol or the like, to  
10 obtain an oxime, then, carrying out dehydration reaction by the action of CDI on the oxime. CDI is used in the amounts of 1 equivalent to 5 equivalents based on the oxime. The reaction temperature is from room temperature to reflux temperature, and the reaction time is from 30 minutes to 24 hours.

15 [Step 1-11]

The present step is a step wherein a cyano group of compound (1c-7) is reduced to obtain compound (1c-8). Compound (1c-8) can be obtained either by carrying out a reduction reaction using a reducing agent such as, for instance, lithium aluminum hydride, diisobutylaluminum hydride or the like, or, by carrying out catalytic  
20 hydrogenation under hydrogen atmosphere using a catalyst such as, for instance, Raney nickel, palladium-carbon or the like. The solvent used is not limited, however, when carrying out reduction reaction using the reducing agent, ethers such as, for instance, tetrahydrofuran, diethyl ether or the like, hydrocarbons such as, for instance, toluene or the like, are used. When carrying out catalytic hydrogenation, alcohols  
25 such as, for instance, methanol, ethanol, propanol or the like are used. The

reducing agent is used in the amounts of 1 equivalent to 10 equivalents based on compound (1c-7). The reaction temperature is from room temperature to 80°C, and the reaction time is from 10 minutes to 24 hours. The reaction pressure when carrying out catalytic hydrogenation is from 1 atm to 4 atm.

5

[Step 1-12]

The present step is a step for obtaining compound (1c-8) from compound (1c-1). Compound (1c-8) can be obtained by carrying out catalytic hydrogenation using a catalyst such as Raney nickel or the like, in a solvent containing ammonia,  
10 under hydrogen atmosphere. Examples of the solvent include, but are not limited to, alcohols such as methanol, ethanol, propanol or the like. The reaction temperature is from room temperature to 80°C, the reaction time is from 1 hour to 36 hour, and the reaction pressure is from 1 atm to 4 atm.

15 [Step 1-13]

The present step is a step wherein an acetal group on compound (1c-9) is deprotected to obtain compound (1c-1). Compound (1c-1) can be obtained by dissolving compound (1c-9) in an organic solvent, and by action of an aqueous solution of acid such as, for instance, hydrochloric acid, sulfuric acid, citric acid or the  
20 like. Examples of the solvent include, but are not limited to, for instance, methanol, ethanol, acetonitrile, tetrahydrofuran or the like. The reaction temperature is from room temperature to 60°C, and the reaction time is from 5 minutes to 24 hours.

[Step 1-14]

25 The present step is a step wherein an hydroxyl group of compound (1c-2) is

converted into an azide group to obtain compound (1c-6). Compound (1c-6) can be obtained by reacting compound (1c-2) and diphenylphosphoryl azide in a solvent such as, for instance, benzene, toluene or the like, in the presence of an organic base such as, for instance, 1,8-diazabicyclo[5,4,0]undec-7-ene. The organic base is  
5 used in the amounts of 1 equivalent to 1.5 equivalents based on compound (1c-2). Diphenylphosphoryl azide is used in the amounts of 1 equivalent to 1.5 equivalents based on compound (1c-2). The reaction temperature is from room temperature to 80°C, and the reaction time is from 10 minutes to 48 hours.

10 [Step 1-26]

The present step is a step wherein compound (1c-30) is formylated to obtain compound (1c-1). Compound (1c-1) can be obtained by action of 1 equivalent to 1.5 equivalents of a strong base based on compound (1c-30) for anionization, then, reacting a formylation agent. Examples of the solvent include, for instance,  
15 tetrahydrofuran, diethyl ether or the like. Examples of the strong base include, for instance, *n*-butyl lithium or the like. Examples of the formylation agent include, for instance, N,N-dimethylformamide, N-formylmorpholine or the like. The formylation agent is used in the amounts of 1 equivalent to 2 equivalents based on compound (1c-30). The reaction temperature is from -80°C to room temperature, and the  
20 reaction time is from 5 minutes to 12 hours.

[Step 1-27]


The present step is a step wherein compound (1c-30) is cyanated to obtain compound (1c-7). Compound (1c-7) can be obtained by reacting compound (1c-30)  
25 and zinc cyanide in a solvent such as, for instance, N,N-dimethylformamide,

N-methylpyrrolidinone or the like, under nitrogen atmosphere, in the presence of a catalyst. Examples of the catalyst include, for instance, tetrakis(triphenylphosphine)palladium(0) or the like. Zinc cyanide is used in the amounts of 1 equivalent to 2 equivalents based on compound (1c-30). The catalyst  
 5 is used in the amounts of 0.01 equivalents to 0.1 equivalents based on compound (1c-30). The reaction temperature is from 50°C to reflux temperature, and the reaction time is from 5 minutes to 24 hours.

Other method: Compound (1c-7) can be obtained by reacting compound (1c-30) and copper cyanide in a solvent such as, for instance,  
 10 N,N-dimethylformamide and N-methylpyrrolidinone, under nitrogen atmosphere. Copper cyanide is used in the amounts of 1 equivalent to excess amount based on compound (1c-30). The reaction temperature is from 50°C to reflux temperature, and the reaction time is from 10 minutes to 72 hours.

15 [Preparation Method 1-2-2] Preparation method for compound (1c-11), which is compound (1c-1)

[0053]



[0054]

20 [wherein Q represents an oxygen atom or a sulfur atom; Ar<sup>1</sup> represents a C<sub>6-10</sub> aryl group that may have 1 or 2 substituents selected from the following substituent group i or an aromatic 5-10 membered heterocyclic group that may have 1 to 3 substituents selected from the following substituent group i.



Substituent group i: a halogen atom, a cyano group, an amino group, a carbamoyl group, a C<sub>1-6</sub> alkyl group, a C<sub>3-8</sub> cycloalkyl group, a C<sub>1-6</sub> alkoxy group, a C<sub>1-6</sub> alkylcarbonyl group, a C<sub>1-6</sub> alkoxy-carbonyl group, a C<sub>1-6</sub> alkylsulfonyl group, a mono-C<sub>6-10</sub> arylamino group, a trifluoromethyl group and a trifluoromethoxy group]

5           Compound (1c-10) which is a commercially available product can be used as is. Compound (1c-10-1) which is a commercially available product can be used as is, or it can also be prepared from a commercially available product by a well known method.


10   [Step 1-15]

          The present step is a step wherein compound (1c-10) and compound (1c-10-1) are reacted to obtain compound (1c-11). Compound (1c-11) can be obtained by reacting compound (1c-10) and compound (1c-10-1) in a solvent such as, for instance, N,N-dimethylformamide, dimethylsulfoxide or the like, in the presence of  
15   a base such as, for instance, sodium hydride, potassium carbonate, cesium carbonate or the like. Compound (1c-10-1) is used in the amounts of 1 equivalent to 2 equivalents based on compound (1c-10). The base is used in the amounts of 1 equivalent to 2 equivalents based on compound (1c-10). The reaction temperature is from 0°C to 80°C, and the reaction time is from 5 minutes to 1 hour.

20

[Preparation Method 1-2-3] Preparation method for compound (1c-13), which is compound (1c-1)

[0055]




[wherein V represents a C<sub>1-6</sub> alkyl group, a C<sub>2-6</sub> alkenyl group, a C<sub>2-6</sub> alkynyl group, a C<sub>3-8</sub> cycloalkyl group, a C<sub>6-10</sub> aryl group, a 5- to 10-membered heterocyclic group, a C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkyl group, a C<sub>6-10</sub> aryl C<sub>1-6</sub> alkyl group or a 5- to 10-membered heterocycle C<sub>1-6</sub> alkyl group; W<sup>a</sup> represents a single bond or an oxygen atom]

Compound (1c-12) which is a commercially available product can be used as is. Compound (1c-12-1) which is a commercially available product can be used as is, or it can also be prepared from a commercially available product by a well known method.

#### [Step 1-16]

The present step is a step wherein compound (1c-12) and compound (1c-12-1) are reacted to obtain compound (1c-13). Examples of the solvent include, for instance, acetic acid or the like. Compound (1c-12-1) is used in the amount of 1 equivalent based on compound (1c-12). The reaction temperature is from 50°C to 110°C, and the reaction time is from 5 minutes to 1 hour.

[Preparation Method 1-2-4] Preparation method for compound (1c-15), which is compound (1c-7)



[0058]

[wherein Q and Ar<sup>1</sup> have the same meanings as defined above.]


Compound (1c-14) which is a commercially available product can be used as is. Compound (1c-10-1) which is a commercially available product can be used as is, or it can also be prepared from a commercially available product by a well known method.

[Step 1-17]

The present step is a step wherein compound (1c-14) and compound (1c-10-1) are reacted to obtain compound (1c-15). Compound (1c-15) can be obtained by reacting compound (1c-14) and compound (1c-10-1) in a solvent such as, for instance, N,N-dimethylformamide, dimethylsulfoxide or the like, in the presence of a base such as, for instance, sodium hydride, potassium carbonate, cesium carbonate or the like. Compound (1c-10-1) is used in the amounts of 1 equivalent to 2 equivalents based on compound (1c-14). The base is used in the amounts of 2 equivalent to 3 equivalents based on compound (1c-14). The reaction temperature is from room temperature to 80°C, and the reaction time is from 1 hour to 72 hours.

[Preparation Method 1-2-5] Preparation method for compound (1c-17), compound (1c-19), compound (1c-21), compound (1c-23) and compound (1c-25), which are compound (1c-7)

[0059]



[0060]

[wherein V and Ar<sup>1</sup> have the same meanings as defined above; R<sup>w1</sup> represents a hydrogen atom or a C<sub>1-6</sub> alkyl group; Hal represents a chlorine atom, a bromine atom or an iodine atom.]

Compound (1c-16), compound (1c-18), compound (1c-20), compound (1c-22), compound (1c-24), compound (1c-16-1), compound (1c-10-1), compound (1c-20-1) and compound (1c-22-1) which are commercially available products can be used as is, or they can also be prepared from commercially available products by a well known method.

## [Step 1-18]

The present step is a step wherein compound (1c-16) and compound (1c-16-1) are subjected to a coupling reaction to obtain compound (1c-17).

- 5 Compound (1c-17) can be obtained by reacting compound (1c-16) and compound (1c-16-1) in a solvent such as, for instance, tetrahydrofuran, 1,4-dioxane, methanol, ethanol, toluene, N,N-dimethylformamide or the like, in the presence of a base such as, for instance, potassium carbonate, cesium carbonate, potassium phosphate or the like, and a catalyst such as, for instance, palladium(II) acetate, 10 tetrakis(triphenylphosphine)palladium(0), dichlorobis(triphenylphosphine)palladium(II), tris(dibenzylideneacetone)dipalladium(0), dichloro(1,1'-bis(diphenylphosphino)ferrocene)nickel(II) or the like. The base is used in the amounts of 1.5 equivalents to excess amount based on compound (1c-16). The catalyst is used in the amounts of 0.05 equivalents to 0.3 equivalents 15 based on compound (1c-16).

## [Step 1-19]

- The present step is a step wherein compound (1c-18) and compound (1c-10-1) are reacted to obtain compound (1c-19). Compound (1c-19) can be 20 obtained by reacting compound (1c-18) and compound (1c-10-1) in a solvent such as dimethylsulfoxide or the like, in the presence of a base such as potassium *tert*-butoxide or the like. The base is used in the amounts of 1 equivalent to excess amount based on compound (1c-18). The reaction temperature is from 80°C to 220°C, and the reaction time is from 30 minutes to 48 hours.

[Step 1-20]

The present step is a step wherein compound (1c-20) and compound (1c-20-1) are reacted to carry out reductive amination and to obtain compound (1c-21). Compound (1c-21) can be obtained by reacting compound (1c-20) and  
5 compound (1c-20-1) in a solvent such as, for instance, tetrahydrofuran, methanol, ethanol or the like, in the presence of a reducing agent such as, for instance, lithium aluminum hydride, sodium borohydride, sodium cyanoborohydride, triacetoxy sodium borohydride or the like, and acetic acid. The reducing agent is used in the amounts of 1 equivalent to 2 equivalents based on compound (1c-20). The reaction  
10 temperature is from room temperature to 60°C, and the reaction time is from 10 minutes to 24 hours.

[Step 1-21]

The present step is a step wherein compound (1c-22) and compound  
15 (1c-22-1) are reacted to carry out reductive amination and to obtain compound (1c-23). Compound (1c-23) can be prepared by a method similar to [Step 1-20].

[Step 1-22]


The present step is a step wherein compound (1c-24) and compound  
20 (1c-10-1) are reacted to obtain compound (1c-25). Compound (1c-25) can be obtained by reacting compound (1c-24) and compound (1c-10-1) in a solvent such as, for instance, dimethylsulfoxide or the like, and in the presence of a base such as, for instance, potassium carbonate, cesium carbonate or the like. The base is used in the amounts of 1 equivalent to 3 equivalents based on compound (1c-24). The  
25 reaction temperature is from room temperature to 80°C, and the reaction time is from

10 minutes to 24 hours.

[Preparation Method 1-2-6] Preparation method for compound (1c-27), which is compound (1c-9), and compound (1c-29), which is compound (1c-1)

5

[0061]



[0062]

[wherein R<sup>w1</sup>, V, Q and Hal have the same meanings as defined above; p1 is an integer of 1 or 2; T represents a hydrogen atom when Q represents an oxygen atom, and a bromine atom when Q represents a sulfur atom.]

10

Compound (1c-26) which is a commercially available product can be used as is. Compound (1c-26-1) and compound (1c-22-1) which are commercially available products can be used as is, or they can also be prepared from commercially available products by a well known method.

15

[Step 1-23]

The present step is a step wherein compound (1c-26) and compound (1c-26-1) are reacted to obtain compound (1c-27). Compound (1c-27) can be

obtained by action of 1 equivalent of a strong base based on compound (1c-26) for anionization, then, reacting with compound (1c-26-1). Examples of the solvent include, for instance, tetrahydrofuran, diethyl ether or the like. Examples of the strong base include, for instance, *n*-butyl lithium or the like. The reaction  
5 temperature is from -80°C to room temperature.

[Step 1-24]

The present step is a step wherein compound (1c-26) and compound (1c-22-1) are reacted to obtain compound (1c-28). Compound (1c-28) can be  
10 obtained by action of 1 equivalent of a strong base based on compound (1c-26) for anionization, then, reacting with compound (1c-22-1). Examples of the solvent include, for instance, tetrahydrofuran, diethyl ether or the like. Examples of the strong base include, for instance, *n*-butyl lithium or the like. In addition, 0.1 equivalent to 1 equivalent of copper(I) iodide or copper(I) bromide based on  
15 compound (1c-26) may be added in the reaction system. The reaction temperature is from -80°C to room temperature.

[Step 1-25]

The present step is a step wherein elimination of a hydroxyl group and  
20 deprotection of an acetal group of compound (1c-28) are carried out simultaneously to obtain compound (1c-29). When compound (1c-28) is treated with trimethylsilyliodide, elimination of the hydroxyl group and deprotection of the acetal group occur simultaneously, to give compound (1c-29). Trimethylsilyliodide is used in the amounts of 2 equivalents to 6 equivalents based on compound (1c-28). In  
25 addition, trimethylsilyliodide may be prepared from trimethylsilyl chloride and sodium




iodide in the reaction solution and used as is. Examples of the solvent include, for instance, acetonitrile or the like. The reaction temperature is from 0°C to 60°C, and the reaction time is from 5 minutes to 6 hours.

Other method: Compound (1c-29) can be obtained by converting the hydroxyl group of compound (1c-28) into the acetyl group, then, eliminating the acetyl group, and carrying out deprotection of an acetal group. Conversion of the hydroxyl group into the acetyl group can be carried out by using an acetylating reagent such as, for instance, acetic anhydride, acetyl chloride or the like. A base such as, for instance, N,N-dimethylaminopyridine, triethylamine, pyridine or the like, in the amounts of 1 equivalent to excess amount based on compound (1c-28) may be added in the reaction system. Examples of the solvent include, for instance, dichloromethane, ethyl acetate, N,N-dimethylformamide or the like. In addition, pyridine, which is added as a base, may be used directly as a solvent. An acetylating reagent is used in the amounts of 1 equivalent to excess amount based on compound (1c-28). The reaction temperature is from 0°C to 60°C, and the reaction time is from 1 hour to 36 hours. Elimination of the acetyl group can be carried out, for instance, under hydrogen atmosphere, in a solvent such as, for instance, ethanol, methanol or the like, and using a catalyst such as for instance, palladium-carbon, Raney nickel or the like. The reaction temperature is from room temperature to 80°C, and the reaction time is from 5 hours to 36 hours. Deprotection of the acetal group can be carried out by an analogous method to [Step 1-13].

[Preparation Method 2] Representative preparation method for compound (2a)

[0063]




[0064]

[wherein each symbol has the same meaning as defined above.]

## 5 [Preparation Method 2-1] Amidation

[0065]



[0066]

[where each symbol has the same meaning as defined above.]

10 Compound (2b) which is a commercially available product can be used as is, or it can also be prepared from a commercially available product by a well known method. In addition, it can also be prepared using a method described in the preparation examples among the examples.


Compound (2c) which is a commercially available product can be used as is,  
 15 or it can also be prepared from a commercially available product by a well known method. In addition, it can also be prepared by a method described in the preparation examples among the examples or [Preparation Method 2-2-1] or the like.

[Step 2]

20 Compound (2a) is obtained by condensing compound (2b) and compound (2c) in a solvent using a condensing agent. Compound (2a) can be prepared by an analogous method to [Step 1].

[Preparation Method 2-2-1] Preparation method for compound (2c-2), which is compound (2c), and compound (2c-5), which is compound (2c)

[0067]



5

[0068]

[wherein E has the same meaning as defined above; p2 represents an integer from 0 to 2.]

Compound (2c-1) which is a commercially available product can be used as is, or it can also be prepared from a commercially available product by a well known method.

10

[Step 2-1]

The present step is a step wherein compound (2c-1) is oxidized to obtain compound (2c-2). Compound (2c-2) can be obtained by dissolving compound (2c-1) in a mixed solvent of *tert*-butanol and water (preferable mixing ratio being *tert*-butanol:water=10:1 to 2:1), and reacting with 3 equivalents to 10 equivalents of 2-methyl-2-butene, 1 equivalent to 2 equivalents of sodium chlorite and 1 equivalent to 2 equivalents of sodium dihydrogenphosphate based on compound (2c-1). The reaction temperature is the room temperature, and the reaction time is from 10 minutes to 2 hours.

20

In addition, compound (2c-2) can also be prepared compound (2c-1) by a method described in the following other method (1), (2) or (3).

Other method (1): Compound (2c-2) can be obtained by dissolving compound (2c-1) in a mixed solvent of *tert*-butanol and water (preferable mixing ratio being *tert*-butanol:water=10:1 to 2:1), and reacting with 1 equivalent to 2 equivalents of potassium permanganate and 1 equivalent to 2 equivalents of potassium dihydrogenphosphate based on compound (2c-1). The reaction temperature is room temperature, and the reaction time is from 10 minutes to 2 hours.

Other method (2): Compound (2c-2) can be obtained by dissolving compound (2c-1) in a sodium hydroxide aqueous solution and ethanol, and reacting with 1.5 equivalents to 2 equivalents of silver nitrate based on compound (2c-1). The reaction temperature is from 0°C to room temperature, and the reaction time is from 5 minutes to 24 hours.

Other method (3): Compound (2c-2) can also be obtained by dissolving compound (2c-1) in methanol and N,N-dimethylformamide, and reacting with 1.0 equivalent to 1.5 equivalents of pyridinium dichromate (PDC) based on compound (2c-1). The reaction temperature is from -10°C to 40°C, and the reaction time is from 10 minutes to 24 hours.

## [Step 2-2]

The present step is a step wherein compound (2c-1) is extended by one carbon to obtain compound (2c-3). Compound (2c-3) can be obtained in a solvent such as, for instance, tetrahydrofuran, 1,4-dioxane or the like, at 10°C to room temperature, by reacting 2 equivalents to 10 equivalents of methoxymethyl triphenylphosphonium chloride and 2 equivalents to 10 equivalents of base (for

instance, potassium *tert*-butoxide) based on compound (2c-1) for 30 minutes to 2 hours, then adding compound (2c-1) to this reaction solution, and reacting at room temperature to reflux temperature for 30 minutes to 4 hours.

5 [Step 2-3]


The present step is a step wherein compound (2c-3) is hydrolyzed to obtain compound (2c-4). Compound (2c-4) can be obtained by dissolving compound (2c-3) in a solvent such as, for instance, hydrous methanol, hydrous 1,4-dioxane or the like, adding an acid such as, for instance, hydrochloric acid, *p*-toluenesulfonic  
10 acid or the like, and reacting at 80°C to reflux temperature for 10 minutes to 16 hours.

[Step 2-4]

The present step is a step wherein compound (2c-4) is oxidized to obtain  
15 compound (2c-5). Compound (2c-5) can be prepared by an analogous method to [Step 2-1].

[Preparation Method 2-2-2] Preparation method for compound (2c-7), which is compound (2c)

20



[wherein E and n have the same meanings as defined above.]

Compound (2c-6) which is a commercially available product can be used as


is, or it can also be prepared from a commercially available product by a well known method.

[Step 2-5]

- 5           The present step is a step wherein compound (2c-6) is hydrolyzed to obtain compound (2c-7). Compound (2c-7) can be obtained in an aqueous solution of acid such as, for instance, hydrochloric acid and sulfuric acid or the like, or, in an aqueous solution of alkali such as, for instance, sodium hydroxide, potassium hydroxide or the like, by hydrolyzing compound (2c-6). The reaction temperature is from room
- 10   temperature to reflux temperature, and the reaction time is from 10 minutes to 10 hours.

[Preparation Method 3] Representative preparation method for compound (3a), compound (3b), compound (3c), compound (3d) and compound (3e)


15           [0071]



[0072]

[wherein each symbol has the same meaning as defined above.]

- 20   [Preparation Method 3-1] Representative preparation method for compound (3a) and compound (3b)



[wherein A, R<sup>Y</sup>, n, E and Hal have the same meanings as defined above; G<sup>1</sup>

5 represent a C<sub>1-6</sub> alkyl group that may be substituted with a C<sub>6-10</sub> aryl group.]

#### [Step 3-1]

The present process is a process wherein compound (1a) is converted into thioamide to obtain compound (3a). Compound (3a) can be obtained by reacting  
 10 compound (1a) and 2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulfide (Lawesson's reagent). Examples of the solvent include, for instance, tetrahydrofuran, toluene or the like. Lawesson's reagent is used in the amounts of 1 equivalent to 5 equivalents based on compound (1a). The reaction temperature is from room temperature to reflux temperature, and the reaction time is from 10  
 15 minutes to 48 hours.

#### [Step 3-2]

The present step is a step wherein compound (3a) is converted into thioimide to obtain compound (3a-2). Compound (3a-2) can be obtained by  
 20 reacting compound (3a) and compound (3a-1). Examples of the solvent include, for instance, acetonitrile, tetrahydrofuran, toluene or the like. For compound (3a-1),

preferably, alkylating agents such as, for instance, iodomethane, benzyl bromide, (bromomethyl)naphthalene or the like can be used. Compound (3a-1) is used in the amounts of 1 equivalent to 10 equivalents based on compound (3a). The reaction temperature is from room temperature to reflux temperature, and the reaction time is

5 from 10 minutes to 48 hours.

## [Step 3-3]


The present step is a step wherein compound (3a-2) is converted into amidine to obtain compound (3b). Compound (3b) can be obtained by reacting

10 compound (3a-2) and compound (3a-3). Examples of the solvent include, for instance, methanol, ethanol, N,N-dimethylformamide, N-methylpyrrolidinone or the like. Examples of compound (3a-3) include, preferably, alkoxy amines such as, for instance, methoxylamine, benzyloxy amine, cyanamide or the like. Compound (3a-3) is used in the amounts of 1 equivalent to 10 equivalents based on compound

15 (3a-2). The reaction temperature is from room temperature to reflux temperature, and the reaction time is from 10 minutes to 48 hours.

## [Preparation Method 3-2] Representative preparation method for compound (3c)

[0075]



[0076]



[wherein each symbol has the same meaning as defined above.]

Compound (3c-1) can be prepared using the method described in [Preparation Method 1-1]. Compound (3c-2) and compound (3c-3) which are commercially available products can be used as is, or they can also be prepared from  
5 commercially available products by a well known method.

[Step 3-4]

The present step is a step wherein compound (3c-1) is reduced to obtain compound (3c). Compound (3c) can be obtained by reacting compound (3c-1) in a  
10 solvent such as, for instance, tetrahydrofuran or the like, in the presence of a reducing agent such as, for instance, lithium aluminum hydride or the like. The reducing agent is used in the amounts of 1 equivalent to 5 equivalents based on compound (3c-1). The reaction temperature is from room temperature to reflux temperature, and the reaction time is from 1 hour to 24 hours.

15


[Step 3-5]

The present step is a step wherein compound (3c-2) and compound (3c-3) are reacted to carry out reductive amination and obtain compound (3c). Compound (3c) can be prepared by an analogous method to [Step 1-20].

20

[Preparation Method 3-3] Representative preparation method for compound (3d)

[0077]



[wherein each symbol has the same meaning as defined above.]

Compound (3d-1) which is a commercially available product can be used as  
 5 is, or it can also be prepared from a commercially available product by a well known method.

#### [Step 3-6]


The present step is a step wherein compound (2a) is reduced to obtain  
 10 compound (3d). Compound (3d) can be prepared by an analogous method to [Step 3-4].

#### [Step 3-7]

The present step is a step wherein compound (2b) and compound (3d-1) are  
 15 reacted to carry out reductive amination and obtain compound (3d). Compound (3d) can be prepared by an analogous method to [Step 1-20].

[Preparation Method 3-3-1] Preparation method for compound (3d-3), which is compound (3d)

20 [0079]




[wherein each symbol has the same meaning as defined above.]

### 5 [Step 3-8]

The present step is a step wherein compound (3d-2) and cyanamide are reacted to obtain compound (3d-3). Examples of the solvent include, for instance, N,N-dimethylformamide, N-methylpyrrolidinone or the like. Cyanamide is used in the amounts of 2 equivalent to 5 equivalents based on compound (3d-2). The reaction temperature is from room temperature to reflux temperature, and the reaction time is from 10 minutes to 48 hours.

[Preparation Method 3-4] Representative preparation method for compound (3e)

[0081]



[0082]

[wherein each symbol has the same meaning as defined above.]

Compound (3e-1), compound (3c-2) and compound (3e-3) which are commercially available products can be used as is, or they can also be prepared from  
5 commercially available products by a well known method.

[Step 3-9]

The present step is a step wherein compound (3e-1) is converted into Grignard reagent, then, the Grignard reagent and compound (3c-2) are reacted to  
10 obtain compound (3e-2). Compound (3e-2) can be obtained by preparing the Grignard reagent of compound (3e-1) in a solvent such as, for instance, tetrahydrofuran or the like, in the presence of metallic magnesium and an initiator such as, for instance, dibromoethane or the like, and reacting the Grignard reagent and compound (3c-2). Metallic magnesium is used in the amounts of 1 equivalent to  
15 1.2 equivalents based on compound (3e-1). The initiator is used in catalytic amounts based on compound (3e-1). The reaction temperature is from 0°C to reflux temperature, and the reaction time is from 10 minutes to 24 hours.

[Step 3-10]

20 The present step is a step wherein a hydroxyl group of compound (3e-2) is oxidized to obtain compound (3e). Compound (3e) can be obtained by reacting compound (3e-2) in a solvent such as, for instance, chloroform, acetone or the like, in the presence of an oxidizing agent such as, for instance, manganese dioxide or the like. The oxidizing agent is used in the amounts of 2 equivalents to 10 equivalents  
25 based on compound (3e-2). The reaction temperature is from room temperature to

reflux temperature, and the reaction time is from 10 minutes to 24 hours.


[Step 3-11]

The present step is a step wherein compound (3e-1) is converted into  
5 Grignard reagent, then, the Grignard reagent and compound (3e-3) are reacted, and  
hydrolysis is carried out to obtain compound (3e). Compound (3e) can be obtained  
by preparing the Grignard reagent of compound (3e-1) in a solvent such as, for  
instance, tetrahydrofuran or the like, in the presence of metallic magnesium and an  
10 initiator such as, for instance, dibromoethane or the like, reacting the Grignard  
reagent and compound (3c-2), and hydrolyzing in the presence of acid. Metallic  
magnesium is used in the amounts of 1 equivalent to 1.2 equivalents based on  
compound (3e-1). The initiator is used in catalytic amounts based on compound  
(3e-1). The reaction temperature is from 0°C to reflux temperature, and the reaction  
time is from 10 minutes to 24 hours.

15

[Preparation Method 4-1] Conversion of substituent on A in compound (I) (with the  
proviso that A represents an aromatic 5- and 10-membered heterocyclic group  
containing at least 1 nitrogen atom) - 1

[0083]



[0084]

[wherein A, X (with the proviso that the case where X represents a group represented

by the formula  $-\text{CH}_2\text{-NH-(CH}_2\text{)}_n\text{-}$  or a group represented by the formula  $-\text{NH-CH}_2\text{-(CH}_2\text{)}_n\text{-}$  are excluded;  $n$  represents an integer from 0 to 3),  $E$  and  $\text{Hal}$  have the same meanings as defined above;  $\text{R}^{\text{N}1}$  represents a  $\text{C}_{1-6}$  alkyl group, a  $\text{C}_{2-6}$  alkenyl group or a  $\text{C}_{2-6}$  alkynyl group;  $\text{R}^{\text{N}2}$  represents a hydrogen atom or a  $\text{C}_{1-6}$  alkyl group;  $\text{R}^{\text{a}}$  represents a  $\text{C}_{2-6}$  alkenyl group, a  $\text{C}_{6-10}$  aryl group or an aromatic 5- and 10-membered heterocyclic group;  $\text{R}^{\text{b}1}$  and  $\text{R}^{\text{b}2}$  may be the same or different from each other, and represent a hydrogen atom or a  $\text{C}_{1-6}$  alkyl group, or form a cyclic boric acid ester together;  $\text{R}^{\text{c}1}$ ,  $\text{R}^{\text{c}2}$ ,  $\text{R}^{\text{c}3}$  and  $\text{R}^{\text{c}4}$  may be the same or different from each other, and represent a hydrogen atom, a  $\text{C}_{1-6}$  alkyl group, a  $\text{C}_{2-6}$  alkenyl group, a  $\text{C}_{2-6}$  alkynyl group, a  $\text{C}_{3-8}$  cycloalkyl group, a  $\text{C}_{6-10}$  aryl group or a 5- to 10-membered heterocyclic group.]

Compound (4-1-1), compound (4-1-2), compound (4-1-3), compound (4-1-4), compound (4-1-5) and compound (4-1-6) which are commercially available products can be used as is, or they can also be prepared from commercially available products by a well known method.

#### [Step 4-1]

The present step is a step wherein compound (4-1) and compound (4-1-1) are reacted to obtain compound (4-2). Examples of the solvent include, for instance, dimethylsulfoxide, tetrahydrofuran, toluene, acetonitrile,  $\text{N,N}$ -dimethylformamide or the like. In addition, the reaction can also be carried out without the solvent. The reaction is preferably carried out in a sealed tube; the reaction time is from 1 hour to 60 hours, and the reaction temperature is from  $50^\circ\text{C}$  to  $200^\circ\text{C}$ . Note that, an organic base such as, for instance,  $\text{N,N}$ -diisopropylethylamine, triethylamine, pyridine, 1,8-diazabicyclo[5,4,0]undec-7-ene, inorganic base such as, for instance, potassium

carbonate, sodium carbonate, can be added in the amounts of 2 equivalents to excess amount based on compound (4-1).

[Step 4-2]

5           The present step is a step wherein compound (4-1) and compound (4-1-2) are reacted to obtain compound (4-3). Compound (4-3) can be obtained by reacting compound (4-1) and compound (4-1-2) in the presence of a catalyst. Examples of the catalyst include tetrakis(triphenylphosphine)palladium(0), dichlorobis(triphenylphosphine)palladium(II) or the like. Examples of the solvent  
10 include, toluene, 1,4-dioxane, xylene or the like. Compound (4-1-2) is used in the amounts of 2 equivalent to 3 equivalents based on compound (4-1). The catalyst is used in the amounts of 0.05 equivalents to 0.3 equivalents based on compound (4-1). The reaction temperature is from 100°C to 140°C, and the reaction time is from 1 hour to 24 hours.

15

[Step 4-3]

          The present step is a step wherein compound (4-1) and compound (4-1-3) are reacted to obtain compound (4-3). Compound (4-3) can be obtained by reacting compound (4-1) and compound (4-1-3) in the presence of a catalyst. Examples of  
20 the catalyst include, for instance, dichloro(1,1'-bis(diphenylphosphino)propane)nickel(II), dichloro(1,1'-bis(diphenylphosphino)ferrocene)nickel(II), tetrakis(triphenylphosphine)palladium(0) or the like. Examples of the solvent include, for instance, tetrahydrofuran, 1,4-dioxane or the like. Compound (4-1-3) is used in  
25 the amounts of 3 equivalents to excess amount based on compound (4-1). The



catalyst is used in the amounts of 0.05 equivalents to 0.3 equivalents based on compound (4-1). The reaction temperature is from 0°C to reflux temperature, and the reaction time is from 5 minutes to 24 hours.

5 [Step 4-4]

The present step is a step wherein compound (4-1) and compound (4-1-4) are reacted to obtain compound (4-3). Compound (4-3) can be obtained by reacting compound (4-1) and compound (4-1-4) in the presence of a catalyst and a base. Examples of the catalyst include, for instance, palladium(II) acetate, 10 tetrakis(triphenylphosphine)palladium(0), dichlorobis(triphenylphosphine)palladium(II), tris(dibenzylideneacetone) dipalladium(0) or the like. To obtain satisfactory results, a phosphorus ligand (for instance, triphenylphosphine, tri-*tert*-butylphosphine or the like) may be added in the amounts of 0.25 equivalents to 1.5 equivalents based on compound (4-1). Examples of the basic include, for instance, potassium carbonate, 15 sodium carbonate, cesium carbonate, potassium fluoride, cesium fluoride, potassium phosphate, sodium hydroxide, barium hydroxide, potassium hydroxide or the like. The present reaction is preferably carried out under an inert gas atmosphere such as, for instance, nitrogen gas and argon gas. Examples of the solvent include, for instance, tetrahydrofuran, 1,4-dioxane, methanol, ethanol, toluene, water or the like. 20 Depending on the reagent used, quaternary ammonium salt such as tetrabutylammonium bromide can be added. The catalyst is used in the amounts of 0.05 equivalents to 0.3 equivalents based on compound (4-1). The base is used in the amounts of 2 equivalents to excess amount based on compound (4-1). The reaction temperature is from room temperature to reflux temperature, and the 25 reaction time is from 30 minutes to 24 hours.

## [Step 4-5]

The present step is a step wherein compound (4-1) and compound (4-1-5) are reacted to obtain compound (4-4). Compound (4-4) can be obtained by reacting  
5 compound (4-1) and compound (4-1-5) in the presence of a catalyst and a base. Examples of the catalyst include, for instance, palladium(II) acetate, tetrakis(triphenylphosphine)palladium(0), dichlorobis(triphenylphosphine)palladium(II), tris(dibenzylideneacetone) dipalladium (0) or the like. Examples of the base include, for instance, triethylamine, N,N-diisopropylethylamine, pyridine or the like.  
10 Examples of the solvent include, for instance, tetrahydrofuran, acetonitrile, 1,4-dioxane, N,N-dimethylformamide, N-methylpyrrolidinone, dimethylsulfoxide, toluene or the like. In addition, to obtain satisfactory results, 0.1 equivalents to 0.3 equivalents of copper (I) iodide or tetrabutylammonium fluoride may be added based on compound (4-1). Compound (4-1-5) is used in the amounts of 1 equivalent to 5  
15 equivalents based on compound (4-1). The catalyst is used in the amounts of 0.05 equivalents to 0.3 equivalents based on compound (4-1). The base is used in the amounts of 2 equivalents to 5 equivalents based on compound (4-1). The reaction temperature is from room temperature to 150°C, and the reaction time is from 30 minutes to 24 hours.

20

## [Step 4-6]

The present step is a step wherein a triple bond in compound (4-4) is reduced into a double bond to obtain compound (4-5). Compound (4-5) can be obtained using a catalyst such as, for instance, Lindlar catalyst, palladium-barium  
25 sulfate or the like, in a solvent such as, for instance, tetrahydrofuran, ethyl acetate,

acetonitrile, methanol, ethanol or the like, under hydrogen atmosphere. The preferable solvent is ethyl acetate. To obtain satisfactory results, 0.1 equivalents to 1 equivalents of quinoline may be added based on compound (4-4). The catalyst is used in catalytic amount to excess amount based on compound (4-4). The reaction  
5 temperature is room temperature, the reaction time is from 15 minutes to 24 hours, and reaction pressure is from 1 atm to 4 atm.

[Step 4-7]

The present step is a step wherein compound (4-5) is reduced to obtain  
10 compound (4-6). Compound (4-6) can be obtained using a catalyst such as, for instance, palladium-carbon, Raney nickel, platinum dioxide or the like, in a solvent such as, for instance, tetrahydrofuran, ethyl acetate, acetonitrile, methanol, ethanol or the like, under hydrogen atmosphere. The catalyst is used in catalytic amount to excess amount based on compound (4-5). The reaction temperature is room  
15 temperature, the reaction time is from 5 minutes to 24 hours, and the reaction pressure is from 1 atm to 4 atm.

[Step 4-8]

The present step is a step wherein compound (4-4) is reduced to obtain  
20 compound (4-6). Compound (4-6) can be prepared by an analogous method to [Step 4-7].

[Step 4-9]

The present step is a step wherein compound (4-1) and compound (4-1-6)  
25 are reacted to obtain compound (4-7). Compound (4-7) can be obtained by reacting


compound (4-1) and compound (4-1-6) in the presence of a catalyst and a base. Examples of the catalyst include, for instance, palladium(II) acetate, tetrakis(triphenylphosphine)palladium(0), dichlorobis(triphenylphosphine)palladium(II), tris(dibenzylideneacetone)dipalladium (0) or the like. Examples of the base include, for instance, triethylamine, N,N-diisopropylethylamine, N,N-dicyclohexylmethylethylamine or the like. Examples of the solvent include, for instance, acetonitrile, tetrahydrofuran, 1,4-dioxane, benzene, toluene, xylene, N,N-dimethylformamide, N-methylpyrrolidinone or the like. In addition, to obtain satisfactory results, 0.25 equivalents to 1.5 equivalents of a phosphorus ligand (for instance, triphenylphosphine, tri-*tert*-butylphosphine, 2-(di-*tert*-butylphosphino)biphenyl or the like) may be added based on compound (4-1). Compound (4-1-6) is used in the amounts of 1 equivalent to 4 equivalents based on compound (4-1). The catalyst is used in the amounts of 0.05 equivalents to 0.3 equivalents based on compound (4-1). The base is used in the amounts of 2 equivalents to 5 equivalents based on compound (4-1). The reaction temperature is from room temperature to 150°C, the reaction time is from 5 minutes to 24 hours.

#### [Step 4-10]

The present step is a step wherein compound (4-7) is reduced to obtain compound (4-8). Compound (4-8) can be prepared by an analogous method to [Step 4-7].

#### [Preparation Method 4-2] Conversion of substituent on E in compound (I) -1

[0085]



[0086]

[wherein A, X (with the proviso that the case where X represents a group represented by the formula  $-\text{CH}_2\text{-NH-(CH}_2\text{)}_n-$  or a group represented by the formula  $-\text{NH-CH}_2\text{-(CH}_2\text{)}_n-$  are excluded; n represents an integer from 0 to 3), E, Hal,  $\text{R}^a$ ,  $\text{R}^{b1}$ ,  $\text{R}^{b2}$ ,  $\text{R}^{c1}$ ,  $\text{R}^{c2}$ ,  $\text{R}^{c3}$ ,  $\text{R}^{c4}$ ,  $\text{R}^{w1}$ , V and  $\text{Ar}^1$  have the same meanings as defined above.]

Compound (4-1-2), compound (4-1-3), compound (4-1-4), compound (4-1-5), compound (4-1-6), compound (1c-20-1) and compound (1c-10-1) which are commercially available products can be used as is, or they can also be prepared from commercially available products by a well known method.

[Step 4-11]

The present step is a step wherein compound (4-9) and compound (4-1-2) are reacted to obtain compound (4-10). Compound (4-10) can be prepared by an analogous method to [Step 4-2].

[Step 4-12]

The present step is a step wherein compound (4-9) and compound (4-1-3) are reacted to obtain compound (4-10). Compound (4-10) can be prepared by an analogous method to [Step 4-3].

[Step 4-13]

The present step is a step wherein compound (4-9) and compound (4-1-4) are reacted to obtain compound (4-10). Compound (4-10) can be prepared by an analogous method to [Step 4-4].

[Step 4-14]

The present step is a step wherein compound (4-9) and compound (4-1-5) are reacted to obtain compound (4-11). Compound (4-11) can be prepared by an analogous method to [Step 4-5].

[Step 4-15]

The present step is a step wherein a triple bond of compound (4-11) is reduced into a double bond to obtain compound (4-12). Compound (4-12) can be prepared by an analogous method to [Step 4-6].

[Step 4-16]

The present step is a step wherein compound (4-12) is reduced to obtain compound (4-13). Compound (4-13) can be prepared by an analogous method to [Step 4-7].

[Step 4-17]

The present step is a step wherein compound (4-11) is reduced to obtain compound (4-13). Compound (4-13) can be prepared by an analogous method to [Step 4-8].

[Step 4-18]

The present step is a step wherein compound (4-9) and compound (4-1-6) are reacted to obtain compound (4-14). Compound (4-14) can be prepared by an analogous method to [Step 4-9].

[Step 4-19]

The present step is a step wherein compound (4-14) is reduced to obtain compound (4-15). Compound (4-15) can be prepared by an analogous method to

5 [Step 4-10].

[Step 4-20]

The present step is a step wherein compound (4-9) and compound (1c-20-1) are reacted to obtain compound (4-16). Compound (4-16) can be obtained by  
10 reacting compound (4-9) and compound (1c-20-1) in a solvent such as, for instance, tetrahydrofuran, benzene, toluene, xylene or the like, in the presence of a catalyst such as for instance, tris(dibenzylideneacetone)dipalladium(0), dichloro(1,1'-bis(diphenylphosphino)ferrocene)palladium(II), palladium(II) acetate, a phosphorus ligand such as, for instance, 2,2-bis(diphenylphosphino)-1,1'-binaphthyl,  
15 and a base such as, for instance, sodium *tert*-butoxide. Compound (1c-20-1) is used in the amounts of 1 equivalent to 3 equivalents based on compound (4-9). Catalyst is used in the amounts of 0.05 equivalents to 0.3 equivalents based on compound (4-9). The base is used in the amounts of 1.5 equivalent to excess amount based on compound (4-9). The phosphorus ligand is used in the amounts  
20 of 0.25 equivalents to 1.5 equivalents based on compound (4-9). The reaction temperature is 50°C to reflux temperature, and the reaction time is from 1 hour to 48 hours.

[Step 4-21]

25 The present step is a step wherein compound (4-9) and compound (1c-10-1)




are reacted to obtain compound (4-17). Compound (4-17) can be obtained by reacting compound (4-9) and compound (1c-10-1) in a solvent such as, for instance, tetrahydrofuran, toluene or the like, in the presence of a catalyst such as, for instance, copper(I) chloride, copper(I) iodide or the like, and a base such as, for instance, potassium carbonate, cesium carbonate, potassium phosphate, pyridine or the like. Compound (1c-10-1) is used in the amounts of 1 equivalent to 3 equivalents based on compound (4-9). Catalyst is used in the amounts of 0.5 equivalents to 3 equivalents based on compound (4-9). The base is used in the amounts of 2 equivalents to 10 equivalents based on compound (4-9). The reaction temperature is 50°C to reflux temperature, and the reaction time is from 1 hour to 48 hours.

Other method for [Step 4-20] and [Step 4-21]

When E represent a furyl group or a thienyl group, compound (4-16) or compound (4-17) can be obtained respectively by reacting compound (4-9) and compound (1c-20-1) or compound (1c-10-1) in a solvent such as, for instance, dimethylsulfoxide, N-methylpyrrolidone or the like, in the presence of a catalyst such as, for instance, copper(I) chloride or the like, a base such as, for instance, cesium carbonate and 2,2,6,6-tetramethyl-3,5-heptanedione. Compound (1c-20-1) or compound (1c-10-1) is used in the amounts of 1 equivalent to 3 equivalents based on compound (4-9). The catalyst is used in the amounts of 0.5 equivalents to 3 equivalents based on compound (4-9). The base is used in the amounts of 2 equivalents to 10 equivalents based on compound (4-9). The reaction temperature is 80°C to reflux temperature, and the reaction time is from 1 hour to 24 hours.

## [Preparation Method 4-3] Conversion of substituent on E in compound (I) -2

[0087]



[0088]

- 5 [wherein A, X (with the proviso that the cases where X represents a group represented by the formula  $\text{---CH}_2\text{---NH---}(\text{CH}_2)_n\text{---}$  or a group represented by the formula  $\text{---NH---CH}_2\text{---}(\text{CH}_2)_n\text{---}$  are excluded; n represents an integer from 0 to 3), E,  $\text{R}^a$ , Hal,  $\text{R}^{w1}$ ,  $p_1$  and V have the same meanings as defined above.]

Compound (4-18-1) and compound (1c-26-1) which are commercially  
 10 available products can be used as is, or they can also be prepared from commercially available products by a well known method.

## [Step 4-22]

The present step is a step wherein compound (4-18) and compound (4-18-1)  
 15 are reacted to obtain compound (4-19). Compound (4-19) can be obtained by reacting compound (4-18) and compound (4-18-1) in the presence of a catalyst and a base. Examples of the catalyst include a cuprous catalyst such as, for instance, copper (II) acetate. Examples of the base include, for instance, triethylamine, N,N-diisopropylethylamine or the like. Examples of the solvent include, for instance,

dichloromethane, tetrahydrofuran, toluene or the like. It is preferable to use dichloromethane. The present reaction is preferably carried out in the presence of oxygen. To obtain satisfactory results, molecular sieves 4A may be added. Compound (4-18-1) is used in the amounts of 1 equivalent to 4 equivalents based on compound (4-18). The catalyst is used in the amounts of 0.1 equivalents to 0.3 equivalents based on compound (4-18). The base is used in the amounts of 2 equivalents to excess amount based on compound (4-18). The reaction temperature is from room temperature to 50°C, the reaction time is from 24 hours to 5 days.

10

## [Step 4-23]

The present step is a step wherein compound (4-18) and compound (1c-26-1) are reacted to obtain compound (4-20). Compound (4-20) can be obtained by reacting compound (4-18) and compound (1c-26-1) in a solvent such as, for instance, N,N-dimethylformamide, N-methylpyrrolidinone, tetrahydrofuran or the like, in the presence of a base such as, for instance, potassium carbonate, cesium carbonate, sodium hydride or the like. To obtain satisfactory results, a catalytic amount of sodium iodide or potassium iodide may be added. The reaction temperature is from room temperature to 160°C, and the reaction time is from 10 minutes to 48 hours.


15  
20

Another method is based on the technique that uses the Mitsunobu reaction. Compound (4-20) can be obtained by reacting compound (4-18), compound (1c-26-1), triphenylphosphine and diethyl azodicarboxylate or diisopropyl azodicarboxylate in a solvent such as, for instance, dichloromethane, tetrahydrofuran or the like. Compound (1c-26-1) is used in the amounts of 1 equivalent to 1.5

25

equivalents based on compound (4-18). Triphenylphosphine is used in the amounts of 1 equivalent to 1.5 equivalents based on compound (4-18). Diethyl azodicarboxylate or diisopropyl azodicarboxylate is used in the amounts of 1 equivalent to 1.5 equivalents based on compound (4-18). The reaction temperature is from room temperature to reflux temperature, and the reaction time is from 5 minutes to 24 hours.

[Preparation Method 4-4] Conversion of substituent on A in compound (I) (with the proviso that A represents an aromatic 5- to 10-membered heterocyclic group containing at least 1 nitrogen atom) - 2



[wherein A, X (with the proviso that the cases where X represents a group represented by the formula  $-\text{CH}_2\text{-NH}-(\text{CH}_2)_n-$  or a group represented by the formula  $-\text{NH-CH}_2-(\text{CH}_2)_n-$  are excluded; n represents an integer from 0 to 3), E and Hal have the same meanings as defined above;  $\text{R}^d$  and  $\text{R}^e$  are the same or different from each other, and represent  $\text{C}_{1-6}$  alkyl groups.]

[Step 4-24]

The present step is a step wherein compound (4-1) and compound (4-24-1) are reacted to obtain compound (4-21). The present reaction is preferably carried out under an inert gas atmosphere; the solvent for use varies depending on the


starting materials and reagents used; for instance, N-methylpyrrolidinone, 1,4-dioxane or the like can be used. Examples of the catalyst include, for instance, palladium(II) acetate, tetrakis(triphenylphosphine)palladium(0), dichlorobis(triphenylphosphine)palladium(II) or

5 tris(dibenzylideneacetone)dipalladium(0) or the like. To obtain satisfactory results, of phosphorus ligand, preferably, for instance, triphenylphosphine, tri-*tert*-butylphosphine, diphenylphosphino ferrocene or the like may be added. Compound (4-24-1) is used in the amounts of 1 equivalent to 10 equivalents based on compound (4-1). The catalyst is used in the amounts of 0.001 equivalents to 0.2

10 equivalents based on compound (4-1). The phosphorus ligand is used in the amounts of 0.001 equivalents to 0.4 equivalents based on compound (4-1). The reaction temperature is from room temperature to reflux temperature, and the reaction time is from 10 minutes to 24 hours.

15 [Preparation Method 4-5] Conversion of substituent on E in compound (I) -3

[0091]



[0092]

[wherein A, X (with the proviso that the cases where X represents a group represented by the formula -CH<sub>2</sub>-NH-(CH<sub>2</sub>)<sub>n</sub>- or a group represented by the formula -NH-CH<sub>2</sub>-(CH<sub>2</sub>)<sub>n</sub>- are excluded; n represents an integer from 0 to 3), E, Hal, R<sup>d</sup> and R<sup>e</sup> have the same meanings as defined above.]

20


## [Step 4-25]

The present step is a step wherein compound (4-9) and compound (4-24-1) are reacted to obtain compound (4-22). Compound (4-22) can be prepared by an analogous method to [Step 4-24].

5

(Preparation method for compound (4-24-1))

[0093]



[0094]

10 [wherein L,  $\text{R}^d$  and  $\text{R}^e$  have the same meanings as defined above.]

Compound (4-24-2), compound (4-24-5) and compound (4-24-6) which are commercially available products can be used as is, or they can also be prepared from commercially available products by a well known method.

15 [Step 4-24-1]

The present step is a step wherein compound (4-24-2) and compound (4-24-5) are reacted to obtain compound (4-24-1). Compound (4-24-1) can be obtained in a solvent such as, for instance, tetrahydrofuran, by abstracting a hydrogen atom from compound (4-24-2) with a strong base such as for instance,  
 20 lithium diisopropyl amide, and then reacting with compound (4-24-5). Examples of compound (4-24-5) include, for instance, chloromethyl ethyl ether, chloromethyl

benzyl ether or the like. The strong base is used in the amounts of 1 equivalent to 2 equivalents based on compound (4-24-2). Compound (4-24-5) is used in the amounts of 1 equivalent to 2 equivalents based on compound (4-24-2). The reaction temperature is from -78°C to reflux temperature, and the reaction time is  
5 from 1 hour to 24 hours.

[Step 4-24-2]

The present step is a step wherein compound (4-24-2) and a formaldehyde equivalent are reacted to obtain compound (4-24-3). Compound (4-24-3) can be  
10 obtained in a solvent such as, for instance, tetrahydrofuran, by abstracting an hydrogen atom from compound (4-24-2) with a base such as, for instance, lithium diisopropyl amide, and then reacting with paraformaldehyde. The strong base is used in the amounts of 1 equivalent to 2 equivalents based on compound (4-24-2). The formaldehyde equivalent is used in the amounts of 1 equivalent to 2 equivalents  
15 based on compound (4-24-2). The reaction temperature is -78°C to reflux temperature, and the reaction time is from 1 hour to 24 hours.

[Step 4-24-3]

The present step is a step wherein a hydroxyl group of compound (4-24-3) is  
20 converted into a leaving group to obtain compound (4-24-4).

When L represents a methane sulfonyloxy group, a *p*-toluenesulfonyloxy group, or the like, compound (4-24-4) can be obtained in a solvent such as, for instance, dichloromethane, by reacting compound (4-24-3) and a sulfonyl halide such as methane sulfonyl chloride or *p*-toluenesulfonyl chloride, in the presence of an  
25 organic base such as, for instance, triethylamine. The organic base is used in the

amounts of 1 equivalent to 3 equivalents based on compound (4-24-3). The sulfonyl halide is used in the amounts of 1 equivalent to 2 equivalents based on compound (4-24-3). The reaction temperature is from 0°C to reflux temperature, and the reaction time is from 10 minutes to 24 hours.

- 5           When L represents a bromine atom or an iodine atom, compound (4-24-4) can be obtained in a solvent such as, for instance, dichloromethane, by action of a halogenation agent such as, for instance, carbon tetrabromide, N-bromosuccinimide or N-iodosuccinimide on compound (4-24-3), in the presence of triphenylphosphine. Triphenylphosphine is used in the amounts of 1 equivalent to 2 equivalents with  
10 based on compound (4-24-3). The halogenation agent is used in the amounts of 1 equivalent to 2 equivalents with respect to compound (4-24-3). The reaction temperature is from 0°C to room temperature, and the reaction time is from 10 minutes to 24 hours.

15   [Step 4-24-4]

- The present step is a process wherein compound (4-24-4) and compound (4-24-6) are reacted to obtain compound (4-24-1). Compound (4-24-1) can be obtained in a solvent such as for instance, N,N-dimethylformamide, abstracting a hydrogen atom from compound (4-24-4) using a base such as, for instance, sodium  
20 hydride, and reacting with compound (4-24-6). Compound (4-24-6) is used in the amounts of 1 equivalent to 10 equivalents based on compound (4-24-4). The base is used in the amounts of 1 equivalent to 10 equivalents based on compound (4-24-4). The reaction temperature is from 0°C to reflux temperature, and the reaction time is from 10 minutes to 24 hours.

25



[Examples]

The compounds according to the present invention can be prepared by a method described in, for instance, the following Preparation Examples and Examples. With the proviso that these are illustrative, and that the compounds according to the present invention is not to be limited in any way to the following specific examples.

Preparation Example A-1. 2-Amino-6-chloro-nicotinic acid

To liquid ammonia (approximately 20mL) was added 2,6-dichloro-nicotinic acid (0.38g, 2mmol) and copper(I) iodide (720mg, 3.8mmol) at -78°C in a sealed tube, and the solution was heated for 25 hours (the temperature of the oil bath was 115°C). The temperature of the oil bath was raised to 125°C, which was further heated for 14 hours 30 minutes. The reaction mixture was allowed to room temperature, and ammonia was evaporated. Methanol was added, insoluble matter was removed by filtration, and the filtrate was concentrated to obtain the title compound (0.25g, 1.45mmol, 72%) as a solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 6.63 (1H, d, J=8.0 Hz), 7.55 (2H, brs), 8.02 (1H, d, J=8.0 Hz).

Preparation Example A-2. 2-Amino-nicotinic acid methyl ester

2-Amino-nicotinic acid (10.0g, 72.4mmol) was dissolved in a mixed solution of methanol (200mL) and sulfuric acid (10mL), and the solution was stirred under reflux for 35 hours. A saturated aqueous solution of sodium bicarbonate was added to the reaction solution at 0°C, which was extracted with ethyl acetate, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by NH silica gel column

chromatography to obtain the title compound (5.26g, 34.6mmol, 48%) as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.89 (3H, s), 6.63 (1H, ddd, J=1.1, 4.8, 7.7Hz), 8.13 (1H, dd, J=1.6, 7.7Hz), 8.22 (1H, dd, J=1.8, 4.8Hz).

5

Preparation Example A-3. 2-Amino-5-nitro-nicotinic acid methyl ester

2-Amino-nicotinic acid methyl ester (1.00g, 6.57mmol) described in Preparation Example A-2 was dissolved at 0°C in a mixed solution of nitric acid (0.7mL) and sulfuric acid (2.6mL), which was stirred at 0°C for 40 minute and at room temperature for 19 hours, then, further stirred at 70°C for 4 hours. A saturated aqueous solution of sodium bicarbonate was added to the reaction solution at 0°C, which was extracted with ethyl acetate and tetrahydrofuran, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*. Methanol was added to the residue, the precipitated solid was filtered to obtain the title compound (459mg, 2.33mmol, 35%) as a white solid.

10

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.86 (3H, s), 8.14 (1H, brs), 8.62 (1H, brs), 8.68 (1H, d, J=2.7Hz), 9.04 (1H, d, J=2.9Hz).

15

Preparation Example A-4. 2-Amino-6-chloronicotinic acid

Tris(2-(2-methoxyethoxy)ethyl)amine (3.0mL, 9.4mmol) was added to a mixture of 2,6-dichloronicotinic acid (40g (90%purity), 0.19 mol), acetamide (80g, 1.4mol), potassium carbonate (78g, 0.56mol), copper(I) chloride (0.93g, 9.4mmol) and xylene (80mL), which was stirred overnight at 145°C. After cooling, copper(I) chloride (0.46g, 4.6mmol) was added to the reaction solution, which was stirred overnight at 145°C. After cooling the reaction solution to 105°C, water (100mL) was

20

25

added, the solution was stirred for 1 hour at the same temperature, and cooled down to room temperature. 5N hydrochloric acid (150mL) was added, the solution was neutralized with a citric acid aqueous solution, then, ethyl acetate was added, and the solution was filtered through Celite pad. The organic layer was washed with brine, then, the solvent was evaporated *in vacuo*. The residue was purified by silica gel column chromatography (ethyl acetate), recrystallization by the ethyl acetate-hexane was carried out to obtain the title compound (1.4g, 8.3mmol, 4.5%) as white crystal. <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 6.61 (1H, d, J=8.1Hz), 7.53 (2H, brs), 8.01 (1H, d, J=8.1Hz).

#### Preparation Example A-5. 2-Amino-6-(2-hydroxy-ethoxy)-nicotinic acid

To ethyleneglycol (0.50mL) was added sodium hydride (70mg, 1.7mmol, 60% in oil), catalytic amount of copper(I) iodide and 2-amino-6-chloronicotinic acid (30mg, 0.17mmol), which was stirred for 3 hours at 110°C, then, further stirred overnight at 80°C. After cooling, water, diethyl ether and aqueous ammonia was added to the reaction solution, which was then partitioned, the aqueous layer was neutralized with citric acid, then, extracted with dichloromethane. The organic layer was washed with brine, then dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo* to obtain the title compound (14mg).

#### Preparation Example A-6. 2-Amino-6-ethoxy-nicotinic acid

The title compound (35mg) was obtained from ethanol (0.50mL) and 2-amino-6-chloro-nicotinic acid (30mg, 0.17mmol) according to an analogous method to Preparation Example A-5.

Preparation Example A-7. 2-Amino-6-isopropoxy-nicotinic acid

The title compound (60mg) was obtained from isopropanol (0.50mL) and 2-amino-6-chloro-nicotinic acid (30mg, 0.17mmol) according to an analogous method to Preparation Example A-5.

5

Preparation Example A-8. 2-Amino-6-chloro-nicotinic acid methyl ester

To methanol (50mL) were added concentrated sulfuric acid (25mL) and 2-amino-6-chloro-nicotinic acid (4.3g, 25mmol) described in Preparation Example A-1 (or A-4) on an ice bath, which was stirred at 70°C for 5 hours. After cooling, an aqueous solution of sodium bicarbonate (90g) was added to neutralize. The resulting solid was filtered to obtain the title compound (3.2g, 17mmol, 68%) as a light brown solid.

10

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.88(3H, s), 6.62(1H, d, J=8.2Hz), 8.05(1H, d, J=8.1Hz).

15

Preparation Example A-9. Tributyl-methoxymethyl-stannane

To a mixture of diisopropylamine (9.4mL, 67mmol) and tetrahydrofuran (150mL) was added n-butyl lithium (2.4M n-hexane solution, 25mL, 61mmol) dropwise at -78°C, which was stirred for 30 minutes at the same temperature. Tributyltin hydride (16mL, 61mmol) was added dropwise at the same temperature, the solution was then stirred for 30 minutes on ice. The reaction solution was brought to -78°C, chloromethyl methyl ether (4.6mL, 61mmol) was added dropwise thereto, the solution was then gradually warmed to room temperature. Water (100mL) was added to the reaction solution, which was then extracted with diethyl ether (300mL). The organic layer was washed with brine, then, evaporated *in vacuo*.

20

25

The residue was purified by neutral silica gel column chromatography (heptane / ethyl acetate = 30 / 1), and the title compound (18g, 0.52mmol, 86%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.88-0.93(15H, m), 1.26-1.35(6H, m),  
5 1.47-1.55(6H, m) , 3.30(3H, s), 3.71(2H, t, J=6.8Hz).

Preparation Example A-10. 2-Amino-6-methoxymethyl-nicotinic acid methyl ester

A mixture of 2-amino-6-chloro-nicotinic acid methyl ester described in Preparation Example A-8 (1.4g, 7.6mmol), tributyl-methoxymethyl-stannane (3.1g,  
10 9.1mmol) described in Preparation Example A-9, tetrakis(triphenylphosphine)palladium (440mg, 0.38mmol) and N-methylpyrrolidinone (20mL) was stirred at 130°C for 3.5 hours. The reaction solution was allowed to cool, an aqueous solution of potassium fluoride and ethyl acetate were added thereto on an ice bath, followed by filtering through Celite pad. The organic layer was  
15 washed with brine, then, evaporated *in vacuo*. The residue was purified by silica gel column chromatography (heptane / ethyl acetate=2/1), and the title compound (0.93g, 4.7mmol, 63%) was obtained as a light brown oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.47(3H, s) , 3.88(3H, s) , 4.41(2H, s) , 6.74(1H, d, J=7.9Hz) , 8.14(1H, d, J=7.9Hz).

20

Preparation Example A-11. 2-Amino-6-methoxymethyl-nicotinic acid

Lithium hydroxide monohydrate (1.2g, 29mmol) was added to a mixture of 2-amino-6-methoxymethyl-nicotinic acid methyl ester described in Preparation Example A-10 (2.9g, 15mmol), tetrahydrofuran (30mL), methanol (7.5mL), and water  
25 (7.5mL), which was then stirred overnight at room temperature. Acetic acid (1.7mL,

29mmol) was added to the reaction solution and the solvent was evaporated *in vacuo*. After filtration using silica gel (methanol/ ethyl acetate=1/3) and the solvent was evaporated *in vacuo*, the residue was washed with water, and the title compound (2.1g,12mmol,80%) was obtained as a pale yellow solid.

- 5 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.32(3H, s) , 4.29(2H, s) , 6.61(1H, d, J=7.9Hz) , 7.16(2H, br s) , 8.02(1H, d, J=7.9Hz).

Preparation Example A-12. 2-(2-Cyanoethyl)-3,3-diaminopropenoic acid ethyl ester

- (1-Ethoxyformimidoyl) 1-acetic acid ethyl ester hydrochloride (50g, 0.26mol)  
10 was suspended in an ammonia-ethanol solution (300mL; prepared by saturating ethanol with ammonia gas at room temperature), which was then stirred at room temperature for 4 hours. After the reaction was completed, the precipitated salt was removed by filtration, and the filtrate was concentrated *in vacuo* at room temperature to reach 1/3 of the amount. Hydrochloric acid-methanol (130mL; hydrochloric acid  
15 content:7.5%) was added to this filtrate, the solution was then concentrated under a reduced pressure to obtain 3,3-diamino-acrylic acid ethyl ester hydrochloride (40g, 0.24mol, 92%) as a solid.

- The resulting 3,3-diamino-acrylic acid ethyl ester hydrochloride (2.2g, 13.2mmol) was suspended in tetrahydrofuran (40mL), triethylamine (2mL, 14.3mmol)  
20 and acrylonitrile (1.2mL, 19.3mmol) were added thereto, and the solution was refluxed for 6 hours. After the reaction was completed, the resulting triethylamine hydrochloride was filtered, and the filtrate was concentrated to obtain the title compound (0.6g, 3.3mmol, 25%).

- <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) :1.26 (3H, t, J=7.2Hz), 2.42-2.49 (2H, m),  
25 2.50-2.57 (2H, m), 4.12 (2H, q, J=7.2Hz), 4.22 (2H, brs), 6.45 (2H, brs).

Preparation Example A-13. 2,6-Diamino-4,5-dihydronicotinic acid ethyl ester

A solution of 2-(2-cyanoethyl) 3,3-diaminopropenoic acid ethyl ester described in Preparation Example A-12 (0.55g, 3mmol) in tetrahydrofuran (7mL) was added dropwise to a suspension of sodium hydride (208mg, 5.2mmol, 60% in oil) in tetrahydrofuran (7mL), and the solution was stirred for 19 hours 20 minutes under reflux. After the reaction was completed, the reaction solution was poured into an ice water, which was then extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, then, concentrated to obtain the title compound as a crude product (0.188g, 1mmol, 34%).

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) :1.27 (3H, t, J=7.2Hz), 2.28-2.34 (2H, m), 2.46-2.52 (2H, m), 4.14 (2H, q, J=7.2Hz).

Preparation Example A-14. 2,6-Diamino-nicotinic acid ethyl ester

To a solution of 2,6-diamino-4,5-dihydronicotinic acid ethyl ester described in Preparation Example A-13 (4.5g, 24.6mmol) in tetrahydrofuran (300mL) was added 2,3-Dichloro-5,6-dicyano-1,4-benzoquinone (5.6g, 24.7mmol), which was then stirred for 40 minutes at room temperature. The residue obtained by concentrating the reaction solution was then purified by NH silica gel column chromatography (ethyl acetate) to obtain a solid of the target compound. This solid was washed with hexane and dried to obtain the title compound (3.1g, 17.1mmol, 69.5%).

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) :1.35 (3H, t, J=7.2Hz), 4.27 (2H, q, J=7.2Hz), 4.60 (2H, brs), 5.82 (1H, d, J=8.4Hz), 7.90 (1H, d, J=8.4Hz).

Preparation Example A-15. 2,6-Diamino-nicotinic acid

2,6-Diamino-nicotinic acid ethyl ester described in Preparation Example A-14 (2g, 11mmol) was dissolved in ethanol (15mL), an aqueous solution of 1N sodium hydroxide (15mL) was added thereto, followed by stirring for 2 hours under reflux. The reaction solution was allowed to room temperature, ethanol was then removed  
5 by evaporation, the residue was cooled on an ice bath, then neutralized with 1N hydrochloric acid. The precipitated solid was collected by filtration, washed with water, then, dried to obtain the title compound (1.72g, 11mmol, quantitatively).

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 5.70 (1H, d, J=8.4Hz), 6.31 (2H, brs), 6.58-7.12 (1H, brs), 7.62 (1H, d, J=8.4Hz).

10

Preparation Example A-16. 2-Amino-6-vinyl-nicotinic acid methyl ester

2-Amino-6-chloro-nicotinic acid methyl ester (2.95g, 15.8mmol), vinyl tri-*n*-butyltin (5.01g, 15.8mmol) and tetrakis(triphenylphosphine)palladium(0) (1.83g, 1.58mmol) were suspended in xylene (15mL), and heated for 1 hour at 130°C. The  
15 reaction mixture was allowed to room temperature, water and ethyl acetate were added thereto, this mixture was filtered through Celite pad, then, the filtrate thereof was partitioned. The organic layer was separated, the solvent was evaporated *in vacuo*, and the residue was purified by silica gel column chromatography to obtain the title compound (1.87g, 10.5mmol, 66%) as a white solid.

20 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.81 (3H, s), 5.54 (1H, dd, J=1.6, 10.4 Hz), 6.24 (1H, dd, J=1.6, 17.2 Hz), 6.65 (1H, dd, J=10.4, 17.2 Hz), 6.76 (1H, d, J=8.0 Hz), 7.16 (1H, brs), 8.04 (1H, d, J=8.0 Hz).

Preparation Example A-17. 2-Amino-6-(2-cyanoethyl)-nicotinic acid methyl ester

25 To a solution of 2-amino-6-vinyl-nicotinic acid methyl ester (760mg,



4.26mmol) in tetrahydrofuran (76mL) was added a solution of diethyl aluminum cyanide in toluene (12.8mL, 12.8mmol) dropwise under sodium chloride – an ice bath cooling, at an internal temperature of -5°C or below, then, the solution was gradually allowed to room temperature and stirred overnight. The reaction solution was partitioned into an aqueous solution of saturated ammonium chloride and ethyl acetate. This organic layer was separated, washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography, and the title compound (180mg, 0.878mmol, 21%) was obtained as a yellow oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 2.80 (2H, t, J=7.2 Hz), 2.97 (2H, t, J=7.2 Hz), 3.88 (3H, s), 6.53 (1H, d, J=8.0 Hz) 8.07 (1H, d, J=8.0 Hz).

Preparation Example A-18. 2-Amino-6-(2-cyanoethyl)-nicotinic acid

To a solvent mixture of 2N-sodium hydroxide aqueous solution (5mL) and methanol (5mL) was added 2-amino-6-(2-cyanoethyl)-nicotinic acid methyl ester (90mg, 0.439mmol), and the solution was stirred for 18 hours at room temperature. This mixed solution was neutralized with 5N-hydrochloric acid, then, extracted with ethyl acetate. This organic layer was dried over anhydrous magnesium sulfate, then, the solvent was evaporated *in vacuo*, and the title compound (68mg, 0.355mmol, 81%) was obtained as a yellowish brown solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.86 (4H, bs), 6.54 (1H, d, J=8.0Hz), 7.18 (2H, bs), 7.98 (1H, d, J=8.0Hz).

Preparation Example A-19. 2-Amino-6-(2-ethoxy-vinyl)-nicotinic acid methyl ester

To a solution of ethyl ethynyl ether (3.6g, 25.7mmol) in tetrahydrofuran

(10mL) was added catechol borane (3.08g, 25.7mmol) on an ice bath. Immediately, the cold bath was removed, the reaction mixture was allowed to room temperature, and then, the solution was stirred for 1.5 hours under reflux. This reaction mixture was allowed to room temperature. To this mixture solution were added  
5 2-amino-6-chloro-nicotinic acid methyl ester (1.6g, 8.57mmol), sodium hydroxide powder (1.13g, 28.3mmol), tetrakis(triphenylphosphine)palladium(0) (0.99g, 0.857mmol) and dioxane (20mL), which was then stirred for 2 hours under reflux. The reaction mixture was allowed to room temperature, and partitioned into water and ethyl acetate. The organic layer was separated, the solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography, and the title  
10 compound (1.30g, 5.85mmol, 68%) was obtained as a brown solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 1.27 (3H, t, J=7.2 Hz), 3.77 (3H, s), 3.95 (2H, q, J=7.2 Hz), 5.75 (1H, d, J=12.8 Hz), 6.49 (1H, d, J=8.4 Hz), 7.02 (2H, brs), 7.63 (1H, d, J=12.8Hz), 7.89 (1H, d, J=8.4Hz).

15

Preparation Example A-20. 2-Amino-6-(2-hydroxy-ethyl)-nicotinic acid methyl ester

2-Amino-6-(2-ethoxy-vinyl)-nicotinic acid methyl ester (1.07g, 4.81mmol) was dissolved in 5N-hydrochloric acid (25mL), ethanol (20mL) and tetrahydrofuran (5mL), and the solution was stirred for 3 hours under reflux. This reaction mixture was  
20 allowed to room temperature, and neutralized with a aqueous solution of saturated sodium bicarbonate, then, sodium borohydride (1g, 26.5mmol) was added to this reaction mixture, which was then stirred for 20 minutes at room temperature. Ethyl acetate was added thereto, followed by filtrating through Celite pad. This filtrate was partitioned. This organic layer was separated, the solvent was evaporated *in vacuo*,  
25 the residue was purified by silica gel column chromatography, and the title compound

(350mg, 1.92mmol, 40%) was obtained as a brown oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.72 (2H, t, J=6.8 Hz), 3.71 (2H, q, J=6.8Hz), 3.79 (3H, s), 4.66 (1H, t, J=6.8Hz), 6.53 (1H, d, J=8.0 Hz), 7.12 (2H, brs), 7.95 (1H, d, J=8.0Hz).

5

Preparation Example A-21. 2-Amino-6-(2-fluoro-ethyl)-nicotinic acid methyl ester

A solution of (bis(2-methoxyethyl)amino)sulfur trifluoride (2.39g, 10.8mmol) in dichloromethane (30mL) was cooled with a dry ice-methanol bath, and 2-amino-6-(2-hydroxyethyl)-nicotinic acid methyl ester (50mg, 0.255mmol) was added thereto dropwise. After dropwise addition the cold bath was immediately removed, and the solution was gradually allowed to room temperature. Water and ethyl acetate were added to the reaction solution, which was then partitioned. The organic layer was separated, the solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography, and the title compound (4mg, 0.020mmol, 7.9%) was obtained as a white solid.

15

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.97 (2H, td, J=6.0, 26.0 Hz), 3.80 (3H, s), 4.77 (2H, td, J=6.0, 47.2 Hz), 6.58 (1H, d, J=8.0 Hz), 7.16 (2H, bs), 8.00 (1H, d, J=8.0 Hz).

20 Preparation Example A-22. 2-Amino-6-(2-fluoro-ethyl)-nicotinic acid

2-Amino-6-(2-fluoro-ethyl)-nicotinic acid methyl ester (77mg, 0.387mmol) was dissolved in an aqueous solution of 2N sodium hydroxide (5mL) and methanol (5mL), and the solution was stirred for 20 minutes at room temperature. This mixture solution was neutralized with 5N hydrochloric acid, then, extracted with ethyl acetate.

25 The organic layer was separated, the solvent was evaporated *in vacuo*, and the title

compound (64mg, 0.348mmol, 90%/65% purity) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.97 (2H, td, J=6.0, 26.0 Hz), 4.77 (2H, td, J=6.0, 47.2 Hz), 6.57 (1H, d, J=8.0 Hz), 7.98 (1H, d, J=8.0 Hz).

5 Preparation Example A-23. Tributyl-ethoxymethyl-stannane

The title compound (2.8g, 8.0mmol, 67%) was obtained as a colorless oil from chloromethyl ethyl ether (1.1mL, 12mmol) according to an analogous method to Preparation Example A-9.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.87-0.92(15H, m), 1.16(3H, t, J=7.0Hz),  
10 1.26-1.35(6H, m), 1.43-1.55(6H, m), 3.36(2H, q, J=7.0Hz), 3.74(2H, t, J=6.5Hz).

Preparation Example A-24. 2-Amino-6-ethoxymethyl-nicotinic acid methyl ester

The title compound (0.35g, 1.7mmol, 39%) was obtained as a pale yellow solid from tributyl-ethoxymethyl-stannane (2.0g, 6.3mmol) described in Preparation  
15 Example A-23 and 2-amino-6-chloro-nicotinic acid methyl ester described in Preparation Example A-8 (0.80g, 4.3mmol) according to an analogous method to Preparation Example A-10.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 1.28(3H, t, J=7.0Hz), 3.61(2H, q, J=7.0Hz),  
3.88(3H, s), 4.45(2H, s), 6.41(2H, br s), 6.78(1H, d, J=7.9Hz), 8.13(1H, d, J=8.1Hz).

20

Preparation Example A-25. 2-Amino-6-ethoxymethyl-nicotinic acid

The title compound (180mg, 0.92mmol, 57%) was obtained as a pale yellow solid from 2-amino-6-ethoxymethyl-nicotinic acid methyl ester (330mg, 1.6mmol) described in Preparation Example A-24 according to an analogous method to  
25 Preparation Example A-11.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 1.15(3H, t, J=7.1Hz), 3.51(2H, q, J=7.0Hz), 4.33(2H, s), 6.64(1H, d, J=7.9Hz), 8.02(1H, d, J=7.9Hz).

Preparation Example A-26. Tributyl-isopropoxymethyl-stannane

5           To a mixture of isopropanol (2mL) and tetrahydrofuran (2mL) was added sodium hydride (66%, 58mg, 1.6mmol) on an ice bath, then, the solution was stirred for 20 minutes at room temperature. To the reaction solution was added a solution of tributyl-iodomethyl-stannane (230mg, 0.53mmol), synthesized according to the document (Synthetic Communications, Vol.24, No.8, pp. 1117-1120), in  
10       tetrahydrofuran (1mL) dropwise on an ice bath, then, N,N-dimethylformamide (0.5mL) was added thereto, followed by stirring overnight at room temperature. The reaction solution was partitioned into water (20mL) and diethyl ether (50mL). The organic layer was separated, washed with brine, and then, evaporated *in vacuo*. The residue was purified by neutral silica gel column chromatography (heptane : ethyl  
15       acetate = 30 : 1), and the title compound (63mg, 0.17mmol, 32%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.87-0.91(15H, m), 1.11(6H, d, J=6.0Hz), 1.26-1.35(6H, m), 1.47-1.53(6H, m), 3.28-3.31(1H, m), 3.69(2H, t, J=7.6Hz).

20       Preparation Example A-27. Butoxymethyl-tributyl-stannane

          The title compound (220mg, 0.58mmol, 99%) was obtained as a colorless oil from tributyl-iodomethyl-stannane (250mg, 0.58mmol) according to an analogous method to Preparation Example A-26.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.87-0.93(18H, m), 1.26-1.38(8H, m),  
25       1.47-1.55(8H, m), 3.30(2H, t, J=6.5Hz), 3.73 (2H, t, J=6.5Hz).

Preparation Example A-28. Tributyl-propoxymethyl-stannane

The title compound (230mg, 0.63mmol, 97%) was obtained as a colorless oil from tributyl-iodomethyl-stannane (280mg, 0.65mmol) according to an analogous  
5 method to Preparation Example A-26.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.87-0.91(18H, m), 1.26-1.35(6H, m), 1.47-1.58(8H, m), 3.27(2H, t, J=6.5Hz), 3.73 (2H, t, J=6.5Hz).

Preparation Example A+-1. Sodium 4-(((2-Aminopyridine-3-carbonyl)-amino)-methyl)-phenolate  
10

To a solution of 4-hydroxybenzaldehyde (10g, 81.9mmol) in methanol (45mL) was added Raney nickel (3g) and 7N aqueous ammonia solution (45mL), and the solution was stirred under hydrogen atmosphere (1 atm) at room temperature for 21 hours. The reaction solution was filtered through Celite pad to remove the catalyst,  
15 the filtrate was concentrated, and 4-aminomethyl-phenol (10g, quantitatively) was obtained as a pale green solid.

Then, a solution of 2-aminonicotinic acid (3.0g, 21.7mmol) in N,N-dimethylformamide (30mL) was cooled with an ice water, 1-hydroxybenzotriazole (3.51g, 26mmol), (3-dimethylaminopropyl)-ethyl-carbodiimide (4.04g, 26mmol) and a  
20 solution of the resulting 4-aminomethyl-phenol (3.0g, 21.7mmol) in N,N-dimethylformamide (20mL) were added, and the solution was stirred for 18 hours at this temperature. The reaction solution was partitioned into brine, the organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated. The residue was dissolved in ethyl acetate, filtration was carried out using NH silica  
25 gel, and the filtrate was concentrated. The residue was dissolved in methanol

(90mL), 1N sodium hydroxide (17.8mL, 17.8mmol) was added thereto, followed by stirring at room temperature for an hour and a half. The reaction solution was concentrated *in vacuo*, and the title compound (5.66g) was obtained as a pale yellow solid.

5

Preparation Example A+-2. (6-Amino-5-((5-(3-fluoro-phenoxy)-thiophene-2-yl)methyl)-carbamoyl)-pyridine-2-yl)-carbamic acid *tert*-butyl ester

To a solution of 6-amino-nicotinic acid (270mg, 2.0mmol) and C-(5-(3-fluorophenoxy)thiophen-2-yl)methylamine (400mg, 1.8mmol) in  
 10 N,N-dimethylformamide (10mL) were added benzotriazole-1-yl-tris(dimethylamino)phosphonium hexafluorophosphate (870mg, 2.0mmol) and triethylamine (0.50mL, 3.6mmol), and the solution was stirred for 30 minutes at 60°C. Water and ethyl acetate were added to the reaction solution, which was then partitioned, and the organic layer was washed twice with water.  
 15 Silica gel was added to the organic layer, the solvent was evaporated *in vacuo* for adsorption, purification was carried out with NH silica gel column chromatography (ethyl acetate, then ethyl acetate : methanol = 10 : 1), and 6-amino-N-(5-(3-fluoro-phenoxy)-thiophen-2-yl)-methylnicotinamide (270mg, 0.79mmol, 43.9%) was obtained.

20 To the resulting 6-amino-N-(5-(3-fluoro-phenoxy)-thiophen-2-yl)-methylnicotinamide (270mg, 0.79mmol) were added di-*tert*-butyldicarbonate (210mg, 0.94mmol) and *tert*-butyl alcohol (15mL), and the solution was stirred for 16.5 hours at room temperature. NH silica gel was added to the reaction solution, solvent was evaporated *in vacuo* for  
 25 adsorption, purification was carried out by NH silica gel column chromatography

(hexane : ethyl acetate = 4:1 then 1:1), and the title compound (250mg, 0.54mmol, 68.3%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 1.46 (9H, s), 4.52 (2H, d, J=5.6Hz), 6.55-6.59 (1H, m), 6.78-6.82 (1H, m), 6.88-7.00 (4H, m), 7.36-7.44 (1H, m), 7.85 (1H, d, J=8.8Hz), 8.15 (1H, d, J=8.8Hz), 8.70 (1H, s), 9.14 (1H, t, J=5.6Hz), 10.1 (1H, s).

Preparation Example A+-3. (6-Amino-5-((5-(3-fluoro-phenoxy)-thiophen-2-ylmethyl)-carbamoyl)-pyridin-2-yl)-methyl-carbamic acid *tert*-butyl ester

To a solution of (6-amino-5-((5-(3-fluoro-phenoxy)-thiophen-2-ylmethyl)-carbamoyl)-pyridine-2-yl)-carbamic acid *tert*-butyl ester described in Preparation Example A+-2 (125mg, 0.27mmol) and methyl iodide (43mg, 0.29mmol) in N,N-dimethylformamide (5mL) was added sodium hydride (12mg, 0.29mmol, 60% in oil) on an ice bath, and the solution was stirred for 1 hour at room temperature. NH silica gel was added to the reaction solution, the solvent was evaporated *in vacuo* for adsorption, purification was carried out by NH silica gel column chromatography (hexane : ethyl acetate = 4 : 1 then 2 : 1), and the title compound (87mg, 0.19mmol, 70.5%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 1.46 (9H, s), 3.31 (3H, s), 4.53 (2H, d, J=5.6Hz), 6.57 (1H, d, J=3.6Hz), 6.81 (1H, d, J=3.6Hz), 6.86-7.00 (3H, m), 7.36-7.44 (1H, m), 7.76 (1H, d, J=8.8Hz), 8.14 (1H, dd, J=2.0, 8.8Hz), 8.80 (1H, d, J=2.0Hz), 9.22 (1H, t, J=5.6Hz).

Preparation Example A+-4. (6-Amino-5-((5-benzyl-thiophen-2-ylmethyl)-carbamoyl)-pyridine-2-yl)-carbamoylmethyl-carbamic acid *tert*-butyl ester



To a solution of 6-aminonicotinic acid (340mg, 2.4mmol) and C-(5-benzyl-thiophen-2-yl)-methylamine described in Preparation Example 42 (450mg, 2.2mmol) in N,N-dimethylformamide (5mL) were added benzotriazol-1-yl-tris(dimethylamino)phosphonium hexafluorophosphate (1.1g, 2.4mmol) and triethylamine(0.62mL, 4.4mmol), and the solution was stirred at 60°C for 1 hour. Water and ethyl acetate were added to the reaction solution, which was then partitioned, and the organic layer was washed twice with water. Silica gel was added to the organic layer, the solvent was evaporated *in vacuo* adsorption, purification was carried out by NH silica gel column chromatography (hexane : ethyl acetate = 1:1, then ethyl acetate, then ethyl acetate : methanol = 10 : 1), and 6-amino-N-(5-benzyl-thiophen-2-ylmethyl)-nicotinamide (210mg, 0.65mmol, 29.5%) was obtained.

Then, according to an analogous method to Preparation Example A+-2, from the resulting 6-amino-N-(5-benzyl-thiophen-2-ylmethyl)-nicotinamide (210mg, 0.65mmol), (5-((5-benzyl-thiophen-2-ylmethyl)-carbamoyl)-pyridin-2-yl)-carbamic acid *tert*-butyl ester (120mg, 0.28mmol, 43.0%) was obtained as a colorless oil.

To a solution of this oil (60mg, 0.14mmol) in N,N-dimethylformamide (5mL) were added sodium hydride (6.8mg, 0.14mmol, 60% in oil) and bromoacetamide (23mg, 0.16mmol), and the solution was stirred for 25 hours room temperature. Water and ethyl acetate were added to the reaction solution, which was then partitioned, the organic layer was evaporated *in vacuo*, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 1:1, then ethyl acetate, then ethyl acetate : methanol = 20 : 1), and the title compound (23mg, 0.047mmol, 33.5%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 1.43 (9H, s), 4.04 (2H, s), 4.49-4.54 (4H, m), 6.69 (1H, d, J=3.6Hz), 6.79 (1H, d, J=3.6Hz), 6.94 (1H, s), 7.16-7.30 (5H, m), 7.38 (1H, s), 7.87 (1H, d, J=8.8Hz), 8.10-8.16 (1H, m), 8.68-8.73 (1H, m), 9.13 (1H, t, J=5.6Hz).

5

Preparation Example A+-5. 6-Chloro-N-(1-(3-fluoro-benzyl)-1H-pyrrol-3-ylmethyl)-nicotinamide

6-chloro-nicotinic acid (100mg, 0.58mmol), triethylamine (0.194mL, 1.39mmol) and benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (308mg, 0.696mmol) were dissolved in N,N-dimethylformamide (3mL), C-(1-(3-fluoro-benzyl)-1H-pyrrol-3-yl)-methylamine (142mg, 0.695mmol) described in Preparation Example 59 was added, and the solution was stirred for 15 hours 10 minutes at room temperature. After the reaction was completed, the reaction solution was poured into brine, and the solution was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, then the solution was concentrated, the resulting residue was purified by NH silica gel column chromatography (ethyl acetate : hexane = 1 : 1), and the title compound (0.14g, 0.39mmol, 67%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.42 (2H, d, J=4.8Hz), 5.02 (2H, s), 5.99-6.09 (1H, m), 6.16-6.18 (1H, m), 6.56 (1H, d, J=8.0 Hz), 6.57 (1H, brs), 6.64-6.68 (2H, m), 6.78-6.83 (1H, m), 6.91-6.94(1H, m), 6.96-7.02 (1H, m), 7.27-7.33 (1H, m), 7.49 (1H, d, J=8.0 Hz).

Preparation Example A+-6. 2-(Ethoxymethyl-amino)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

25

To a solution of 2-aminonicotinic acid (3245mg, 23.49mmol) in N,N-dimethylformamide (200mL) were added C-(5-phenoxy-thiophen-2-yl)methylamine (5305mg, 25.84mmol) described in Preparation Example 24, benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (12.49g, 28.19mmol) and triethylamine (7.86mL, 56.38mmol), and the solution was stirred for 2 days at room temperature. Water was added to the reaction mixture, which was then extracted with ethyl acetate, the organic layer was sequentially washed with an aqueous solution of saturated sodium bicarbonate, water, then brine, the solvent was evaporated *in vacuo*, and 2-amino-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide (4999mg, 15.36mmol, 65%) was obtained as crude product.

To a solution of the resulting 2-amino-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide (602mg, 1.85mmol) in ethanol (40mL) were added 5,5-dimethylhydantoin (260mg, 2.04mmol), 37% formic acid aqueous solution (3.00mL, 23.6mmol), and the solution was stirred under reflux for 1 hour. Water was added to the reaction mixture, which was then extracted with ethyl acetate, the organic layer was sequentially washed with water and brine, dried over anhydrous sodium sulfate, and then, evaporated *in vacuo*. The resulting residue was purified by silica gel chromatography (hexane-ethyl acetate), and the title compound (430mg, 1.12mmol, 61%) was obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) :1.03 (3H, t, J=6.8Hz), 3.41 (2H, q, J=6.8Hz), 4.48 (2H, d, J=5.6Hz), 4.90 (2H, d, J=7.2Hz), 6.49 (1H, d, J=3.6Hz), 6.68 (1H, dd, J=4.8, 7.6Hz), 6.77 (1H, d, J=3.6Hz), 7.06-7.14 (3H, m), 7.33-7.38(2H, m), 7.94 (1H, dd, J=1.6, 7.6Hz), 8.19 (1H, dd, J=1.6, 4.8Hz), 8.91 (1H, t, J=7.2Hz), 9.16-9.20 (1H, m).

Preparation Example A+-7. 2-Chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-  
nicotinamide

To a solution of 2-chloro nicotinic acid (1182mg, 7.50mmol) in  
 5 N,N-dimethylformamide (3mL) were added C-(5-phenoxy-thiophen-2-yl)methylamine  
 described in Preparation Example 24 (1693mg, 8.25mmol), benzotriazol-1-yloxy  
 tris(dimethylamino)phosphonium hexafluorophosphate (3987mg, 9.0mmol) and  
 triethylamine (2.5mL, 18.0mmol), and the solution was stirred at 60°C for 2 days.  
 Water was added to the reaction mixture, which was then extracted with ethyl acetate,  
 10 and then concentrated. The residue was purified by silica gel column  
 chromatography, and the title compound (1181mg, 3.43mmol, 46%) was obtained.  
<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.51(2H, d, J=5.2Hz), 6.52 (1H, d, J=3.6Hz),  
 6.80 (1H, d, J=3.6Hz), 7.07-7.16 (3H, m), 7.30-7.45 (2H, m), 7.47 (1H, dd, J=4.8,  
 7.6Hz), 7.87 (1H, dd, J=1.6, 7.6Hz), 8.46 (1H, dd, J=1.6, 4.8Hz), 9.20 (1H, t,  
 15 J=5.2Hz).

Preparation Example A+-8. N-(4-Benzyloxy-benzyl)-6-(ethoxymethyl-amino)-  
nicotinamide

To a solution of 6-aminonicotinic acid (130mg, 0.941mmol) and  
 20 4-benzyloxy-benzylamine (201mg, 0.941mmol) described in Preparation Example 1  
 in N,N-dimethylformamide (5mL), was added  
 benzotriazol-1-yl-tris(dimethylamino)phosphonium hexafluorophosphate (624mg,  
 1.41mmol) and triethylamine (394μl, 2.82mmol), and the solution was stirred for 40  
 minutes at 80°C. Water and ethyl acetate were added to the reaction solution,  
 25 which was then partitioned, and the organic layer was washed twice with water. The

solvent was evaporated *in vacuo*, ethyl acetate was added to the residue, a white insoluble matter was collected by filtration, and 6-amino-N-(4-benzyloxy-benzyl)-nicotinamide (202mg, 0.606mmol, 64%) was obtained.

5 To a solution of the resulting 6-amino-N-(4-benzyloxy-benzyl)-nicotinamide (200mg, 0.556mmol) in ethanol (10mL) were added 5,5-dimethylimidazophospho-2,4-dione (85mg, 0.66mmol) and 37% formaldehyde aqueous solution (1mL), and the solution was stirred for 1 hour under reflux. Water and ethyl acetate were added to the reaction solution, which was then partitioned,  
10 and the organic layer was washed twice. NH silica gel was added to the organic layer, the solvent was evaporated *in vacuo* for adsorption, purification was carried out by NH silica gel column chromatography (ethyl acetate), and the title compound (95mg, 0.243mmol, 40%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 1.20 (3H, t, J=7.0Hz), 3.55 (2H, q, J=7.0Hz), 4.56  
15 (2H, d, J=5.5Hz), 4.84 (2H, s), 5.07 (2H, s), 5.68 (1H, brs), 6.14 (1H, brs), 6.62 (1H, d, J=8.8Hz), 6.96 (2H, d, J=8.8Hz), 7.22-7.35 (4H, m), 7.27-7.44 (3H, m), 7.93 (1H, dd, J=2.4, 8.8Hz), 8.54 (1H, d, J=2.4Hz).

Preparation Example A+-9. 2-Amino-5-nitro-N-(5-phenoxy-thiophen-2-ylmethyl)-  
20 nicotinamide

2-Amino-5-nitro-nicotinic acid methyl ester (48.4mg, 0.245mmol) described in Preparation Example A-3 and lithium hydroxide monohydrate (10.3mg, 0.245mmol) were dissolved in a solvent mixture of tetrahydrofuran (1mL), methanol (0.1mL) and water (0.1mL), and the solution was stirred for 17 hours at room temperature. The  
25 solvent was evaporated *in vacuo*, and 2-amino-5-nitro-nicotinic acid was obtained as

a lithium salt.

Then, the resulting lithium salt of 2-amino-5-nitro-nicotinic acid, C-(5-phenoxy-thiophen-2-yl)-methylamine (60mg, 0.29mmol), benzotriazol-1-yl-tris(dimethylamino)phosphonium hexafluorophosphate (162mg, 0.367mmol) and triethylamine (103μl, 0.735mmol) were dissolved in N,N-dimethylformamide (2.0mL), and the solution was stirred for 6 hours at room temperature. Water was added to the reaction solution, which was then extracted with ethyl acetate, and the organic layer was washed with water and brine. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 1 : 1), and the title compound (87mg, 0.24mmol, 96%) was obtained as a pale yellow solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.49 (2H, d, J=5.5Hz), 6.50 (1H, d, J=3.7Hz), 6.80 (1H, d, J=3.1Hz), 7.08 (2H, d, J=7.7Hz), 7.13 (1H, t, J=7.5Hz), 7.37 (2H, t, J=7.5Hz), 8.76 (1H, d, J=2.2Hz), 8.96 (1H, d, J=1.7Hz), 9.51 (1H, t, J=5.5Hz).

15

Preparation Example A+-10. 2,5-Diamino-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

2-Amino-5-nitro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide (74mg, 0.20mmol) described in Preparation Example A+-9, an iron powder (56mg, 1.0mmol) and ammonium chloride (21mg, 0.40mmol) were suspended in a solvent mixture of ethanol (2mL) and water (0.5mL), the solution was stirred at 60°C for 3 hours, then, it was stirred at 90°C for 6 hours. The reaction solution was cooled to room temperature, then, it was filtered through Celite pad. The filtrate was evaporated *in vacuo*, the residue was purified by reverse phase high performance liquid chromatography (acetonitrile-water: mobile phase (containing 0.1% trifluoroacetic

25

acid) was used), and the title compound (54.4mg) was obtained as a trifluoroacetic acid salt.

MS m/e (ESI) 341.26(MH<sup>+</sup>)

5 Preparation Example A+-11. 2-Amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

2-Amino-6-chloro-nicotinic acid (400mg, 2.31mmol) described in Preparation Example A-1 was dissolved in N,N-dimethylformamide (10mL), triethylamine (0.78mL, 5.6mmol), benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate  
10 (1.23g, 2.8mmol) and C-(5-phenoxy-thiophen-2-yl)-methylamine (572mg, 2.8mmol) described in Preparation Example 35 were added thereto, and the solution was stirred for 13 hours 30 minutes at room temperature. After the reaction was completed, the reaction solution was poured into brine, and the solution was extracted with ethyl acetate. The organic layer was dried over  
15 anhydrous magnesium sulfate, then concentrated, the resulting residue was purified by silica gel column chromatography (ethyl acetate : hexane = 1 : 1), and the title compound (380mg, 1.05mmol, 46%) was obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.47 (2H, d, J=6.0Hz), 6.50 (1H, d, J=4.0Hz), 6.64 (1H, d, J=8.0Hz), 6.78 (1H, d, J=4.0Hz), 7.07-7.17 (3H, m), 7.36-7.41 (2H, m),  
20 7.53 (2H, brs), 7.93 (1H, d, J=8.0Hz), 9.11 (1H, t, J=6.0Hz).

Preparation Example A+-12. 2-Amino-6-(2-amino-ethylamino)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

2-Amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide (150mg,  
25 0.417mmol) described in Preparation Example A+-11, ethane-1,2-diamine (418μl,

6.25mmol) were dissolved in a mixture solution of dimethylsulfoxide (2mL) and N,N-diisopropylethylamine (1mL), and the solution was stirred at 120°C for 15 hours. The reaction solution was cooled to room temperature, water was added, then extracted with ethyl acetate and tetrahydrofuran, the organic layer was washed with  
5 brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, tetrahydrofuran and NH silica gel were added to the residue, the solvent was evaporated *in vacuo* adsorption, purification was carried out by NH silica gel column chromatography (ethyl acetate : methanol = 10 : 1), and the title compound (95mg,0.25mmol,59%) was obtained as a white solid.

10 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.64 (2H, t, J=6.4Hz), 3.16-3.22 (2H, m), 4.41 (2H, d, J=5.7Hz), 5.70 (1H, d, J=8.6Hz), 6.48 (1H, d, J=3.8Hz), 6.67 (1H, brs), 6.72 (1H, d, J=3.8Hz), 7.02 (2H, s), 7.08 (2H, d, J=8.6Hz), 7.14 (1H, t, J=7.3Hz), 7.38 (2H, dd, J=7.3, 8.6Hz), 7.59 (1H, d, J=8.6Hz), 8.38 (1H, t, J=5.7Hz).

15 Preparation Example A+-13. 2-Amino-6-(2-(4-nitro-phenylamino)-ethylamino)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

2-Amino-6-(2-amino-ethylamino)-N-(5-phenoxy-thiophen-2-ylmethyl)-  
nicotinamide (25mg, 65μmol) described in Preparation Example A+-12 ,  
4-fluoronitrobenzene (7.6μl, 71μmol) and N,N-diisopropylethylamine (22.7μl,  
20 130μmol) were dissolved in dimethylsulfoxide (0.5mL), the solution was stirred for 3.  
5 hours at room temperature, and then, the solution was stirred for 15.5 hours at  
70°C. The reaction solution was cooled to room temperature, water was added, the  
solution was extracted with ethyl acetate, the organic layer was washed with brine  
and dried over anhydrous magnesium sulfate. The solvent was evaporated *in*  
25 *vacuo*, residue was purified by reverse phase high performance liquid



chromatography (mobile phase:acetonitrile-water (containing 0.1% trifluoroacetic acid) was used), and the title compound (23mg) was obtained as a trifluoroacetic acid salt.

MS m/e (ESI) 505.37(MH<sup>+</sup>)

5

Preparation Example A+-14. 2-Amino-6-(1-ethoxyvinyl)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

1-Ethoxyvinyl(tri-*n*-butyl)tin (0.47mL, 1.4mmol) was added to a mixture of 2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide (170mg, 0.46mmol) described in Preparation Example A+-11, tetrakis(triphenylphosphine)palladium(0) (54mg, 0.046mmol) and xylene (5mL), and the solution was stirred for 2.5 hours at 130°C. After cooling, water and ethyl acetate were added to the reaction solution for extraction, and the solution was washed with brine. The solvent was evaporated *in vacuo*, then, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 2 : 1), and the title compound (150mg, 0.38mmol, 82%) was obtained as a pale yellow solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) :1.42 (3H, t, J=7.0Hz), 3.93 (2H, q, J=7.0Hz), 4.35 (1H, d, J=1.8Hz), 4.65 (2H, d, J=5.3Hz), 5.37 (1H, d, J=1.8Hz), 6.30-6.32 (3H, m), 6.39 (1H, d, J=3.7Hz), 6.74 (1H, d, J=3.8Hz), 7.00 (1H, d, J=8.1Hz), 7.08-7.13 (3H, m), 7.30-7.35 (2H, m), 7.59 (1H, d, J=8.1Hz).

20

Preparation Example A+-15. 2-Amino-N-(5-phenoxy-thiophen-2-ylmethyl)-6-tributylstanyl-nicotinamide

To a mixture of 2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide (1.1g, 3.0mmol) described in Preparation Example A+-11,

25

tetrakis(triphenylphosphine)palladium(0) (170mg, 0.15mmol) and xylene (5mL) was added bis(tri *n*-butyl tin) (9.1mL, 18mmol), and the solution was stirred at 135°C for 2 hours. After cooling, the reaction solution was directly purified by neutral silica gel column chromatography (hexane : ethyl acetate = 3 : 1), then, the resulting crude product was washed with hexane cooled to 0°C to obtain the title compound (600mg, 0.98mmol, 33%) as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.86-0.90 (9H, m), 1.05-1.09 (6H, m), 1.27-1.36 (6H, m), 1.50-1.58 (6H, m), 4.64 (1H, d, J=5.5Hz) , 6.26-6.30 (4H, m), 6.38 (1H, d, J=3.8Hz) , 6.73-6.74 (2H, m) , 7.08-7.12 (3H, m), 7.31-7.36 (3H, m).

#### Preparation Example A+-16. 2-Amino-5-iodo-N-(5-phenoxy-thiophene-2-ylmethyl)-nicotinamide

2-Amino-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide (250mg, 0.768mmol) obtained from 2-aminonicotinic acid and C-(5-phenoxy-thiophen-2-yl)methylamine described in Preparation Example 24 according to an analogous method to Preparation Example A+-5, and N-iodosuccinimide (190mg, 0.845mmol) were dissolved in tetrahydrofuran (5mL), and the solution was stirred for 16 hours at room temperature. An aqueous solution of saturated sodium bicarbonate was added to the reaction solution at 0°C, which was then extracted with ethyl acetate, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 1 : 1), and title compound (45mg, 0.10mmol, 13%) was obtained as a pale yellow solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.45 (2H, d, J=5.7Hz), 6.49 (1H, d, J=3.7Hz), 6.77 (1H, d, J=3.8Hz), 7.08 (2H, d, J=7.7Hz), 7.13 (1H, t, J=7.3Hz), 7.23 (2H, s), 7.37

(2H, t, J=7.3Hz), 8.15 (1H, d, J=2.2Hz), 8.21 (1H, d, J=1.8Hz), 9.13 (1H, d, J=5.7Hz).

Preparation Example A+-17. 2-Amino-N-(3-hydroxybenzyl)-nicotinamide

The title compound (0.63g, 2.6mmol, 53%) was obtained as a white solid  
5 from 3-aminomethylphenol (0.60g, 4.9mmol) described in Preparation Example 130  
and 2-aminonicotinic acid (0.67g, 4.9mmol) according to an analogous method to  
Preparation Example Q+-1.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.34 (2H, d, J=5.9Hz), 6.56-6.61 (2H, m),  
6.69-6.71 (2H, m), 7.06-7.10 (3H, m), 7.93 (1H, dd, J=1.8, 7.8Hz), 8.06 (1H, dd,  
10 J=1.8, 4.8Hz), 8.91 (1H, t, J=6.0Hz), 9.30 (1H, s).

Preparation Example A+-18. 2-Amino-N-(4-benzyloxy-benzyl)-6-chloro-nicotinamide

The title compound (0.43g, 1.2mmol, 28%) was obtained as a white solid  
from 14-benzyloxy-benzylamine (0.90g, 4.2mmol) described in Preparation Example  
15 and 2-amino-6-chloro-nicotinic acid (1.5g, 8.4mmol) described in Preparation  
Example A-1 (or A-4) according to an analogous method to Preparation Example  
Q+-1.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.33 (2H, d, J=5.7Hz), 5.06 (2H, s), 6.61 (1H,  
d, J=8.1Hz), 6.94 (2H, d, J=8.6Hz), 7.20 (2H, d, J=8.4Hz), 7.28-7.31 (1H, m),  
20 7.34-7.38 (2H, m) , 7.41 (2H, d, J=7.5Hz), 7.49 (2H, brs), 7.95 (1H, d, J=8.1Hz),  
8.92-8.95 (1H, m).

Preparation Example A+-19. 5-Bromo-thiophene-2-carbaldehyde oxime

To a mixture of 2-bromo-5-formylthiophene (2.5mL, 21mmol) and pyridine  
25 (25mL) was added hydroxylamine hydrochloride (2.2g, 32mmol) on an ice bath, then,

the solution was stirred overnight at room temperature. This reaction mixture was concentrated *in vacuo*, then, partitioned with water (50mL), ethyl acetate (50mL) then 1N hydrochloric acid solution (50mL). The organic layer was separated, sequentially washed with an aqueous solution of saturated sodium bicarbonate and brine, and then, dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was washed with heptane-ethyl acetate (30:1), and the title compound (4.3g, 21mmol, 98%) was obtained as a light brown solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 7.07(1H, d, J=4.0Hz), 7.11(1H, d, J=4.0Hz), 7.62(1H, s), 8.35(1H, br s).

#### Preparation Example A+-20, 5-Bromo-thiophene-2-carbonitrile

To a mixture of 5-bromo-thiophene-2-carbaldehyde oxime described in Preparation Example A+-19 (1.3g, 6.2mmol) and tetrahydrofuran (15mL) were acetic acid (1.4mL, 25mmol) and acetic anhydride (1.5mL, 15mmol) at room temperature, then, stirred at 50°C for 2 hours, and further stirred at 70°C for 8 hours. After cooling this reaction mixture, the solution was concentrated *in vacuo*. The residue was purified by NH-silica gel column chromatography (heptane : ethyl acetate = 8 : 1), and the title compound (1.0g, 5.4mmol, 88%) was obtained as a colorless solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 7.11(1H, d, J=4.0Hz), 7.40(1H, d, J=4.0Hz).

#### Preparation Example A+-21. C-(5-Bromo-thiophen-2-yl)-methylamine

To a mixture of lithium aluminum hydride (1.6g, 41mmol) and tetrahydrofuran (45mL) was aluminum chloride (6.1g, 46mmol) on an ice bath, then, the solution was stirred for 1 hour at room temperature. The reaction solution was cooled to -20°C, a solution of 5-bromo-thiophene-2-carbonitrile described in Preparation Example A+-20

(4.3g, 25mmol) in tetrahydrofuran (5mL) was added dropwise at the same temperature. After stirring at 2°C for 20 minutes, the reaction solution was cooled to -10°C, and while maintaining the internal temperature at 0°C or lower, tetrahydrofuran (300mL) and 28% aqueous ammonia solution (5mL) were added thereto.

5 Anhydrous magnesium sulfate was added to the reaction solution, which was then filtered using a filter paper, and then, concentrated *in vacuo*. The residue was purified by NH-silica gel column chromatography (ethyl acetate), and the title compound (840mg, 4.4mmol, 85%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.99(2H, s), 6.67(1H, d, J=3.7Hz), 6.88(1H, d,  
10 J=3.7Hz).

Preparation Example A+-22. 2-Amino-N-(5-bromo-thiophen-2-ylmethyl)-  
6-methoxymethyl-nicotinamide

To a solution of C-(5-bromo-thiophen-2-yl)-methylaniline (250mg, 1.3mmol) in  
15 N,N-dimethylformamide (5mL) were added triethylamine (0.54mL, 3.9mmol),  
2-amino-6-methoxymethyl-nicotinic acid (240mg, 1.3mmol) described in Preparation  
Example A-11, and benzotriazol-1-yloxytris(dimethylamino)phosphonium  
hexafluorophosphate (690mg, 1.6mmol) sequentially on an ice bath, then, the  
solution was stirred for 2 days at room temperature. The reaction solution was  
20 partitioned in water and ethyl acetate. The organic layer was separated, the solvent  
was evaporated *in vacuo*. The residue was purified by silica gel column  
chromatography (heptane : ethyl acetate = 4 : 1), and the title compound (370mg,  
1.0mmol, 79%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.46(3H, s), 4.40(2H, s), 4.67(2H, d, J=5.7Hz),  
25 6.33(1H, br s), 6.38(2H, br s), 6.71(1H, d, J=7.9Hz), 6.79(1H, d, J=3.7Hz), 6.91(1H, d,

J=3.9Hz), 7.59(1H, d, J=7.9Hz).

Preparation Example A+-23. 2-Amino-N-(4-bromo-benzyl)-6-methoxymethyl-nicotinamide

5           The title compound (1.9g, 5.4mmol, 86%) was obtained as a pale yellow solid from 2-amino-6-methoxymethyl-nicotinic acid (1.2g, 6.3mmol) described in Preparation Example A-11 and 4-bromobenzylamine hydrochloride (1.5g, 6.9mmol) according to an analogous method to Preparation Example A+-22.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.46(3H, s), 4.40(2H, s), 4.55(2H, d, J=5.7Hz),  
10   6.30(1H, br s), 6.39(2H, br s), 6.70(1H, d, J=7.9Hz), 7.22(2H, d, J=8.2Hz), 7.48(2H, d, J=8.4Hz), 7.60(1H, d, J=7.9Hz).

Preparation Example A+-24. 2-Amino-N-(4-((Z)-2-ethoxy-vinyl)-benzyl)-6-methoxymethyl-nicotinamide

15           A mixture of (2-ethoxy-vinyl)-tributyl-stannane (37mg, 0.10mmol) synthesized according to WO02/018368, 2-amino-N-(4-bromo-benzyl)-6-methoxymethyl-nicotinamide (30mg, 0.086mmol) described in Preparation Example A+-23, tri-*o*-tolylphosphine (6.5mg, 0.021mmol), palladium acetate (0.96mg, 0.0043mmol), tetrabutylammonium chloride (24mg, 0.086mmol) and N-methylpyrrolidinone (1mL)  
20   was stirred at 125°C for 1 hour. The reaction solution was allowed to cool, an aqueous solution of potassium fluoride was added on an ice bath, the solution was filtered through Celite pad. The filtrate was partitioned with ethyl acetate. The organic layer was separated, washed with brine, then, evaporated *in vacuo*. The residue was purified by silica gel column chromatography (heptane : ethyl acetate =  
25   1 : 2), and the title compound (12mg, 0.035mmol, 35%) was obtained as a colorless

solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 1.36(3H, d, J=7.1Hz), 3.45(3H, s), 3.99(2H, q, J=7.1Hz), 4.39(2H, s), 4.55(2H, d, J=5.5Hz), 5.21(1H, d, J=7.0Hz), 6.21-6.23(2H, m), 6.40(2H, br s), 6.68(1H, d, J=7.9Hz), 7.26(2H, d, J=8.3Hz), 7.57-7.59(3H, m).

5

Preparation Example A+-25. 2-Amino-6-methoxymethyl-N-(3-hydroxybenzyl)-nicotinamide

2-Amino-6-methoxymethylcotinic acid (500mg, 2.74mmol),  
 3-hydroxybenzylamine (405mg, 3.29mmol),  
 10 benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (1.33mg, 3.01mmol) and triethylamine (555mg, 5.48mmol) were added to dimethylsulfoxide (20mL), and the solution was stirred for 15 minutes at 60°C. Water and ethyl acetate were added to the reaction mixture, which was then partitioned. The aqueous layer was extracted once with ethyl acetate, the ethyl acetate layers were  
 15 combined and the solvent was evaporated *in vacuo*. The residue was purified by silica gel column chromatography (hexane:ethyl acetate=1:1, then ethyl acetate). The resulting residue was purified by NH silica gel column chromatography (ethyl acetate). The resulting residue was solidified from hexane-ethyl acetate, and the title compound (490mg, 1.71mmol, 62.2%) was obtained as a pale yellow solid.  
 20 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.34 (3H, s), 4.30 (2H, s), 4.35 (2H, d, J=6.0 Hz), 6.61 (2H, d, J=8.0 Hz), 6.70 (1H, s), 6.71 (1H, d, J=8.0 Hz), 7.10 (1H, dd, J=8.0, 8.0 Hz), 7.13 (2H, brs), 7.99 (1H, d, J=8.0 Hz), 8.91 (1H, t, J=6.0 Hz), 9.31 (1H, s).

Preparation Example A+-26. 2-Amino-6-methoxymethyl-N-(4-hydroxybenzyl)-nicotinamide

25

The title compound (506mg, 1.76mmol, 64.3%) was obtained as a pale yellow solid from 2-amino-6-methoxymethylcotinic acid (500mg, 2.74mmol) and 4-hydroxybenzylamine (506mg, 4.11mmol) according to an analogous method to Preparation Example A+-25.

5 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.34 (3H, s), 4.29 (2H, s), 4.30 (2H, d, J=6.0), 6.59 (1H, d, J=7.6 Hz), 6.70 (2H, d, J=7.4 Hz), 7.10 (2H, d, J=7.4 Hz), 7.12 (2H, brs), 7.95 (1H, d, J=7.6 Hz), 8.84 (1H, t, J=6.0 Hz), 9.27 (1H, s).

Preparation Example AA-1. 3,5-Diamino-pyrazine-2-carboxylic acid methyl ester

10 To a solution of 3,5-diamino-6-chloro-pyrazine-2-carboxylic acid methyl ester (8.00g, 39.5mmol) in tetrahydrofuran (150mL) were added tetrakis(triphenylphosphine)palladium(0) (2.28g, 1.98mmol), formic acid (2.24mL, 59.3mmol) and triethylamine (16.5mL, 119mmol) at 0°C under nitrogen atmosphere, then, the solution was stirred at 125°C for 12 hours. The reaction solution was  
15 cooled to room temperature, and a solid precipitated. This solid was collected by filtration, and the title compound (10.7g, quantitatively) was obtained as a crude product of a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.70(3H, s), 6.95(2H, brs), 7.21(1H, s).

20 Preparation Example AA-2. 3,5-Diamino-pyrazine-2-carboxylic acid

Lithium hydroxide monohydrate (2.50g, 59.5mmol) was added to 3,5-diamino-pyrazine-2-carboxylic acid methyl ester (10.0g, 59.5mmol) described in Preparation Example AA-1 in a mixture solvent of tetrahydrofuran (100mL), methanol (10mL) and water (10mL) at room temperature. The solution was stirred at room  
25 temperature for 17 hours, then, an aqueous solution of 5N sodium hydroxide (15mL)



was added thereto, followed by further stirring for 4.5 hours under reflux. The reaction solution was cooled to room temperature, and partitioned in 5N hydrochloric acid solution and ethyl acetate. The organic layer was separated, washed with brine, and then, dried over anhydrous magnesium sulfate. The organic layer was filtered, then, the solvent was evaporated *in vacuo*, and the title compound (3.34g, 36%) was obtained as a crude product of a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 6.93(2H, brs), 7.20(1H, s).

Preparation Example B-1. 4-Amino-5-hydroxy-carbonyl-2-*n*-propylamino-pyrimidine

4-Amino-2-chloro-5-cyano-pyrimidine (300mg, 1.94mmol) and *n*-propyl amine (5g, 84.6mmol) were mixed and stirred for 10 minutes at 60°C. The reaction solution was directly purified by silica gel column chromatography, and a 2-propylamino compound (300mg, 1.69mmol, 101%) was obtained as a brown solid. This solid was suspended in concentrated sulfuric acid (3mL) and water (3mL), and the solution was stirred for 1.5 hours at 130°C. This mixture was alkalized with an aqueous solution of saturated sodium bicarbonate, then, the aqueous layer was washed with ethyl acetate. Then, this aqueous layer was neutralized with citric acid, and extracted with a mixture solvent of acetate-methanol. This organic layer was separated, then, the solvent was evaporated, and the title compound (44mg, 0.224mmol, 12%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 0.87 (3H, t, J=7.2 Hz), 1.49 (2H, qt, J=7.2, 7.2 Hz), 3.18 (2H, q, J=7.2Hz), 6.98 (2H, bs), 8.14 (1H, bs), 8.35 (1H, s).

Preparation Example C-1. Potassium salt of 2-Cyano-3-hydroxy-acrylic acid ethyl ester

To a solution of potassium ethoxide (9.8g, 116mmol) in ethanol (180mL) were added a solution of cyano ethyl acetate ester (13.2g, 117mmol) and formic acid ethyl ester (30g, 405mmol) in ethanol (20mL), and the solution was stirred for 2 hours under reflux. The reaction solution was allowed to room temperature, the precipitated solid was collected by filtration and dried, and the title compound (18g, 100mmol, 85%) was obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) :1.13 (3H, t, J=7.2 Hz), 3.96 (2H, q, J=7.2Hz), 9.18 (1H, s).

10 Preparation Example C-2. 1-(Pyrrolidino)-2-(2-carbethoxy-2-cyanoethylene)cyclopentene

Potassium salt of 2-cyano-3-hydroxy-acrylic acid ethyl ester described in Preparation Example C-1 (18g, 0.1mol) was dissolved in dichloromethane (80mL), phosphorus pentachloride (20.9g, 0.1mol) was added, and the solution was stirred for 15 2 hours under reflux. After the reaction was completed, the residue obtained by evaporating dichloromethane was subjected to distillation *in vacuo*, and ethyl(chloromethylene)cyanoacetate (9.5g, 56mmol) was obtained.

To a solution of 1-pyrrolidinocyclopentene (10.2g, 74mmol) and triethylamine (10mL, 72mmol) in dichloromethane (200mL) was added a solution of the resulting 20 ethyl(chloromethylene)cyanoacetate (9.5g, 56mmol) in dichloromethane (20mL) dropwise while stirring at from -20°C to -25°C. The solution was stirred at room temperature for 50 minutes, water (20mL) was added thereto, followed by further stirring for 5 minutes. The reaction mixture was partitioned, the organic layer was dried over anhydrous magnesium sulfate, then concentrated, and the title compound 25 (6g, 23mmol, 23%) was obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) :1.19 (3H, t, J=7.2 Hz), 1.76-1.86 (2H, m), 1.86-2.04 (4H, m), 2.73 (2H, t, J=7.6 Hz), 2.88 (2H, t, J=7.2 Hz), 3.60-3.71 (4H, m), 4.09 (2H, q, J=7.2Hz), 7.97 (1H, brs).

5 Preparation Example C-3. 1-Amino-2-(2-carbethoxy-2-cyanoethylene)cyclopentene

1-(Pyrrolidino)-2-(2-carbethoxy-2-cyanoethylene)cyclopentene (6g, 23mmol) described in Preparation Example C-2 was dissolved in ethanol saturated with ammonia (75mL; was saturated at room temperature using ammonia gas), and the solution was stirred for 19 hours at room temperature. The reaction solution was concentrated and the title compound (4.8g, 23mmol, quantitatively) was obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) :1.21 (3H, t, J=7.2 Hz), 1.74-1.83 (2H, m), 2.48-2.54 (2H, m), 2.72-2.78 (2H, m), 4.12 (2H, q, J=7.2Hz), 8.09-8.47 (1H, brs).

Preparation Example C-4. 2-Amino-6,7-dihydro-5H-cyclopenta[b]pyridine-3-

15 carboxylic acid ethyl ester

1-Amino-2-(2-carboethoxy-2-cyanoethylene)cyclopentene (0.8g, 3.9mmol) described in Preparation Example C-3 was dissolved in ethanol (27mL), sodium ethoxide (0.12g, 1.8mmol) was added thereto, followed by stirring for 16 hours under reflux. The reaction mixture was allowed to room temperature, poured into brine, and the solution was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, then, the residue obtained by concentration was purified by silica gel column chromatography (ethyl acetate : hexane = 1 : 2), and the title compound (0.63g, 3.1mmol, 79%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 1.38 (3H, t, J=7.2Hz), 2.04-2.13 (2H, m), 2.79-2.88 (4H, m), 4.32 (2H, q, J=7.2 Hz), 6.16-6.32 (2H, brs), 7.96 (1H, s).

Preparation Example C-5. 2-Amino-6,7-dihydro-5H-cyclopenta[b]pyridine-3-carboxylic acid

2-Amino-6,7-dihydro-5H-cyclopenta[b]pyridine-3-carboxylic acid ethyl ester  
5 (0.2g, 0.97mmol) described in Preparation Example C-4 was dissolved in ethanol (15mL), an aqueous solution of 1N sodium hydroxide (7.5mL) was added thereto, followed by heating at 100°C for 30 minutes. The reaction solution was allowed to room temperature, then cooled on an ice bath, and neutralized with 1N hydrochloric acid. The precipitated solid was collected by filtration, rinsed with water, then dried,  
10 and the title compound (143mg, 0.8mmol, 83%) was obtained.  
<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) :1.94-2.03 (2H, m), 2.71-2.76 (4H, m), 7.84 (1H, s).

Preparation Example D-1. [1,5]Naphthylidene-2-carboxylic acid

15 5-Amino-2-chloro pyridine (10g, 0.078mol) and oxalacetic acid diethyl ester (14.7g, 0.078mol) were stirred at 90°C for 7 hours. The reaction solution was allowed to room temperature, ethyl acetate was added thereto, the precipitated yellow solid was removed by filtration, and the filtrate was concentrated. The residue was purified by silica gel column chromatography (hexane : ethyl acetate),  
20 and 2-(6-chloro-pyridin-3-ylamino)-buto-2-enedioic acid diethyl ester (4.8g, 21%) was obtained as a yellow oil.

The resulting 2-(6-chloro-pyridin-3-ylamino)-buto-2-enedioic acid diethyl ester (4.8g, 16.1mmol) was added to Dowtherm A (Dowtherm A; trademark) (300mL), and the solution was stirred at 210°C for 5 hours. The reaction solution was allowed to  
25 room temperature, hexane was added thereto, the precipitated solid was collected,

washed with hexane, then, dried *in vacuo* to obtain 6-chloro-4-hydroxy-[1,5]naphthylidene-2-carboxylic acid ethyl ester (1.38g, 34%) as a pale brown solid.

To the resulting 6-chloro-4-hydroxy-[1,5]naphthylidene-2-carboxylic acid ethyl ester (502mg, 1.99mmol) was added thionyl chloride (10mL), and the solution was stirred for 7 hours under reflux. Excess thionyl chloride was evaporated *in vacuo*, 4,6-dichloro-[1,5]naphthylidene-2-carboxylic acid ethyl ester (522mg, 97%) was obtained as a pale brown solid. Under nitrogen atmosphere, a portion of the resulting solid (200mg, 0.738mmol) was dissolved in dimethylsulfoxide (30mL), tetrakis(triphenylphosphine)palladium(0) (171mg, 0.148mmol) and formic acid sodium (251mg, 3.69mmol) were added thereto, followed by stirring at 100°C for 4 hours. The reaction solution was allowed to room temperature, ethyl acetate and water were added for partitioning, the organic layer was washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated, then, the residue was purified by silica gel column chromatography (hexane : ethyl acetate), and [1,5]naphthylidene-2-carboxylic acid ethyl ester (49mg, 33%) was obtained as a colorless solid.

The resulting solid was dissolved in methanol (1.0mL), an aqueous solution of 1N sodium hydroxide (0.3mL) was added thereto, followed by stirring at room temperature for 30 minutes. Water was added to the reaction solution, the pH was adjusted from 3 to 4 using 1N hydrochloric acid, the solution was extracted with ethyl acetate, dried over anhydrous magnesium sulfate, and the solvent was evaporated to obtain the title compound (29mg, 69%) as a white solid.

Preparation Example E-1. Quinoline-6-carboxylic acid cyanomethyl-amide

To a solution of quinoline-6-carboxylic acid (500mg, 2.9mmol) and amino acetonitrile hydrochloride (320mg, 3.4mmol) in N,N-dimethylformamide (10mL) were added benzotriazol-1-yl-tris(dimethylamino)phosphonium hexafluorophosphate (1.5g, 3.48mmol) and triethylamine (1.2mL, 8.7mmol), and the solution was stirred at 60°C for 10 minutes. Ethyl acetate and water was added to the reaction solution, which was then partitioned, the organic layer was washed twice with water. Silica gel was added to the organic layer, the solvent was evaporated *in vacuo* for adsorption, purified by NH silica gel column chromatography (ethyl acetate), and the title compound (420mg, 2.0mmol, 68.9%) was obtained as a light brown solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.43 (2H, d, J=5.6Hz), 7.65 (1H, dd, J=4.0, 8.4Hz), 8.14 (1H, d, J=8.8Hz), 8.18-8.22 (1H, m), 8.30-8.35 (1H, m), 8.58 (1H, d, J=1.2Hz), 9.02-9.05 (1H, m), 9.49 (1H, t, J=5.6Hz).

#### Preparation Example E+-1. Quinoline-6-carboxylic acid 4-hydroxybenzylamide

Trifluoroacetic acid (5mL) and thioanisole (3 drops) were added to quinoline-6-carboxylic acid 4-benzyloxybenzylamide (2.67g, 7.25mmol) synthesized in Example E-8, the solution was stirred at room temperature for 14 hours, followed by stirring at 50°C for 4 hours, lastly, and the solution was stirred at 70°C for 3 hours. The reaction solution was allowed to room temperature, then, was neutralized with an aqueous solution of saturated sodium bicarbonate, extracted with ethyl acetate and dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by NH silica gel column chromatography (ethyl acetate), and the title compound (433mg, 22%) was obtained as a colorless solid.

Preparation Example E+-2. Quinoline-6-carboxylic acid

(5-bromo-furan-2-ylmethyl)-amide

The title compound (1.0g, 3.0mmol, 75.5%) was obtained as a white solid from C-(5-bromo-furan-2-yl)-methylamine (700mg, 4.0mmol) and quinoline-6-carboxylic acid (700mg, 4.0mmol) according to an analogous method to

## 5 Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.49 (2H, d, J=5.6Hz), 6.38-6.41 (1H, m), 6.50 (1H, d, J=3.6Hz), 7.60 (1H, dd, J=4.0, 8.4Hz), 8.06 (1H, d, J=8.8Hz), 8.17 (1H, dd, J=2.0, 8.8Hz), 8.44-8.48 (1H, m), 8.52 (1H, d, J=2.0Hz), 8.97 (1H, dd, J=1.6, 4.0Hz), 9.23 (1H, t, J=5.6Hz).

10

Preparation Example E+-3. Quinoline-6-carboxylic acid 4-benzylamino-benzylamide

4-Benzylamino-benzonitrile (472mg, 2.27mmol) described in Preparation Example 89 was dissolved at 0°C in tetrahydrofuran (20mL), and lithium aluminum hydride (430mg, 11.3mmol) was added thereto. The solution was stirred overnight at room temperature, then, at 0°C, water (430μl), an aqueous solution of 5N sodium hydroxide (430μl) and water (1.29mL) were sequentially added to the solution. The reaction solution was filtered through Celite pad, the solvent was then evaporated *in vacuo*, (4-aminomethyl-phenyl)-benzylamine (475mg, 2.24mmol, 99%) was obtained as an oil.

20 The resulting (4-aminomethyl-phenyl)-benzylamine (162mg, 0.763mmol), quinoline-6-carboxylic acid (132mg, 0.736mmol), benzotriazol-1-yl-tris(dimethylamino)phosphonium hexafluorophosphate (506mg, 1.14mmol) and triethylamine (319μl, 2.29mmol) were dissolved in N,N-dimethylformamide (4.0mL), and the solution was stirred for 2 hours at room temperature. Water was added to the reaction solution, which was then extracted

25

with ethyl acetate, the organic layer was washed with water and brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 1 : 1), and the title compound (224mg, 0.610mmol, 80%) was obtained as a white solid.

5 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.23 (2H, d, J=6.0Hz), 4.33 (2H, d, J=6.0Hz), 6.18 (1H, t, J=6.1Hz), 6.51 (2H, d, J=8.6Hz), 7.03 (2H, d, J=8.6Hz), 7.18 (1H, t, J=7.0Hz), 7.25-7.34 (4H, m), 7.58 (1H, dd, J=4.1, 8.3Hz), 8.04 (1H, d, J=8.8Hz), 8.16 (1H, dd, J=1.8, 9.0Hz), 8.43 (1H, d, J=7.0Hz), 8.49 (1H, d, J=2.0Hz), 8.95 (1H, dd, J=1.8, 5.0Hz), 9.04 (1H, t, J=5.5Hz).

10

Preparation Example E+-4. Quinoline-6-carboxylic acid 3-hydroxybenzylamide

Thioanisole (1.7mL, 14mmol) was added to a mixture of quinoline-6-carboxylic acid 3-benzyloxybenzylamide (1.3g, 3.6mmol) and trifluoroacetic acid(8mL) on an ice bath, and the solution was stirred for 4 hours at  
15 room temperature. The solvent was evaporated *in vacuo*, then water, an aqueous solution of saturated sodium bicarbonate, ethyl acetate and tetrahydrofuran were added to the residue for extraction, the organic layer was washed with brine, then, dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, recrystallized from ethyl acetate-methanol, and the title compound (0.64g, 2.3mmol,  
20 64%) was obtained as a white crystal.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.45 (2H, d, J=5.9Hz), 6.60-6.63 (1H, m), 6.75-6.77 (2H, m), 7.10 (1H, t, J=8.1Hz), 7.60 (1H, dd, J=4.2, 8.2Hz), 8.07 (1H, d, J=8.8Hz), 8.20 (1H, dd, J=2.0, 8.8Hz), 8.45-8.47 (1H, m), 8.54 (1H, d, J=1.8Hz) , 8.97 (1H, dd, J=1.7, 4.2Hz), 9.23 (1H, t, J=5.8Hz), 9.33 (1H, s).

25



Preparation Example E+-5. Quinoline-6-carboxylic acid 4-phenylethynyl-benzylamide

Quinoline-6-carboxylic acid 4-bromobenzylamide(1.3g, 68%) was obtained from quinoline-6-carboxylic acid (1.0g, 5.8mmol) and 4-bromobenzylamine hydrochloride (1.3g, 5.8mmol) according to an analogous method to Preparation  
 5 Example A+-5 (with the proviso that reaction was carried out at 80°C).

N,N-diisopropylethylamine (0.31mL, 1.8mmol) was added to a mixture of the resulting quinoline-6-carboxylic acid 4-bromobenzylamide (200mg, 0.59mmol) , ethynylbenzene(0.077mL, 0.70mmol), copper(I) iodide (catalytic amount), tetrakis(triphenylphosphine)palladium(0) (68mg, 0.059mmol) and  
 10 N-methylpyrrolidinone (4mL), and the solution was stirred at 100°C for 30 minutes, and at 120°C for 50 minutes. After cooling, water, ethyl acetate, tetrahydrofuran and 29% aqueous ammonia solution were added to the reaction mixture for extraction, the organic layer was washed with brine, then, the solvent was evaporated *in vacuo*. The residue was purified by silica gel column chromatography (hexane : ethyl acetate  
 15 = 1 : 4), then, the resulting crudely purified product was washed with diethyl ether to obtain the title compound (50mg, 0.14mmol, 24%) as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.57 (2H, d, J=5.7Hz), 7.39-7.41 (5H, m), 7.52 (4H, d, J=6.8Hz), 7.60 (1H, dd, J=3.8, 8.1Hz), 8.08 (1H, d, J=8.6Hz), 8.21 (1H, d, J=8.8Hz), 8.47 (1H, d, J=7.9Hz), 8.56 (1H, s), 8.97 (1H, d, J=4.0Hz), 9.33 (1H, brs).

20

Preparation Example E+-6. Quinoline-6-carboxylic acid  
4-[1,3]dioxolan-2-yl-benzylamide

The title compound (1.31g, 3.92mmol, 77%) was obtained as a white solid from 4-[1,3]dioxolan-2-yl-benzylamine described in Preparation Example 120 (970mg,  
 25 5.60mmol) and quinoline-6-carboxylic acid (913mg, 5.09mmol) according to an

analogous method to Preparation Example Q+-1.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.90-4.02 (4H, m), 4.54 (2H, d, J=5.9Hz),  
5.69 (1H, s), 7.35-7.40 (4H, m), 7.60 (1H, dd, J=4.2, 8.2Hz), 8.07 (1H, d, J=9.0Hz),  
8.2 (1H, dd, J=1.9, 9.0Hz), 8.46 (1H, d, J=8.1Hz), 8.54 (1H, d, J=1.5Hz), 8.97 (1H, dd,  
5 J=1.7, 4.2Hz), 9.29 (1H, t, J=5.7Hz).

Preparation Example E+-7. Quinoline-6-carboxylic acid 4-formyl-benzylamide

Quinoline-6-carboxylic acid 4-[1,3]dioxolan-2-yl-benzylamide described in  
Preparation Example E+-6 (1.30g, 3.89mmol) was dissolved in a mixture solution of  
10 tetrahydrofuran (20mL), water (10mL) and sulfuric acid (3mL), the solution was  
stirred for 2 hours under reflux. An aqueous solution of saturated sodium  
bicarbonate was added to the reaction solution at 0°C, the solution was extracted  
with ethyl acetate and tetrahydrofuran. The organic layer was washed with brine,  
dried over anhydrous magnesium sulfate, and the solvent was evaporated *in vacuo*.  
15 Ethyl acetate was added to the residue, the precipitated solid was filtered, and the  
title compound (700mg, 2.41mmol, 62%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.63 (2H, d, J=5.7Hz), 7.57 (2H, d, J=7.9Hz),  
7.61 (1H, t, J=4.1Hz), 7.88 (2H, dd, J=1.8, 8.4Hz), 8.09 (1H, d, J=8.8Hz), 8.21 (1H,  
dd, J=2.0, 8.8Hz), 8.48 (1H, d, J=7.9Hz), 8.56 (1H, d, J=1.5Hz), 8.98 (1H, dd, J=1.8,  
20 4.2Hz), 9.39 (1H, t, J=6.0Hz), 9.97 (1H, s).

Preparation Example E+-8. Quinoline-6-carboxylic acid 3-bromobenzylamide

The title compound (1.4g, 4.0mmol, 70%) was obtained as a white solid from  
3-bromobenzylamine hydrochloride (1.3g, 5.8mmol) and quinoline-6-carboxylic acid  
25 (1.0g, 5.8mmol) according to an analogous method to Preparation Example Q+-1.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.68 (2H, d, J=5.7Hz), 6.75 (1H, brs), 7.23 (1H, t, J=7.8Hz), 7.31-7.34 (1H, m), 7.43-7.45 (1H, m), 7.47 (1H, dd, J=4.2, 8.2Hz), 7.53-7.54 (1H, m), 8.07 (1H, dd, J=2.0, 8.8Hz), 8.16 (1H, d, J=8.8Hz), 8.22-8.24 (1H, m), 8.34 (1H, d, J=1.8Hz), 8.99 (1H, dd, J=1.8, 4.2Hz).

5

Preparation Example E+-9. Quinoline-6-carboxylic acid 4-bromobenzylamide

The title compound (1.3g, 3.9mmol, 68%) was obtained as a white solid from 4-bromobenzylamine hydrochloride (1.3g, 5.8mmol) and quinoline-6-carboxylic acid (1.0g, 5.8mmol) according to an analogous method to Preparation Example Q+-1.

10 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.67 (2H, d, J=5.9Hz), 6.63 (1H, brs), 7.26-7.29 (2H, m), 7.47-7.51 (3H, m), 8.06 (1H, dd, J=2.0, 8.8Hz), 8.16 (1H, d, J=8.8Hz), 8.22-8.25 (1H, m), 8.33 (1H, d, J=2.0Hz), 8.99 (1H, dd, J=1.7, 4.2Hz).

Preparation Example E+-10. Quinoline-6-carbothioic acid 4-benzyloxy-benzylamide

15 To a solution of 6-quinoline carboxylic acid (100mg, 0.577mmol) in tetrahydrofuran (50mL) was added N,N'-dicyclohexylcarbodiimide (1.90g, 11.7mmol), and the solution was stirred for 1 hour at room temperature. Then, a solution of 4-benzyloxybenzylamine described in Preparation Example 1 (2.49g, 11.7mmol) in tetrahydrofuran was added thereto, followed by stirring overnight at room  
20 temperature. The solvent was evaporated, the residue was purified by NH silica gel column chromatography (hexane:ethyl acetate), and quinoline-6-carboxylic acid 4-benzyloxy-benzylamide (4.31g, quantitatively) was obtained as a white solid.

A mixture of the resulting quinoline-6-carboxylic acid 4-benzyloxy-benzylamide (310mg, 0.84mmol), 2,4-bis  
25 (4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulphide (Lawesson's reagent)

(1.4g, 3.4mmol) and tetrahydrofuran (10mL) was refluxed for 1 hour. After cooling, the solvent was evaporated *in vacuo*, dichloromethane was added to the residue, which was purified by NH silica gel column chromatography (hexane : ethyl acetate = 1 : 1), and the title compound (55mg, 0.14mmol, 17%) was obtained as a pale yellow solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.93 (2H, d, J=4.6Hz), 5.09 (2H, s), 6.99 (2H,d, J=8.8Hz), 7.26-7.44 (7H, m) , 7.58 (1H, dd, J=4.2, 8.2Hz) , 8.01 (1H, d, J=9.0Hz) , 8.13 (1H, dd, J=2.1, 8.9Hz), 8.29 (1H, d, J=1.8Hz) , 8.46 (1H, d, J=8.2Hz) , 8.94 (1H, dd, J=1.6, 4.2Hz), 10.9(1H, brs).

#### Preparation Example F-1. 3-Acetyl-4-amino-benzoic acid methyl ester

To a solution of 4-amino-3-iodo-benzoic acid methyl ester (11.30g, 40.77mmol) in toluene (300mL) were added tributyl(1-ethoxyvinyl)tin (16.5mL, 48.9mmol) and tetrakis(triphenylphosphine)palladium(0) (9422mg, 8.154mmol) under nitrogen atmosphere, and the solution was stirred at 105°C for 7 hours. After cooling to room temperature, water was added, the mixture was extracted with ethyl acetate-tetrahydrofuran, the organic layer was washed with water, and then, evaporated. The residue was dissolved in 280mL of tetrahydrofuran, 2N hydrochloric acid (80mL) was added thereto, followed by stirring for 3 hours at room temperature. The reaction mixture was cooled on an ice bath, an aqueous solution of 2N sodium hydroxide (80mL) was added, an aqueous solution of saturated sodium bicarbonate was further added, and the solution was extracted with ethyl acetate. An aqueous solution of 10% potassium fluoride was added to the organic layer, and the solution was stirred for 3 hours at room temperature. The organic layer was separated, washed with brine, then evaporated, the residue was purified by silica gel

column chromatography (hexane-ethyl acetate), and title compound (6.42g, 33.2mmol, 81.4%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 2.64 (3H, s), 3.89 (3H, s), 6.63 (1H, d, J=8.8Hz), 7.91 (1H, dd, J=2.0, 8.8Hz), 8.47 (1H, d, J=2.0Hz).

5

Preparation Example F-2. 4-Oxo-1,4-dihydro-cinnoline-6-carboxylic acid methyl ester

To a solution of 3-acetyl-4-amino-benzoic acid methyl ester (2063mg, 10.68mmol) in acetic acid (39mL) was added sulfuric acid (6.5mL) on an ice bath, then, an aqueous solution (6.5mL) of sodium nitrite (922mg, 13.35mmol) was added thereto, followed by stirring on the ice for 1 hour, and at room temperature for 2 days. The reaction mixture was concentrated until it reached half of the amount, water was added, then an aqueous solution of 2N sodium hydroxide was added on an ice bath to adjust the pH to 5. The insoluble matter was separated by filtration, then, the filtrate was extracted with ethyl acetate, the organic layer was dried over sodium sulfate and then evaporated. Diethyl ether was added to the resulting residue for solidification, the resulting solid was washed with diethyl ether, and 365mg of the title compound (1.78mmol, 16.6%) was obtained.

10

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.88 (3H, s), 7.66 (1H, d, J= 8.8Hz), 7.82 (1H, s), 8.24 (1H, d, J=8.8Hz), 8.58 (1H, s), 13.7(1H, brs).

15

Preparation Example F-3. 4-Chloro-cinnoline-6-carboxylic acid methyl ester

Thionyl chloride (5mL, 68.5mmol) and N,N-dimethylformamide (3 drops) were added to 4-oxo-1,4-dihydro-cinnoline-6-carboxylic acid methyl ester (212mg, 1.04mol), and the solution was stirred for 1.15 hours under reflux. Toluene was added to the reaction mixture, which was then evaporated *in vacuo*. Ethyl acetate

20

was added to the residue the organic layer was washed with an ice water, dried over anhydrous magnesium sulfate, then, evaporated *in vacuo*, and the title compound (192mg, 0.862mmol, 82.9%) was obtained. This was used in the next reaction without purification.

5 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.97 (3H, s), 8.43 (1H, d, J=8.8Hz), 8.66 (1H, d, J=8.8Hz), 8.72 (1H, s), 9.73 (1H, s).

#### Preparation Example F-4. Cinnoline-6-carboxylic acid methyl ester

To a solution of 4-chloro-cinnoline-6-carboxylic acid methyl ester (192mg, 0.863mmol) in dimethylsulfoxide (30mL) were added sodium formate (70mg, 1.04mmol), tetrakis(triphenylphosphine)palladium(0) (198mg, 0.702mmol) and N,N-diisopropylethylamine (0.21mL, 1.21mmol), and the mixture was stirred for 1.5 hours at 90°C. After cooling to room temperature, water was added, ethyl acetate extraction was carried out, the organic layer was washed with water and brine, then, after drying over anhydrous sodium sulfate, the organic layer was evaporated. The resulting solid was washed with diethyl ether, then, purified by silica gel column chromatography (hexane-ethyl acetate), and the title compound (16mg, 0.089mmol, 10%) was obtained.

20 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.03 (3H, s), 7.97 (1H, dd, J=0.8, 6.0Hz), 8.42 (1H, J=0.8, 8.0Hz), 8.59-8.63 (2H, m), 9.43(1H, dd, J= 0.8, 6.0Hz).

#### Preparation Example G-1. Isoquinoline-6-carboxylic acid

A solution prepared by adding (4-bromobenzylidene)-(2,2-diethoxyethyl) amine (synthesized from 4-bromobenzaldehyde, according to the method described in J. Org. Chem., vol. 48, 3344-3346 (1983)) (51.4g, 0.189mmol) to an ice-cold

concentrated sulfuric acid (20g) was added to a solution prepared by adding diphosphorus pentoxide (40g) to an ice-cold concentrated sulfuric acid (360g), and the solution was stirred at 160°C for 2 hours. The reaction solution was gradually cooled to 0°C, the solution was filtered through Celite pad, the filtrate was neutralized  
5 with sodium carbonate. This solution was further filtrated through Celite pad, this filtrate was extracted with ethyl acetate and dried over anhydrous magnesium sulfate. The solvent was evaporated, and the residue was purified by silica gel column chromatography (hexane:ethyl acetate), and 6-bromoisoquinoline (482mg, 1.2%) was obtained as an orange oil.

10       Next, to a solution of 6-bromoisoquinoline (382mg, 1.84mmol) in N,N-dimethylformamide (3.8mL) were added zinc cyanide (431mg, 3.67mmol) and tetrakis(triphenylphosphine)palladium(0) (42mg, 0.0367mmol) under nitrogen atmosphere, and the mixture was stirred at 100°C for 1 hour. Tetrakis(triphenylphosphine)palladium(0) (42mg, 0.0367mmol) was further added,  
15 and the mixture was stirred for 2.5 hours at 100°C. The reaction mixture was allowed to room temperature, ethyl acetate and water were added for extraction, the organic layer was washed with water and dried over anhydrous magnesium sulfate. The residue was purified by silica gel column chromatography (hexane : ethyl acetate), and isoquinoline-6-carbonitrile (234mg, 83%) was obtained as a yellow  
20 solid.

      Lastly, isoquinoline-6-carbonitrile (51mg, 0.331mmol) was dissolved in diethyleneglycol (1.0mL), potassium hydroxide (9mg, 0.166mmol) was added thereto, followed by stirring at 160°C for 3 hours. The reaction mixture was allowed to room temperature, neutralized using hydrochloric acid, extracted with ethyl acetate, dried  
25 over anhydrous magnesium sulfate, then, the solvent was evaporated. Water was

added to the residue, the precipitated solid was collected, washed with water, dried *in vacuo*, so as to obtain the title compound (12mg, 21%) as a yellow solid.

Preparation Example H-1. 4-Chloro-quinazoline-6-carboxylic acid ethyl ester

5           The title compound (380mg, 1.61mmol, 88%) was obtained from 4-oxo-dihydroquinazoline-6-carboxylic acid ethyl ester (396mg, 1.81mmol) according to an analogous method to Preparation Example F-3.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) :1.34(3H, t, J=7.2Hz), 4.35 (2H, q, J=7.2Hz), 7.78 (1H, d, J=8.4Hz), 8.29 (1H, dd, J=2.0, 8.4Hz), 8.37 (1H, s), 8.64 (1H, d, J=2.0Hz).

10

Preparation Example H-2. Quinazoline-6-carboxylic acid ethyl ester

          The title compound (79mg, 0.39mmol, 24%) was obtained from 4-chloro-quinazoline-6-carboxylic acid ethyl ester (380mg, 1.61mmol) according to an analogous method to Preparation Example F-4.

15

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) :1.47 (3H, t, J=7.6Hz), 4.48 (2H, t, J=7.6Hz), 8.11 (1H, d, J=8.8Hz), 8.53 (1H, dd, J=2.0, 8.8Hz), 8.71 (1H, d, J=2.0Hz), 9.42 (1H, s), 9.52 (1H, s).

20   Preparation Example H-3. Quinazoline-6-carboxylic acid

          To a solution of quinazoline-6-carboxylic acid ethyl ester (79mg, 0.391mmol) in ethanol (4mL) was added an aqueous solution of 1N sodium hydroxide (4mL), and the solution was stirred for 1 hour at room temperature. 1N Hydrochloric acid was added to the reaction mixture to adjust the pH to 4, and the solution was evaporated *in vacuo*. Ethanol was added to the residue, and the organic layer was

25



concentrated. The residue was dissolved in an ethyl acetate-tetrahydrofuran mixture solvent, dried over anhydrous magnesium sulfate, then, evaporated *in vacuo*, and quinazoline-6-carboxylic acid (15mg, 0.086mmol, 22%) was obtained. This was used in the next reaction without purification.

- 5 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 8.09 (1H, d, J=8.8Hz), 8.44 (1H, dd, J=2.0, 8.8Hz), 8.83 (1H, d, J=2.0Hz), 9.39 (1H, s), 9.79 (1H, s).

#### Preparation Example I-1. Quinoxaline-6-carboxylic acid

To a solution of quinoxaline-6-carboxylic acid methyl ester (2084mg, 11.07mmol) in ethanol (25mL) was added an aqueous solution of 1N sodium hydroxide (25mL), and the solution was stirred for 4 hours under reflux. 1N Hydrochloric acid was added to the reaction mixture to adjust the pH to 4, then, the precipitated solid was collected by filtration, washed with water and isopropanol, then dried to obtain the title compound (1477mg, 8.479mmol, 76.6%) as a solid.

- 15 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 8.18 (1H, d, J=8.4Hz), 8.29 (1H, dd, J=8.4, 1.2Hz), 8.61 (1H, d, J=1.2Hz), 9.00-9.07 (2H, m).

#### Preparation Example J-1. 2,2-Dimethyl-N-pyridin-2-yl-propionamide

2-Aminopyridine (3.1g, 33mmol) and triethylamine(6.9mL, 49mmol) was dissolved in dichloromethane (40mL), 2,2-dimethylpropionyl chloride (4.5mL, 36mmol) was added on an ice bath, and the solution was stirred for 2 hours at the same temperature. Water was added thereto for extraction, the organic layer was sequentially washed with an aqueous solution of saturated sodium bicarbonate and brine, then, dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, and the title compound (6.0g, 34mmol, 102%) was obtained as a white

solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) :1.27 (9H, s), 7.03 (1H, ddd, J=1.1, 4.9, 7.3Hz), 7.68-7.72 (1H, m), 8.02 (1H, s), 8.23-8.27 (2H, m).

5 Preparation Example J-2. N-(3-formylpyridin-2-yl)-2,2-dimethylpropionamide

To a mixture solution of *tert*-butyl lithium (1.5M pentane solution, 10mL, 15mmol) and diethyl ether (50mL) was added a solution of 2,2-dimethyl-N-pyridin-2-yl-propionamide described in Preparation Example J-1 (900mg, 5.0mmol) in diethyl ether (10mL) dropwise at -78°C, and the solution was stirred for 90 minutes at the same temperature. At the same temperature, morpholine-4-carbaldehyde (1.0mL, 10mmol) was added dropwise, and the solution was warmed gradually to room temperature. Water and tetrahydrofuran were added to the reaction mixture for extraction, the organic layer was washed with brine, then, the solvent was evaporated *in vacuo*. The residue was purified by silica gel column chromatography (hexane : ethyl acetate = 1 : 4), and the title compound (880mg, 4.3mmol, 85%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) :1.38 (9H, s), 7.21 (1H, dd, J=4.9, 7.6Hz), 8.05 (1H, dd, J=2.0, 7.5Hz), 8.69 (1H, dd, J=2.0, 4.9Hz), 9.94 (1H, s), 10.9 (1H, brs).

20 Preparation Example J-3. (2-Aminopyridin-3-yl)-methanol

A mixture solution of N-(3-formylpyridin-2-yl)-2,2-dimethylpropionamide described in Preparation Example J-2 (500mg, 2.4mmol) and an aqueous solution of 5N sodium hydroxide (7mL) was refluxed for 90 minutes. After cooling, ethyl acetate and tetrahydrofuran were added for extraction, the organic layer was washed with brine, then, the solvent was evaporated *in vacuo*. The residue was purified by

silica gel column chromatography (methanol : ethyl acetate = 1 : 5), and the title compound (160mg, 1.2mmol, 53%) was obtained as a pale yellow solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.31 (2H, s), 5.13 (1H, brs), 5.62 (2H, s), 6.51 (1H, dd, J=5.0, 7.3Hz), 7.34-7.36 (1H, m), 7.81-7.82 (1H, m).

5

Preparation Example J-4. 2-Aminopyridine-3-carbaldehyde

To a mixture of (2-aminopyridin-3-yl)-methanol described in Preparation Example J-3 (130mg, 1.1mmol) and dichloromethane (10mL) was added manganese dioxide (1.3g, 15mmol) at room temperature, and the solution was stirred overnight.

10 The reaction solution was filtered through Celite pad, then, the solvent was evaporated *in vacuo*, and the title compound (108mg, 0.88mmol, 83%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 6.75 (1H, dd, J=4.9, 7.5Hz), 7.83 (1H, dd, J=1.9, 7.5Hz), 8.23 (1H, dd, J=1.9, 4.9Hz), 9.86 (1H, s).

15

Preparation Example J-5. 2-Hydroxy-[1,8]naphthylidene-3-carboxylic acid ethyl ester

To a mixture of 2-aminopyridine-3-carbaldehyde described in Preparation Example J-4 (8.0mg, 0.066mmol) and ethanol (2mL) were added diethylmalonate (0.50mL, 3.3mmol) and piperidine (0.20mL, 2.0mmol), and the solution was stirred overnight at 70°C. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (methanol : ethyl acetate = 1 : 10), and the title compound (9.2mg, 0.042mmol, 64%) was obtained as a white solid.

20

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 1.43 (3H, t, J=7.1Hz), 4.45 (2H, q, J=7.1Hz), 7.28 (1H, dd, J=4.9, 7.8Hz), 8.04 (1H, dd, J=1.7, 7.9Hz), 8.48 (1H, s), 8.87-8.88 (1H, m), 12.16 (1H, brs).

25

Preparation Example J-6. 2-Trifluoromethane sulfonyloxy-[1,8]naphthylidene-3-carboxylic acid ethyl ester

To a mixture of 2-hydroxy-[1,8]naphthylidene-3-carboxylic acid ethyl ester described in Preparation Example J-5 (95mg, 0.44mmol), dichloromethane(4mL) and N,N-dimethylformamide (0.5mL) were added N-phenyl-bis(trifluoromethanesulfonimide) (230mg, 0.65mmol), triethylamine (0.18mL, 1.3mmol) and catalytic amount of 4-(dimethylamino)pyridine, and the solution was stirred for 2.5 hours at room temperature. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 1 : 1), and the title compound (22mg, 0.063mmol, 14%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 1.49 (3H, t, J=7.1Hz), 4.54 (2H, q, J=7.1Hz), 7.69 (1H, dd, J=4.3, 8.2Hz), 8.41 (1H, dd, J=2.0, 8.4Hz), 9.09 (1H, s), 9.28 (1H, dd, J=2.0, 4.2Hz).

Preparation Example J-7. [1,8]Naphthylidene-3-carboxylic acid ethyl ester

To a mixture of 2-trifluoromethanesulfonyloxy-[1,8]naphthylidene-3-carboxylic acid ethyl ester described in Preparation Example J-6 (22mg, 0.063mmol), tetrakis(triphenylphosphine)palladium(0) (7.3mg, 0.0063mmol), and 1-methyl-2-pyrrolidinone (1.5mL) were added N,N-diisopropylethylamine (0.033mL, 0.19mmol) and formic acid (0.0036mL, 0.094mmol), and the solution was stirred for 45 minutes at 100°C. After cooling, filtration was carried out using NH silica gel, and the filtrate was evaporated *in vacuo*. Water, ethyl acetate and tetrahydrofuran were added to the residue for extraction, the organic layer was washed with brine, the

solvent was then evaporated *in vacuo*. The residue was purified by silica gel column chromatography (ethyl acetate), and the title compound (8.1mg, 0.040mmol, 64%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 1.49 (3H, t, J=7.1Hz), 4.51 (2H, q, J=7.1Hz), 7.61 (1H, dd, J=4.3, 8.2Hz), 8.34 (1H, dd, J=2.0, 8.2Hz), 8.91 (1H, d, J=2.2Hz), 9.25 (1H, dd, J=2.0, 4.2Hz), 9.69 (1H, d, J=2.4Hz).

Preparation Example K-1. 2-Methyl-benzoxazole-6-carboxylic acid methyl ester

To a solution of 4-amino-3-hydroxy-benzoic acid methyl ester (2085mg, 12.47mmol) in xylene (200mL) were added acetyl chloride (1.06mL, 14.96mmol), pyridinium *p*-toluenesulfonate (940mg, 3.74mmol) and triethylamine (2.09mL, 14.96mmol), and the solution was stirred for 8.5 hours under reflux. Ethyl acetate was added to the reaction mixture, which was then washed with water, dried over anhydrous magnesium sulfate, then, evaporated *in vacuo*, the resulting residue was purified by silica gel chromatography (hexane-ethyl acetate), and the title compound (1917mg, 10.02mmol, 80.4%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 2.69 (3H, s), 3.96 (3H, s), 7.68 (1H, d, J=8.4Hz), 8.05 (1H, dd, J=1.2, 8.4Hz), 8.17 (1H, d, J= 1.2Hz).

Preparation Example K-2. 2-Methyl-benzoxazole-6-carboxylic acid

To a solution of 2-methyl-benzoxazole-6-carboxylic acid methyl ester (301mg, 1.57mmol) in ethanol (10mL) was added an aqueous solution of 2N sodium hydroxide (10mL), and the mixture was stirred for 2 hours at room temperature. 2N Hydrochloric acid was added to the reaction mixture to adjust the pH to 4, and the solution was extracted with ethyl acetate. The organic layer was washed with brine,

dried over anhydrous sodium sulfate, then, evaporated *in vacuo*, and the title compound (270mg, 1.52mmol, 97%) was obtained. This was used in the next reaction without purification.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.64 (3H, s), 7.73 (1H, d, J=8.0Hz), 7.93(1H, dd, J=1.2, 8.0Hz), 8.15 (1H, d, J= 1.2Hz).

Preparation Example O-1. 2,3-Dihydro-1H-pyrrolo[2,3-b]pyridine-5-carbonitrile

1H-Pyrrolo[2,3-b]pyridine (1.0g, 8.46mmol) and 10% palladium-carbon (500mg) were dissolved in a mixture of formic acid (10mL) and triethylamine (10mL), and the solution was stirred at 70°C for 17 hours. To this reaction mixture was further added 10% palladium-carbon (270mg), and the mixture was stirred at 70°C for 2 hours. The reaction mixture was cooled to room temperature, then an aqueous solution of 5N sodium hydroxide was added thereto, the solution was extracted with ethyl acetate and tetrahydrofuran, and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (ethyl acetate : methanol = 10 : 1), and 2,3-dihydro-1H-pyrrolo[2,3-b]pyridine (614mg, 5.11mmol, 60%) was obtained as a pale yellow solid.

The resulting 2,3-dihydro-1H-pyrrolo[2,3-b]pyridine (614mg, 5.11mmol) and N-bromosuccinimide (1.09g, 6.13mmol) were dissolved in N,N-dimethylformamide (12mL), and the solution was stirred for 2.5 hours at room temperature. An aqueous solution of saturated sodium bicarbonate was added to the reaction solution at 0°C, which was then extracted with ethyl acetate, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by NH silica gel column chromatography (hexane :

ethyl acetate = 3 : 1), and 5-bromo-2,3-dihydro-1*H*-pyrrolo[2,3-*b*]pyridine (370mg, 1.86mmol, 36%) was obtained as a white solid.

The resulting 5-bromo-2,3-dihydro-1*H*-pyrrolo[2,3-*b*]pyridine (345mg, 1.73mmol), zinc cyanide (305mg, 2.60mmol) and tetrakis(triphenylphosphine)palladium(0) (200mg, 0.173mmol) were dissolved in dimethylsulfoxide (7mL), and the solution was stirred at 120°C for 4 hours under nitrogen atmosphere. The reaction solution was cooled to room temperature, water and ethyl acetate were added to the reaction solution, the organic layer was separated, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (ethyl acetate), and the title compound (167mg, 1.15mmol, 66%) was obtained as a light brown solid.

<sup>1</sup>H-NMR Spectrum (DMSO-*d*<sub>6</sub>) δ(ppm) : 3.01 (2H, t, J=8.6Hz), 3.58 (2H, t, J=8.6Hz), 7.46 (1H, s), 7.63 (1H, s), 8.10 (1H, s).

15

Preparation Example O-2. 2,3-Dihydro-1*H*-pyrrolo[2,3-*b*]pyridine-5-carboxylic acid

The title compound (259mg, quantitatively) was obtained as a white solid from 2,3-dihydro-1*H*-pyrrolo[2,3-*b*]pyridine-5-carbonitrile described in Preparation Example O-1 (167mg, 1.15mmol) according to an analogous method to Preparation Example T-6.

20

<sup>1</sup>H-NMR Spectrum (DMSO-*d*<sub>6</sub>) δ(ppm) : 3.00 (2H, t, J=8.6Hz), 3.56 (2H, t, J=8.6Hz), 7.25 (1H, s), 7.59 (1H, s), 8.30 (1H, s).

Preparation Example P-1. 6-Oxo-1,6-dihydro-pyridine-3-carboxylic acid ethyl ester

To a suspension of 6-hydroxy-nicotinic acid (5.00g, 35.9mmol) in ethanol

25

(60mL) was added 1N hydrochloric acid (20mL), which was then stirred at 110°C for 3 hours. The reaction solution was cooled to 0°C, then, an aqueous solution of saturated sodium bicarbonate was added, the solution was extracted with ethyl acetate and tetrahydrofuran, and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (ethyl acetate), and the title compound (3.90g, 23.3mmol, 65%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 1.27 (3H, t, J=7.1Hz), 4.23 (2H, q, J=7.1Hz), 6.36 (1H, d, J=9.7Hz), 7.79 (1H, dd, J=2.6, 9.7Hz), 8.03 (1H, d, J=2.6Hz).

10

Preparation Example P-2. 5-Iodo-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid ethyl ester

The title compound (2.82g, 9.62mmol, 80%) was obtained as a white solid from 6-oxo-1,6-dihydro-pyridine-3-carboxylic acid ethyl ester described in Preparation Example P-1 (2.00g, 12.0mmol) according to an analogous method to Preparation Example A+-16.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 1.28 (3H, t, J=7.1Hz), 4.23 (2H, q, J=7.1Hz), 8.09 (1H, s), 8.36 (1H, d, J=2.4Hz).

20 Preparation Example P-3. 6-Oxo-5-trimethylsilanylethynyl-1,6-dihydro-pyridine-3-carboxylic acid ethyl ester

5-Iodine-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid ethyl ester described in Preparation Example P-2 (1.00g, 3.41mmol), trimethylsilylacetylene (626μl, 4.43mmol), palladium(II) acetate (7.66mg, 34μmol), triphenylphosphine (17.9mg, 68μmol), copper(I) iodide (13mg, 68μmol) and butylamine (674μl, 6.82mmol) were

25



suspended in tetrahydrofuran (6mL), and the mixture was stirred at 40°C for 16 hours under nitrogen atmosphere. The reaction mixture was cooled to room temperature, then, water was added thereto, the solution was extracted with ethyl acetate, and the organic layer was washed with brine. The organic layer was dried over anhydrous magnesium sulfate, then, the solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 1 : 1), and the title compound (551mg, 2.09mmol, 61%) was obtained as a light brown solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 0.22 (9H, s), 1.27 (3H, t, J=7.1Hz), 4.23 (2H, q, J=7.1Hz), 7.91 (1H, d, J=2.4Hz), 8.07 (1H, d, J=2.4Hz).

10

Preparation Example P-4. Furo[2,3-b]pyridine-5-carboxylic acid ethyl ester

6-Oxo-5-trimethylsilanylethynyl-1,6-dihydro-pyridine-3-carboxylic acid ethyl ester described in Preparation Example P-3 (545mg, 2.07mmol) and copper(I) iodide (5.9mg, 31μmol) were suspended in ethanol (7mL) and triethylamine (3mL), and the mixture was stirred at 75°C for 20 hours. The reaction mixture was cooled to room temperature, then, potassium carbonate (572mg, 4.14mmol) was added to the reaction mixture, which was further stirred at 75°C for 5 hours. The reaction mixture was cooled to 0°C, then, water was added, the precipitated solid was filtered, and the title compound (303mg) was obtained as a brown solid. In addition, ethyl acetate was added to the mother liquor for extraction, the organic layer was washed with brine, then, dried over anhydrous magnesium sulfate, the solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 1 : 1), and the title compound (33mg, 0.17mmol) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 1.36 (3H, t, J=7.1Hz), 4.38 (2H, q, J=7.1Hz),

25

7.16 (1H, d, J=2.4Hz), 8.25 (1H, d, J=2.2Hz), 8.69 (1H, d, J=1.8Hz), 8.87 (1H, d, J=2.0Hz).

Preparation Example Q+-1. Imidazo[1,2-a]pyridine-6-carboxylic acid

5 (5-(3-fluorophenoxy)thiophene-2-ylmethyl) amide

To a solution of imidazo[1,2-a]pyridine-6-carboxylic acid (87mg, 0.54mmol) and C-(5-(3-fluorophenoxy)thiophen-2-yl)methylamine (120mg, 0.54mmol) in N,N-dimethylformamide (5mL) were added benzotriazol-1-yl-tris(dimethylamino)phosphonium hexafluorophosphate (240mg, 10 0.54mmol) and triethylamine(0.15mL, 1.08mmol), and the solution was stirred for 40 minutes at 80°C. Water and ethyl acetate were added to the reaction mixture for extraction, and the organic layer was washed twice with water. Silica gel was added to the organic layer, solvent was evaporated *in vacuo* for adsorption, purification was carried out by silica gel column chromatography (hexane : ethyl acetate = 1 : 1, then 15 ethyl acetate), and the title compound (90mg, 0.25mmol, 45.4%) was obtained as a light brown oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.55 (2H, d, J=5.6Hz), 6.58 (1H, d, J=4.0Hz), 6.83 (1H, d, J=4.0Hz), 6.90-7.00 (3H, m), 7.40 (1H, ddd, J=8.0, 8.0, 8.0Hz), 7.57-7.66 (3H, m), 8.04 (1H, s), 9.12 (1H, d, J=0.8Hz), 9.20 (1H, t, J=5.6Hz).

20

Preparation Example R-1. 2,6-Diamino-5-iodo-nicotinic acid ethyl ester

To a solution of 2,6-diamino-nicotinic acid ethyl ester described in Preparation Example A-14 (1.4g, 7.7mmol) in N,N-dimethylformamide (15mL) was added N-iodosuccinimide (2.09g, 9.3mmol), and the solution was stirred for 1 hour at room 25 temperature. The reaction mixture was poured into an aqueous solution of

saturated sodium thiosulfate pentahydrate, and the solution was extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, and then evaporated. The resulting residue was purified by silica gel column chromatography (ethyl acetate : hexane = 1 : 1) and the title  
5 compound (0.84g, 2.7mmol, 35.5%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 1.36 (3H, t, J=7.2Hz), 4.27 (2H, q, J=7.2Hz), 5.10 (2H, brs), 8.23 (1H, s).

10 Preparation Example R-2. 6-Amino-1*H*-pyrrolo[2,3-*b*]pyridine-5-carboxylic acid ethyl ester

Catechol borane (2.7mL, 1M tetrahydrofuran solution, 2.7mmol) was added dropwise to ethoxyacetylene (0.7mL, 40% hexane solution, 2.83mmol) on an ice bath, and the mixture was stirred for 1 hour at room temperature. The mixture was further heated at 70°C and stirred for 2 hours, and allowed to room temperature. A solution  
15 of 2,6-diamino-5-iodo-nicotinic acid ethyl ester described in Preparation Example R-1 (415mg, 1.35mmol) in tetrahydrofuran (5.5mL), tetrakis(triphenylphosphine)palladium (0) (48mg, 0.042mmol) and sodium hydroxide (160mg, 4mmol, powder) were added thereto, followed by stirring for 7 hours 30 minutes under reflux. The reaction mixture was allowed to room temperature, 2N hydrochloric acid (4.7mL, 9.4mmol)  
20 was added thereto, followed by stirring for 60 hours at room temperature. After the reaction was completed, the reaction mixture was evaporated, and extracted using diethyl ether. The aqueous layer was fractionated, neutralized on an ice bath with an aqueous solution of 5N sodium hydroxide, then, was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, then evaporated,  
25 the resulting residue was purified by silica gel column chromatography (ethyl

acetate : hexane = 2 : 1), and the title compound (97mg, 0.47mmol, 35%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 1.41(3H, t, J=7.2Hz), 4.36(2H, q, J=7.2Hz), 6.28-6.42(3H, m), 6.99-7.02(1H, m), 8.49(1H, s), 9.19(1H, brs).

5

Preparation Example R-3. 2,3-Dihydro-1H-pyrrolo[2,3-b]pyridine

1H-Pyrrolo[2,3-b]pyridine (1.00g, 8.46mmol) and 10% palladium-carbon (500mg) were dissolved in a mixture of formic acid (10mL) and triethylamine (10mL), and the mixture was stirred at 70°C for 87 hours. To this reaction mixture was  
10 further added 10% palladium-carbon (400mg), and the solution was stirred for 9.5 hours at 70°C. The reaction mixture was cooled to room temperature, an aqueous solution of 5N sodium hydroxide was added thereto, the solution was extracted with ethyl acetate and tetrahydrofuran, and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified  
15 by silica gel column chromatography (ethyl acetate : methanol = 10 : 1), and the title compound (219mg, 1.82mmol, 22%) was obtained as a pale yellow solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.94 (2H, t, J=8.4Hz), 3.43 (2H, t, J=8.4Hz), 6.27 (1H, s), 6.39 (1H, dd, J=5.3, 7.0Hz), 7.22 (1H, d, J=7.0Hz), 7.66 (1H, d, J=4.9Hz).

20

Preparation Example R-4. 5-Bromo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine

2,3-Dihydro-1H-pyrrolo[2,3-b]pyridine described in Preparation Example R-3 (15mg, 0.13mmol) and N-bromosuccinimide (24mg, 0.14mmol) were dissolved in N,N-dimethylformamide (0.5mL), and the solution was stirred for 15 hours at room  
25 temperature. An aqueous solution of saturated sodium bicarbonate was added to

the reaction solution at 0°C, which was then extracted with ethyl acetate, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 5 : 1), and the title compound (12mg, 60μmol, 48%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.98 (2H, t, J=8.8Hz), 3.48 (2H, t, J=8.8Hz), 6.60 (1H, s), 7.37 (1H, d, J=1.1Hz), 7.71 (1H, d, J=2.4Hz).

Preparation Example R-5. 5-Bromo-1H-pyrrolo[2,3-b]pyridine

5-Bromo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine (600mg, 3.01mmol) and 2,3-dichloro-5,6-dicyano-1,4-benzoquinone described in Preparation Example R-4 (753mg, 3.31mmol) was dissolved in toluene (15mL), and the solution was refluxed for 40 minutes under nitrogen atmosphere. The reaction solution was cooled to room temperature, water and ethyl acetate were added thereto, the organic layer was partitioned, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 5 : 1), and the title compound (260mg, 1.32mmol, 44%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 6.40-6.48 (1H, m), 7.50-7.60 (1H, m), 8.20 (1H, s), 8.30 (1H, s), 11.9 (1H, s).

Preparation Example R-6. 1H-Pyrrolo[2,3-b]pyridine-5-carbonitrile

5-Bromo-1H-pyrrolo[2,3-b]pyridine described in Preparation Example R-5 (90mg, 0.46mmol), zinc cyanide (80mg, 0.69mmol) and tetrakis(triphenylphosphine)palladium(0) (53mg, 46μmol) were dissolved in

N-methyl-2-pyrrolidinone (2mL), and the mixture was stirred for 4.5 hours at 110°C under nitrogen atmosphere. The reaction mixture was cooled to room temperature, water and ethyl acetate were added to the reaction mixture, the organic layer was partitioned, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 5 : 1), and the title compound (48mg, 0.34mmol, 73%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 6.55-6.68(1H, m), 7.65-7.78 (1H, m), 8.52 (1H, s), 8.60 (1H, s), 12.3 (1H, brs).

10

Preparation Example R-7. 1H-Pyrrolo[2,3-b]pyridine-5-carboxylic acid

The title compound (47mg, 0.29mmol, 88%) was obtained as a white solid from 1H-pyrrolo[2,3-b]pyridine-5-carbonitrile described in Preparation Example R-6 (47mg, 0.33mmol) according to an analogous method to Preparation Example T-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 6.57-6.63 (1H, m), 7.55-7.62 (1H, m), 8.51 (1H, s), 8.79 (1H, s), 12.0 (1H, s), 12.7 (1H, brs).

Preparation Example S-1. 3-Amino-2-bromopyridine

2-Bromo-3-nitropyridine (3g, 15mmol) was dissolved in a mixture solution of tetrahydrofuran (15mL) and water (5mL), then iron powder (1g, 18mmol) and ammonium chloride (2g, 37mmol) were added thereto, followed by stirring at from 60°C to 70°C for 5 hours. After the reaction was completed, the reaction mixture was filtered through Celite pad, brine was added, and the solution was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, then evaporated, the resulting residue was purified by silica gel column chromatography

(ethyl acetate : hexane = 1 : 4), and the title compound (2.6g, 15mmol, quantitatively) was obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 5.47 (2H, brs), 7.07-7.09 (2H, m), 7.54 (1H, dd, J=2.0, 3.6Hz).

5

Preparation Example S-2. (2-Bromo-pyridin-3-yl)carbamic acid ethyl ester

3-Amino-2-bromopyridine described in Preparation Example S-1 (1.4g, 8.1mmol) was dissolved in pyridine (10mL), ethyl chloroformate (0.93mL, 9.7mmol) was added dropwise on an ice bath, and the solution was stirred for 2 hours at room temperature. After the reaction was completed, the reaction solution was poured into brine, and the solution was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, then evaporated, the resulting residue was purified by silica gel column chromatography (ethyl acetate : hexane = 1 : 4), and the title compound (0.56g, 2.3mmol, 28%) was obtained.

10

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 1.22 (3H, t, J=7.2Hz), 4.12 (2H, q, J=7.2Hz), 7.43 (1H, dd, J=4.8, 8.0Hz), 7.92 (1H, dd, J=1.6, 8.0Hz), 8.17 (1H, dd, J=1.6, 4.8Hz), 9.10 (1H, brs).

15

Preparation Example S-3. (2-Trimethylsilanylethynyl-pyridin-3-yl)carbamic acid ethyl ester

20

A mixture of (2-bromo-pyridin-3-yl)carbamic acid ethyl ester described in Preparation Example S-2 (395mg, 1.6mmol), dichlorobis(triphenylphosphine)palladium(II) (20mg, 0.028mmol), triethylamine (0.25mL, 1.8mmol), copper(I) iodide (10mg, 0.05mmol) and trimethylsilylacetylene (0.131mL, 2.4mmol) was placed in a sealed tube, and heated at 100°C for 4 hours.

25

After the reaction was completed, the reaction mixture was poured into water, which was then extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, then evaporated, the resulting residue was purified by silica gel column chromatography (ethyl acetate : hexane = 1 : 2), and the title compound  
5 (0.42g, 1.6mmol, quantitatively) was obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 0.24 (9H, s), 1.21 (3H, t, J=7.2Hz), 4.12 (2H, q, J=7.2Hz), 7.38 (1H, dd, J=4.8, 8.4Hz), 7.88-7.96 (1H, m), 8.29 (1H, dd, J=1.6, 4.8Hz), 8.82 (1H, brs).

10 Preparation Example S-4. 1H-Pyrrolo[3, 2-b]pyridine

(2-Trimethylsilanylethynyl-pyridin-3-yl)carbamic acid ethyl ester described in Preparation Example S-3 (0.42g, 1.6mmol) was dissolved in ethanol (8mL), sodium ethoxide (204mg, 3mmol) was added thereto, followed by stirring for 1 hour under reflux. After the reaction was completed, the reaction mixture was poured into brine,  
15 and the solution was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, then evaporated, the resulting solid was washed with solvent (diethylether : hexane = 1 : 2), and the title compound (0.12g, 1mmol, 63.5%) was obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 6.50-6.54 (1H, m), 7.06 (1H, dd, J=4.8, 8.4Hz), 7.58-7.62 (1H, m), 7.72-7.76 (1H, m), 8.26-8.30 (1H, m), 11.2 (1H, brs).  
20

Preparation Example T-1. 3-Dichloromethyl-2-nitro-thiophene

To a solution of potassium *tert*-butoxide (23.0mL, 1.0M tetrahydrofuran solution, 23.2mmol) in N,N-dimethyl formamide (20mL) were added a mixture  
25 solution of 2-nitro-thiophene (1.00g, 7.74mmol) in chloroform (682μl, 8.51mmol) and



N,N-dimethylformamide (2mL) dropwise at -78°C, the solution was stirred for 5 minutes, and then, methanol and acetic acid were added at 0°C. Brine was added to the reaction solution, which was then extracted with ethyl acetate. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 20 : 1), and the title compound (1.54g, 7.26mmol, 94%) was obtained as a light brown solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 7.54 (1H, d, J=5.7Hz), 7.57 (1H, d, J=5.7Hz), 7.64 (1H, s).

10 Preparation Example T-2. 2-nitro-thiophene-3-carbaldehyde

3-Dichloromethyl-2-nitro-thiophene described in Preparation Example T-1 (1.54g, 7.26mmol) was dissolved in formic acid (10mL), and the solution was refluxed for 24 hours under nitrogen atmosphere. An aqueous solution of 5N sodium hydroxide was added to the reaction solution at 0°C, which was then extracted with ethyl acetate, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 10 : 1), and the title compound (472mg, 3.00mmol, 41%) was obtained as a light brown solid.

15 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 7.49 (1H, d, J=5.5Hz), 7.54 (1H, d, J=5.7Hz),  
20 10.62 (1H, s).

Preparation Example T-3. 2-(2-Nitro-thiophen-3-yl)-[1,3]dioxolane

2-Nitro-thiophene-3-carbaldehyde described in Preparation Example T-2 (367mg, 2.33mmol), ethane-1,2-diol (651μl, 11.7mmol) and toluene-4-sulfonic acid monohydrate (40mg, 0.233mmol) were dissolved in toluene (8mL), and the solution

was stirred for 2.5 hours under reflux. An aqueous solution of saturated sodium bicarbonate was added to the reaction solution at 0°C, which was then extracted with ethyl acetate, the organic layer was washed with brine, and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 5 : 1), and the title compound (304mg, 1.51mmol, 65%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.07-4.15 (4H, m), 6.51 (1H, s), 7.25 (1H, d, J=5.5Hz), 7.45 (1H, d, J=5.5Hz).

10

#### Preparation Example T-4. 2-Amino-thiophene-3-carbaldehyde

2-(2-Nitro-thiophen-3-yl)-[1,3]dioxolane described in Preparation Example T-3 (150mg, 0.746mmol), iron powder (208mg, 3.73mmol) and ammonium chloride (80mg, 1.49mmol) were suspended in a mixture solvent of ethanol (3mL) and water (0.75mL), and the mixture was stirred at 90°C for 5 hours. The reaction mixture was cooled to room temperature, then, filtered through Celite pad. The filtrate was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 5 : 1), and the title compound (38mg, 0.30mmol, 40%) was obtained as a red oily substance.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 6.19 (1H, d, J=5.7Hz), 6.67 (2H, brs), 6.90 (1H, d, J=5.7Hz), 9.69 (1H, s).

20

#### Preparation Example T-5. 6-Amino-thieno[2,3-b]pyridine-5-carbonitrile

2-Amino-thiophene-3-carbaldehyde described in Preparation Example T-4 (38mg, 0.30mmol) and malononitrile (20mg, 0.30mmol) were dissolved in ethanol

25

(1mL) to which piperidine (several drops) had been added, and the solution was stirred for 1 hour under reflux. The reaction solution was cooled to room temperature, and the solvent was evaporated *in vacuo*. The residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 10 : 1), and the title  
5 compound (50mg, 0.29mmol, 96%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 7.00 (2H, s), 7.18 (1H, d, J=6.0Hz), 7.42 (1H, d, J=6.0Hz), 8.40 (1H, s).

Preparation Example T-6. 6-Amino-thieno[2,3-b]pyridine-5-carboxylic acid

10 6-Amino-thieno[2,3-b]pyridine-5-carbonitrile described in Preparation Example T-5 (104mg, 0.594mmol) was dissolved in a mixture solution of water (1.5mL) and sulfuric acid (1.5mL), and the solution was stirred for 3 hours under reflux. An aqueous solution of 5N sodium hydroxide was added to the reaction solution at 0°C, to neutralize the solution. The precipitated solid was filtered, and  
15 title compound (65mg, 0.33mmol, 56%) was obtained as a pale yellow solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 7.19 (1H, d, J=5.9Hz), 7.25 (1H, d, J=6.0Hz), 8.48 (1H, s).

Preparation Example T-7. 6-Amino-thieno[2,3-b]pyridine-5-carboxylic acid methyl  
20 ester

6-Amino-thieno[2,3-b]pyridine-5-carboxylic acid (44mg, 0.23mmol) was dissolved in a mixture solution of methanol (1mL) and sulfuric acid (0.5mL), and the solution was stirred for 24 hours under reflux. An aqueous solution of saturated sodium bicarbonate was added to the reaction solution at 0°C, which was then  
25 extracted with ethyl acetate, the organic layer was washed with brine and dried over

anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 2 : 1), and the title compound (34mg, 0.16mmol, 72%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.93 (3H, s), 7.09 (1H, d, J=6.0Hz), 7.11 (1H, d, J=6.0Hz), 8.54 (1H, s).

Preparation Example T-8. 6-Oxo-6,7-dihydro-thieno[2,3-b]pyridine-5-carboxylic acid methyl ester

6-Amino-thieno[2,3-b]pyridine-5-carboxylic acid methyl ester described in Preparation Example T-7 (10mg, 48μmol) and sodium nitrite (10mg, 144μmol) were dissolved in phosphinic acid (0.5mL), and the solution was stirred at 0°C for 1 hour. An aqueous solution of saturated sodium bicarbonate was added to the reaction solution at 0°C, which was then extracted with ethyl acetate, and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 1 : 1), and the title compound (10mg, 48μmol, quantitatively) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.03 (3H, s), 7.21 (1H, d, J=5.9Hz), 7.34 (1H, d, J=6.0Hz), 8.61 (1H, s), 11.4 (1H, s).

Preparation Example T-9. 6-trifluoromethanesulfonyloxy-thieno[2,3-b]pyridine-5-carboxylic acid methyl ester

6-Oxo-6,7-dihydro-thieno[2,3-b]pyridine-5-carboxylic acid methyl ester described in Preparation Example T-8 (9mg, 43μmol), N-phenyltrifluoromethanesulfonimide (23mg, 65μmol) and

dimethyl-pyridin-4-yl-amine (catalytic amount) were dissolved in dichloromethane (0.5mL), and the solution was stirred for 18.5 hours at room temperature. Water was added to the reaction solution, which was then extracted with ethyl acetate, the organic layer was washed with brine and dried over anhydrous magnesium sulfate.

5 The solvent was evaporated *in vacuo*, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 3 : 1), and the title compound (10mg, 29μmol, 68%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.03 (3H, s), 7.43 (1H, d, J=5.9Hz), 7.73 (1H, d, J=5.9Hz), 8.87 (1H, s).

10

Preparation Example T-10. Thieno[2,3-b]pyridine-5-carboxylic acid methyl ester

6-Trifluoromethanesulfonyloxy-thieno[2,3-b]pyridine-5-carboxylic acid methyl ester described in Preparation Example T-9 (10mg, 29μmol), tetrakis(triphenylphosphine)palladium(0) (3.4mg, 2.9μmol), formic acid (1.7μl, 15 44μmol) and N,N-diisopropylethylamine (15μl, 87μmol) were dissolved in 1-methyl-2-pyrrolidone (0.5mL), and the mixture was stirred for 1.5 hours at 100°C. The reaction mixture was cooled to room temperature, water and ethyl acetate were added, the organic layer was partitioned, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in* 20 *vacuo*, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 5 : 1), and the title compound (7mg, quantitatively) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.99 (3H, s), 7.36 (1H, d, J=6.4Hz), 7.62 (1H, d, J=6.0Hz), 8.70 (1H, d, J=1.6Hz), 9.17 (1H, d, J=2.0Hz).

25

Preparation Example U-1. Thiophen-3-yl-carbamic acid *tert*-butyl ester

Thiophene-3-carboxylic acid (2.50g, 19.5mmol), diphenylphosphoryl azide (4.62mL, 21.5mmol), triethylamine (3.26mL, 23.4mmol) were dissolved in *tert*-butanol (50mL), and the solution was stirred for 3.5 hours under reflux. Water was added to the reaction solution at 0°C, which was then extracted with ethyl acetate, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 10 : 1), and the title compound (3.33g, 16.7mmol, 86%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 1.46 (9H, s), 6.97 (1H, d, J=5.2Hz), 7.16 (1H, s), 7.38 (1H, m), 9.61 (1H, s).

Preparation Example U-2. (2-Formyl-thiophen-3-yl)-carbamic acid *tert*-butyl ester

Thiophen-3-yl-carbamic acid *tert*-butyl ester described in Preparation Example U-1 (1.00g, 5.02mmol) was dissolved in tetrahydrofuran (20mL), to which *n*-butyl lithium (2.47M hexane solution, 4.47mL, 11.0mmol) was added at -78°C, and the mixture was stirred at -78°C for 1 hour. N,N-dimethylformamide (466μL, 6.02mmol) was added to the reaction mixture at -78°C, and the solution was stirred for 1 hour at room temperature. Water was added to the reaction solution, which was then extracted with ethyl acetate, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 10 : 1), and the title compound (1.14g, quantitatively) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 1.50 (9H, s), 7.60 (1H, d, J=5.3Hz), 8.02 (1H,

d, J=5.3Hz), 9.94 (1H, s), 10.1 (1H, s).

Preparation Example U-3. 5-Amino-thieno[3,2-b]pyridine-6-carbonitrile

(2-Formyl-thiophen-3-yl)-carbamic acid *tert*-butyl ester described in  
5 Preparation Example U-2 (500mg, 2.20mmol) and malononitrile (153mg, 2.31mmol)  
were dissolved in a solution of ethanol (10mL) to which piperidine (catalytic amount)  
had been added, and the mixture was stirred for 1 hour under reflux. The reaction  
mixture was cooled to room temperature, the precipitated solid was filtered, and title  
compound (215mg, 1.23mmol, 56%) was obtained as a white solid.

10 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 6.76 (2H, s), 7.22 (1H, dd, J=0.73, 5.5Hz),  
8.22 (1H, d, J=5.5Hz), 8.64 (1H, s).

Preparation Example U-4. 5-Amino-thieno[3,2-b]pyridine-6-carboxylic acid

The title compound (200mg) was obtained as a white solid from  
15 5-amino-thieno[3,2-b]pyridine-6-carbonitrile described in Preparation Example U-3  
(208mg, 1.19mmol) according to an analogous method to Preparation Example T-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 7.27 (1H, dd, J=0.73, 5.5Hz), 8.28 (1H, d,  
J=5.5Hz), 8.92 (1H, s).

20 Preparation Example U+-1. 5-Oxo-4,5-dihydro-thieno[3,2-b]pyridine-6-carboxylic acid  
(5-phenoxy-thiophen-2-ylmethyl)-amide

The title compound (17mg, 44μmol, 46%) was obtained as a white solid from  
5-amino-thieno[3,2-b]pyridine-6-carboxylic acid  
(5-phenoxy-thiophen-2-ylmethyl)-amide described in Preparation Example U-4 (37mg,  
25 97μmol) according to an analogous method to Preparation Example T-8.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.59 (2H, d, J=5.9Hz), 6.49 (1H, d, J=3.8Hz), 6.79 (1H, d, J=3.7Hz), 7.07-7.15 (4H, m), 7.37 (2H, t, J=7.7Hz), 8.15 (1H, d, J=5.5Hz), 8.94 (1H, s), 10.3 (1H, m), 13.0 (1H, s).

5     Preparation                      Example                      U+-2.                      Trifluoromethanesulfonic                      acid  
6-((5-phenoxy-thiophen-2-ylmethyl)-carbamoyl)-thieno[3,2-b]pyridin-5-yl ester

The title compound (11mg, 21μmol, 68%) was obtained as a white solid from 5-oxo-4,5-dihydro-thieno[3,2-b]pyridine-6-carboxylic acid (5-phenoxy-thiophen-2-ylmethyl)-amide described in Preparation Example U+-1 (12mg, 31μmol) according to an analogous method to Preparation Example T-9 (with the proviso that N,N-dimethylformamide was used instead of dichloromethane).

10     <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.53 (2H, d, J=5.5Hz), 6.53 (1H, d, J=3.8Hz), 6.83 (1H, d, J=4.4Hz), 7.09 (2H, d, J=8.6Hz), 7.13 (1H, t, J=7.7Hz), 7.37 (2H, t, J=7.7Hz), 7.65 (1H, d, J=5.5Hz), 8.50 (1H, d, J=5.7Hz), 8.97 (1H, s), 9.39-9.44(1H, 15     m).

Preparation Example W-1. Dithiocarbonic acid O-ethyl ester S-quinolin-6-yl ester

To a solution of quinolin-6-yl amine (2.88g, 20mmol) and HBF<sub>4</sub> aqueous solution (48% W/W, 11mL) in tetrahydrofuran (100mL) was added 20     3-methyl-1-nitrosooxy-butane (10.7mL, 60mmol) dropwise under nitrogen atmosphere at from -10°C to -15°C, then, the mixture was stirred for 1 hour. At from -10°C to -15°C, diethyl ether (200mL) was added dropwise, the precipitated solid was collected by filtration, washed with diethyl ether, dried, and quinolin-6-diazonium tetrafluoroborate (6.85g) was obtained as a yellow-red solid.

25             Next, to a solution of potassium O-ethyl dithiocarbonate (802mg, 5mmol),



water (20mL) and diethyl ether (30mL) was added quinolin-6-diazonium tetrafluoroborate (665mg, 2mmol) dropwise on an ice bath, then, the solution was stirred for 18 hours. To this reaction mixture was added water (100mL), then the solution was extracted with diethyl ether (50mL), and washed with brine (150mL).

5 The organic layer was dried over anhydrous magnesium sulfate, filtered, then, evaporation *in vacuo* was carried out, and a residue of red oil (0.462g) was obtained. This residue in the amount of 0.2g was purified by thin layer silica gel chromatography (hexane : ethyl acetate = 1 : 1), and the title compound (50mg, 0.2mmol) was obtained as a pale yellow oil.

10 <sup>1</sup>H-NMR Spectrum (Acetone-d<sub>6</sub>) δ(ppm) :1.31(3H, t, J=7.2 Hz), 4.66(2H, q, J=7.2 Hz), 7.60(1H, dd, J=8.4, 4.4 Hz), 7.86(1H, m), 8.13(1H, d, J=8.8 Hz), 8.18(1H, m), 8.41(1H, m), 9.01(1H, m).

MS m/e (ESI) 360(MH<sup>+</sup>)

15 Preparation Example X-1. 4-Chloro-quinazoline

Phosphorus oxychloride (64mL, 687mmol) and phosphorus pentachloride (14.89g, 71.50mmol) were added to 4-hydroxyquinazoline (7.94g, 52.3mmol), and the solution was stirred for 20 minutes under reflux. The reaction mixture was evaporated *in vacuo*, and the residue was dissolved in chloroform. The chloroform

20 solution was poured on an ice bath, concentrated aqueous ammonia was added to adjust the pH to 10, then, the solution was partitioned. The aqueous layer was extracted with chloroform, and the combined chloroform layers were washed with water. The organic layer was dried over anhydrous magnesium sulfate, then, evaporated *in vacuo*, and the title compound (8.03g, 48.8mmol, 93%) was obtained.

25 This was used in the next reaction without purification.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 7.64(1H, dd, J=7.2, 8.0Hz), 7.81(1H, d, J=8.0Hz), 7.93(1H, ddd, J=0.8, 7.2, 8.0Hz), 8.16(1H, dd, J=0.8, 8.0Hz), 8.86 (1H, brs).

5 Preparation Example Z-1. Quinoline-6-carbaldehyde

*n*-Butyl lithium (19.2mL; 1.5M hexane solution) was added dropwise to a solution of 6-bromoquinoline (5g, 24.0mmol) in diethyl ether (20mL) that had been cooled to -70°C under nitrogen atmosphere, a solution of N,N-dimethylformamide (3.7mL, 48.0mmol) in diethyl ether was further added, and the solution was stirred at  
10 -70°C for 2 hours. The reaction solution was allowed to room temperature, then, an aqueous solution of saturated ammonium chloride was added, the solution was extracted with ethyl acetate and dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by silica gel column chromatography (hexane : ethyl acetate), and the title compound (320mg, 8.5%) was  
15 obtained as a yellow oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 7.52-7.56(1H, m), 8.21-8.22(2H, m), 8.33-8.36(1H, m), 8.37(1H, s), 9.06-9.07(1H, m), 10.2(1H, s).

Preparation Example Z-2. 4-Bromo-1-butyl-2-methylbenzene

20 Aluminum chloride (38.9g, 0.292mol) was portionwise added to a solution of 3-bromotoluene (25g, 0.146mol) and *n*-butyryl chloride (22.9mL, 0.129mol) in carbon disulfide (100mL) that had been cooled with an ice water, and the solution was stirred for 18 hours while gradually allowing to room temperature. The reaction solution was poured into a mixture solution of an ice water (120mL) and concentrated  
25 hydrochloric acid (10mL), and the solution was stirred for 10 minutes. Then, the

solution was extracted with hexane, washed with water and an aqueous solution of saturated sodium bicarbonate and dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by silica gel column chromatography (hexane : ethyl acetate), and

5 1-(4-bromo-2-methyl-1-phenyl)-butane-1-one (7.77g, 22%) was obtained as a pale yellow oil. To a solution of the resulting compound in diethyleneglycol (40mL), were added hydrazine monohydrate (4.52g, 90.2mmol) and potassium hydroxide (4.16g, 74.1mmol), then the solution was stirred at 80°C for 2 hours, and further stirred at 160°C overnight. The reaction solution was allowed to room temperature, hexane

10 and water were added for partitioning, the organic layer was washed with water, and then, dried over anhydrous magnesium sulfate. The solvent was evaporated, then, the residue was purified by silica gel column chromatography (hexane : ethyl acetate), and the title compound (5.01g, 69%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.90(3H, t, J=2Hz), 1.32-1.36(2H, m),

15 1.51-1.55(2H, m), 2.27(3H, s), 2.52(2H, t, J=7.6Hz), 6.98(1H, d, J=8.0Hz), 7.23(1H, dd, J=2.0Hz, 8.0Hz), 7.26-7.27(1H, m).

#### Preparation Example Z+-1. (4-Butyl-3-methyl-phenyl)-quinolin-6-yl-metanol

Magnesium (83mg, 3.40mmol), 4-bromo-1-butyl-2-methylbenzene (722mg, 3.18mmol) and, as an initiator, catalytic amount of 1,2-dibromoethane were added to tetrahydrofuran (2.5mL) under nitrogen atmosphere, and the mixture was stirred for

20 10 minutes under reflux. This mixture was cooled to 0°C, a solution of quinoline-6-carbaldehyde (100mg, 6.36mmol) in tetrahydrofuran (12mL) was added thereto, followed by stirring for 1 hour at this temperature. An aqueous solution of

25 saturated ammonium chloride was added to the reaction mixture, which was

extracted with chloroform and dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by silica gel column chromatography (hexane : ethyl acetate), and the title compound (152mg, 78%) was obtained as a yellow solid.

5 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.93(3H, t, J=7.2Hz), 1.34-1.43(2H, m), 1.49-1.57(2H, m), 1.64(1H, brs), 2.28(3H, s), 2.57(2H, t, J=8.0Hz), 5.97(1H, brs), 7.10-7.17(3H, m), 7.38-7.42(1H, m), 7.66-7.68(1H, m), 7.93(1H, s), 8.03-8.05(1H, m), 8.16-8.18(1H, m), 8.87-8.88(1H, m).

#### 10 Preparation Example Z+-2. Quinoline-6-carboxylic acid (4-benzyloxy phenyl)-amide

To a solution of 6-quinolinecarboxylic acid (500mg, 2.89mmol) and 4-benzyloxyphenylamine (681mg, 2.89mmol) in dichloromethane (25mL) were added benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (1.53g, 3.47mmol) and triethylamine (0.96mL, 6.94mmol), and the solution was stirred  
15 overnight at room temperature. The solvent was evaporated, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate), and the title compound (194mg, 19%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 5.11(2H, s), 7.00-7.02(2H, m), 7.32-7.45(5H, m), 7.48-7.51(1H, m), 7.57-7.59(2H, m), 7.86(1H, brs), 8.11-8.14(1H, m), 8.19-8.21(1H,  
20 m), 8.26-8.28(1H, m), 8.38-8.39(1H, m), 9.00-9.01(1H, m).

#### Preparation Example Z+-3. 4-Benzyloxy-N-quinolin-6-yl-benzamide

To a solution of 6-aminoquinoline (500mg, 3.47mmol) and 4-benzyloxybenzoic acid (792mg, 3.47mmol) in dichloromethane (25mL) were added  
25 benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (1.84g,

4.16mmol) and triethylamine (0.58mL, 4.16mmol), and the solution was stirred overnight at room temperature. The solvent was evaporated, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate), and the title compound (218mg, 18%) was obtained as a white solid.

5 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 5.16(2H, s), 7.08-7.10(2H, m), 7.36-7.47(6H, m), 7.65-7.68(1H, m), 7.89-7.91(2H, m), 7.97(1H, brs), 8.08-8.10(1H, m), 8.15-8.17(1H, m), 8.49-8.50(1H, m), 8.84-8.86(1H, m).

#### Preparation Example 1. 4-Benzyloxybenzylamine

10 Potassium phthalimide (20g, 0.108mol) was added to a solution obtained by dissolving 4-benzyloxybenzyl chloride (25g, 0.107mol) in N,N-dimethylformamide (75mL), and the solution was stirred for 3 hours under reflux. The reaction solution was allowed to room temperature, then, ethyl acetate and water were added for partitioning, the organic layer was washed with water and dried over anhydrous  
15 magnesium sulfate. The solvent was evaporated, and 2-(4-benzyloxybenzyl)-isoindol-1,3-dione (37g, quantitatively) was obtained as a pale brown solid.

Next, to a solution of the resulting 2-(4-benzyloxybenzyl)-isoindol-1,3-dione (37g, 0.107mol) in ethanol (1L) was added hydrazine monohydrate (8.04g, 0.161mol),  
20 and the solution was stirred for 8 hours under reflux. The reaction solution was allowed to room temperature, then, water was added, and ethanol was evaporated *in vacuo*. Ethyl acetate and water were added to the residue for partitioning, the organic layer was washed with water, an aqueous solution of 2N sodium hydroxide and water, in this order, and dried over anhydrous magnesium sulfate. The solvent  
25 was evaporated, the residue was purified by NH silica gel column chromatography

(hexane : ethyl acetate = 2 : 1; hereinafter, the NH silica gel used was manufactured by Fuji Silysia), and the title compound (15g, 64%) was obtained as a white solid.

#### Preparation Example 2. 3-Benzyloxybenzylamine

5           To a solution of 3-benzyloxybenzyl alcohol (3.0g, 14.0mmol) in dichloromethane (30mL) was added methanesulfonyl chloride (1.39mL, 16.8mmol) and triethylamine (2.34mL, 16.8mmol) on an ice bath, and the solution was stirred overnight. The reaction solution was diluted with dichloromethane, washed with aqueous solution of 5% sodium bicarbonate, and then, dried over anhydrous  
10   magnesium sulfate. The solvent was evaporated, then, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 10 : 1), and 1-benzyloxy-3-chloromethyl-benzene (2.2g, 67%) was obtained as a colorless oil.

          Next, to a solution of iminodicarboxylic acid di-*tert*-butyl ester (2.12g, 8.76mmol) in N,N-dimethylformamide (13mL) was added sodium hydride (0.39g,  
15   9.86mmol, 60% in oil), the solution was stirred at 60°C for 6 hours, 1-benzyloxy-3-chloromethyl-benzene (1.0g, 4.30mmol) was added, and the solution was further stirred at 60°C for 4 hours. The reaction solution was allowed to room temperature, then, dichloromethane and water were added for partitioning, and the organic layer was dried over anhydrous magnesium sulfate. The solvent was  
20   evaporated, then, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 10 : 1), and (3-benzyloxybenzyl) iminodicarboxylic acid di-*tert*-butyl ester (691mg, 39%) was obtained as a pale yellow oil.

          Lastly, (3-benzyloxybenzyl) iminodicarboxylic acid di-*tert*-butyl ester (691mg, 1.67mmol) was cooled on an ice bath, trifluoroacetic acid (3mL) was added, and the  
25   solution was stirred for 30 minutes. An aqueous solution of saturated sodium

bicarbonate was added to the reaction solution, which was then extracted with ethyl acetate and dried over anhydrous magnesium sulfate. The solvent was evaporated, and the title compound (292mg, 82%) was obtained as a white waxy solid. This was used in the next reaction without further purification.

5

#### Preparation Example 3. 4-Phenoxybenzylamine

A solution of sodium borohydride (2.20g, 58.3mmol) and concentrated sulfuric acid in diethyl ether (1.6mL) was added to a solution of 4-phenoxybenzoic acid (5.0g, 23.3mmol) in tetrahydrofuran (20mL) that had been cooled on an ice bath, and the solution was stirred for 4 hours at room temperature. The reaction solution was cooled on an ice bath, methanol was added, then, the solution was allowed to room temperature and stirred for 30 minutes. This reaction solution was cooled again, ethyl acetate and an aqueous solution of 2N sodium hydroxide were added for partitioning, the organic layer was washed with 10% sodium chloride water, dried over anhydrous magnesium sulfate, and the solvent was evaporated to obtain 4-phenoxybenzyl alcohol (4.66g, quantitatively) as a colorless solid. This 4-phenoxybenzyl alcohol was used to carry out an analogous reaction to Preparation Example 2, and the title compound (886mg) was obtained as a pale brown solid.

#### 20 Preparation Example 4. 3-Phenoxybenzylamine

The title compound was obtained as a pale brown solid from 3-phenoxybenzyl alcohol according to a similar method to Preparation Example 2.

#### Preparation Example 5. C-Biphenyl-3-yl-methylamine

25 To a solution of 3-cyanophenylboronic acid (1.0g, 6.81mmol) and

bromobenzene (1.07g, 6.81mmol) in N,N-dimethylformamide (100mL) were added tetrakis(triphenylphosphine)palladium(0) (0.393g, 0.341mmol) and cesium carbonate (2.77g, 8.51mmol) under nitrogen atmosphere, and the mixture was stirred for 4 hours under reflux. The reaction mixture was allowed to room temperature, ethyl acetate and water were added for partitioning, the organic layer was washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 10 : 1), and biphenyl-3-carbonitrile (821mg, 67%) was obtained as a yellow solid.

Next, a solution of the resulting biphenyl-3-carbonitrile (821mg, 4.58mmol) in tetrahydrofuran (5mL) was added to a solution of lithium aluminum hydride (0.435g, 11.5mmol) in tetrahydrofuran (5mL) that had been cooled on an ice bath, and the solution was stirred for 6 hours at room temperature. The reaction solution was cooled on an ice bath, a mixture solution of methanol and water (9:1) was added thereto, an aqueous solution of saturated ammonium chloride was further added, filtration was carried out through Celite pad and insoluble matter was removed. The filtrate was partitioned, the organic layer was dried over anhydrous magnesium sulfate, and the title compound (527mg, 63%) was obtained as a brown oil. This was used in the next reaction without further purification.

#### 20 Preparation Example 6. 4-(3-Fluorobenzyloxy)-benzylamine

To a solution of 4-cyanophenol (3.0g, 25.2mmol) and 3-fluorobenzyl bromide (3.1mL, 25.2mmol) in N,N-dimethylformamide (30mL) was added potassium carbonate (8.71g, 63.0mmol), and the mixture was stirred for 1 hour at room temperature. Ethyl acetate and water were added to the reaction mixture, which was then partitioned, the organic layer was washed with water, then, dried over



anhydrous magnesium sulfate, and the solvent was evaporated to obtain 4-(3-fluorobenzyloxy)-benzonitrile (5.31g, 93%) as a colorless solid.

Next, to a solution of lithium aluminum hydride (1.25g, 133.0mmol) in tetrahydrofuran (15mL) was added a solution of 4-(3-fluorobenzyloxy)-benzonitrile (218mg, 0.615mmol) in tetrahydrofuran (12mL) on an ice bath, and the solution was stirred at room temperature for 19 hours. A mixture solvent of methanol and water (9:1) was added to the reaction solution, an aqueous solution of saturated ammonium chloride was further added, and the solution was stirred on an ice bath for 30 minutes. This solution was filtered through Celite pad, and insoluble matter was removed. The filtrate was partitioned, and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated, and the title compound (1.33g, 44%) was obtained as a yellow solid. This was used in the next reaction without further purification.

Preparation Example 7. C-(4-Phenoxy-pyridin-2-yl)-methylamine

To a solution of 4-phenoxy-pyridine (3.0g, 17.5mmol) in dichloromethane (500mL) was added 3-chloro-perbenzoic acid (5.18g, 21.0mmol) on an ice bath, and the solution was stirred for 22 hours. An aqueous solution of saturated sodium thiosulfate and an aqueous solution of saturated sodium bicarbonate were added to the reaction solution, the solution was stirred at room temperature for 10 minutes, then, the organic layer was separated, washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated, and 4-phenoxy-pyridine N-oxide (3.3g, quantitatively) was obtained as a pale yellow solid.

The resulting solid (3.3g, 17.6mmol) was dissolved in acetonitrile (18mL), trimethylsilyl cyanide (6.6mL, 52.8mmol) and triethylamine (4.9mL, 35.2mmol) were

added, and the solution was stirred for 5 hours under reflux. The solvent was evaporated, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 4 : 1), and 4-phenoxy-pyridine-2-carbonitrile (2.5g, 73%) was obtained as a pale yellow solid.

5        Next, to a solution of lithium aluminum hydride (725mg, 19.1mmol) in tetrahydrofuran (6.0mL) was portionwise added a solution of the resulting 4-phenoxy-pyridine-2-carbonitrile (1.5g, 7.65mmol) in tetrahydrofuran (3mL) on an ice bath, and the solution was stirred at room temperature for 15 hours. A mixture solvent of methanol and water (9:1) was added to the reaction solution, an aqueous  
10       solution of saturated ammonium chloride was further added, the solution was extracted with ethyl acetate, dried over anhydrous magnesium sulfate, then, the solvent was evaporated to obtain the title compound (730mg, 48%) as a pale brown oil. This was used in the next reaction without further purification.

15       Preparation Example 8. 3-(4-Fluorophenoxy)-benzylamine

The title compound (790mg, quantitatively) was obtained as a pale yellow solid from 3-(4-fluorophenoxy)benzyl bromide (944mg, 3.36mmol) according to a similar technique to Preparation Example 1.

20       Preparation Example 9. 3-(4-Methoxyphenoxy)benzylamine

To a solution of 3-(4-methoxyphenoxy)benzaldehyde (5.0g, 21.9mmol) in methanol (35mL) was added a solution of sodium borohydride (0.86g, 22.8mmol) in an aqueous solution of 2N sodium hydroxide (2.5mL), and the mixture was stirred at room temperature for 30 minutes. Water was added to the reaction mixture, which  
25       was neutralized with acetic acid, then, extracted with ethyl acetate, washed with brine,

then, dried over anhydrous magnesium sulfate, and the solvent was evaporated to obtain (3-(4-methoxy-phenoxy)-phenyl)-metanol (5.3g, quantitatively) as a colorless oil.

To a solution of the obtained (3-(4-methoxy-phenoxy)-phenyl)-metanol (2.0g, 8.73mmol) in dichloromethane (20mL) was added methanesulfonyl chloride (0.81mL, 10.5mmol) and triethylamine (1.46mL, 10.5mmol) on an ice bath, and the solution was stirred for 19 hours. The reaction solution was diluted with dichloromethane, washed with an aqueous solution of 5% sodium bicarbonate, dried over anhydrous magnesium sulfate, and the solvent was evaporated to obtain methanesulfonic acid 3-(4-methoxy-phenoxy)benzyl ester (2.4g, 89%) as a pale brown oil.

Next, the title compound (859mg, 89%) was obtained as a pale yellow solid from the resulting methanesulfonic acid 3-(4-methoxy-phenoxy)benzyl ester (2.4g, 7.78mmol) according to a similar technique to Preparation Example 1.

#### 15 Preparation Example 10. 3-(3-Trifluoromethyl-phenoxy)-benzylamine

The title compound (2.63g) was obtained as a brown oil from 3-(3-(trifluoromethyl)phenoxy)benzaldehyde (5.01g, 18.8mmol) according to an analogous method to Preparation Example 9.

#### 20 Preparation Example 11. 3-(3-Fluoro-phenoxy)-benzylamine

To a solution of 3-fluoro-phenol (500mg, 4.46mmol) and 3-fluoro-benzonitrile (540mg, 4.46mmol) in dimethylsulfoxide (1.0mL) was added potassium *tert*-butoxide (500mg, 4.46mmol), and the solution was stirred at 140°C for 3 hours. The reaction solution was allowed to room temperature, ethyl acetate and water were added for partitioning, the organic layer was washed with water and dried over anhydrous

magnesium sulfate. The solvent was evaporated, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 4 : 1), and 3-(3-fluoro-phenoxy)-benzonitrile (313mg, 33%) was obtained as a yellow solid.

Next, to a solution of lithium aluminum hydride (139mg, 3.68mmol) in tetrahydrofuran (3.0mL) was added a solution of the resulting 3-(3-fluoro-phenoxy)-benzonitrile (313mg, 1.47mmol) in tetrahydrofuran (1mL) on an ice bath, and the solution was stirred at room temperature for 18 hours. A mixture solvent of methanol and water (9:1) was added to the reaction solution, an aqueous solution of saturated ammonium chloride was further added thereto, followed by stirring at room temperature for 10 minutes, then, extracted with ethyl acetate, dried over anhydrous magnesium sulfate, then, the solvent was evaporated to obtain the title compound (285mg, 89%) as a yellow oil.

#### Preparation Example 12. 4-(Furan-2-ylmethoxy)-benzylamine

To a solution of 4-cyanophenol (2.0g, 16.8mmol) in dichloromethane (20mL) were added triphenyl phosphine (6.6g, 25.2mmol), furfuryl alcohol (1.65g, 16.8mmol) and diethyl azodicarboxylate (3.97mL, 25.2mmol) on an ice bath, and the solution was stirred at room temperature for 16 hours. The reaction solution was directly purified by silica gel column chromatography (hexane : ethyl acetate), the obtained crudely purified product was further purified by NH silica gel column chromatography (hexane : ethyl acetate), 4-furan-2-ylmethoxy)-benzonitrile (106mg, 3%) was obtained as a pale yellow solid.

Next, to a solution of lithium aluminum hydride (50mg, 1.33mmol) in tetrahydrofuran (1.0mL) was added a solution of the resulting 4-furan-2-ylmethoxy)-benzonitrile (106mg, 0.532mmol) in tetrahydrofuran (1mL) on

an ice bath, and the solution was stirred at room temperature for 4 hours. A mixture solvent of methanol and water (9 : 1) was added to the reaction solution, an aqueous solution of saturated ammonium chloride was further added, the solution was extracted with ethyl acetate, dried over anhydrous magnesium sulfate, then, the solvent was evaporated to obtain the title compound (76mg, 70%) as a yellow solid.

Preparation Example 13. 4-(Thiophen-2-ylmethoxy)-benzylamine

To a solution of 2-thiophenemethanol (2.0g, 17.5mmol) in dichloromethane (20mL) were added methanesulfonyl chloride (1.63mL, 21.0mmol) and triethylamine (2.93mL, 21.0mmol) on an ice bath, and the solution was stirred for 13 hours. The reaction solution was diluted with dichloromethane, washed with an aqueous solution of 5% sodium bicarbonate, dried over anhydrous magnesium sulfate, and the solvent was evaporated to obtain methanesulfonic acid 2-thiophen-2-ylmethyl ester (2.4g) as a brown oil.

Next, to a solution of the resulting methanesulfonic acid 2-thiophen-2-ylmethyl ester (2.4g, 12.6mmol) and *p*-cyanophenol (1.50g, 12.6mmol) in *N,N*-dimethylformamide (25mL) was added potassium carbonate (4.35g, 32.5mmol), and the solution was stirred at room temperature for 13 hours. Ethyl acetate and water were added to the reaction solution, which was then separated, the organic layer was washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 4 : 1), and 4-(thiophen-2-ylmethoxy)-benzonitrile (1.88g) was obtained as a white solid.

In addition, to a solution of lithium aluminum hydride (220mg, 5.80mmol) in tetrahydrofuran (2.5mL) was added a solution of the resulting

4-(thiophene-2-ylmethoxy)-benzonitrile (500mg, 2.32mmol) in tetrahydrofuran (1mL) on an ice bath, and the solution was stirred at room temperature for 4 hours. A mixture solvent of methanol and water (9:1) was added to the reaction solution, an aqueous solution of saturated ammonium chloride was further added, the solution  
 5 was extracted with ethyl acetate, dried over anhydrous magnesium sulfate, then, the solvent was evaporated to obtain the title compound (415mg, 82%) as a colorless solid.

Preparation Example 14. 4-(Thiophen-3-ylmethyl)-benzylamine

10 The title compound (419mg) was obtained as a pale brown solid from 3-thiophenemethanol according to an analogous method to Preparation Example 13.

Preparation Example 15. 4-((S)-1-Phenyl-ethoxy)-benzylamine

To a solution of 4-bromobenzonitrile (500mg, 2.75mmol) and  
 15 S-(-)- $\alpha$ -phenylethylalcohol (403mg, 3.30mmol) in toluene (5mL) were added sodium hydride (220mg, 5.49mmol; 60% in oil), tris(dibenzylideneacetone)dipalladium(0) (38mg, 0.0413mmol) and 2,2-bis (di-*p*-tolylphosphino)-1,1-binaphthyl (67mg, 0.099mmol), and the solution was stirred at 70°C for 4 hours. The reaction solution was allowed to room temperature, ethyl acetate and water were added for partitioning,  
 20 the organic layer was washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by silica gel column chromatography (hexane : ethyl acetate), and 4-(1-phenyl-ethoxy)-benzonitrile (159mg, 26%) was obtained as a colorless oil.

Next, to a solution of lithium aluminum hydride (68mg, 1.78mmol) in  
 25 tetrahydrofuran (5.0mL) was added a solution of the resulting

4-(1-phenyl-ethoxy)-benzonitrile (159mg, 0.712mmol) in tetrahydrofuran (1mL) on an ice bath, which was stirred under reflux for 2 hours. The reaction solution was allowed to room temperature, a mixture solvent of methanol and water (9:1) was added, an aqueous solution of saturated ammonium chloride was further added, then, ethyl acetate and water were added for partitioning, the organic layer was dried over anhydrous magnesium sulfate, then, the solvent was evaporated to obtain the title compound (172mg, quantitatively) as a yellow oily substance.

Preparation Example 16. C-(6-Phenoxy-pyridin-2-yl)-methylamine

To a solution of 2,6-dibromopyridine (20g, 84.4mmol) and phenol (7.94g, 84.4mmol) in dimethylsulfoxide (200mL) was added potassium *tert*-butoxide (9.47g, 84.4mmol), and the solution was stirred at 160°C for 7 hours. The reaction solution was allowed to room temperature, ethyl acetate and water were added for partitioning, the organic layer was washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 10 : 1), and 2-bromo-6-phenoxy-pyridine (19.6g, 93%) was obtained as a yellow solid.

Next, to a solution of the resulting 2-bromo-6-phenoxy-pyridine (1.0g, 4.0mmol) in N,N-dimethylformamide (30mL) were added zinc cyanide (940mg, 8.0mmol) and tetrakis(triphenylphosphine)palladium(0) (924mg, 0.8mmol) under nitrogen atmosphere, and the mixture was stirred at 100°C for 1 hour. The reaction mixture was allowed to room temperature, ethyl acetate and water were added for partitioning, the organic layer was washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 10 : 1), and

6-phenoxy-pyridine-2-carbonitrile (524mg, 67%) was obtained as a white solid.

10% Palladium-carbon (50mg) was added to a solution of the resulting 6-phenoxy-pyridine-2-carbonitrile (100mg, 0.51mmol) in methanol (5.0mL), the mixture was stirred at room temperature for 24 hours under hydrogen atmosphere (1 atm). The catalyst was removed by filtration, and the filtrate was concentrated to obtain the title compound (65mg, 64%) as a colorless oil.

Preparation Example 17. C-(5-(3-bromophenoxy)-thiophen-2-yl)-methylamine

To a solution of 5-nitrothiophene-2-carbonitrile (1.79g, 11.6mmol) and 3-bromophenol (2.00g, 11.6mmol) in dimethylsulfoxide (22mL) was added potassium carbonate (1.76g, 12.8mmol), and the solution was stirred at 70°C for 3 hours. The reaction solution was allowed to room temperature, ethyl acetate and water were added for partitioning, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 10 : 1), and 5-(3-bromophenoxy)-thiophene-2-carbonitrile (2.00g, 62%) was obtained as a yellow oil.

Next, to a solution of lithium aluminum hydride (204mg, 5.39mmol) in tetrahydrofuran (10mL) was added a solution of the resulting 5-(3-bromophenoxy)-thiophene-2-carbonitrile (1.01g, 3.59mmol) in tetrahydrofuran (10mL), and the solution was stirred at room temperature for 2 hours. Then, lithium aluminum hydride (68mg, 1.80mmol) was added, and the solution was further stirred at room temperature for 1 hour. Water was added to the reaction solution, which was then extracted with ethyl acetate, dried over anhydrous magnesium sulfate, and the solvent was evaporated to obtain a mixture of the title compound and



debrominated compound (740mg) as a pale brown oil. As this mixture cannot be purified, it was used in the next reaction without purification.

Preparation Example 18. C-(5-(3-Benzoyloxy-phenoxy)-thiophen-2-yl)-methylamine

5           To a solution of resorcinol (10g, 90.8mmol) in N,N-dimethylformamide (100mL) was added potassium carbonate (12.6g, 90.8mmol) and benzyl bromide (10.8mL, 90.8mmol), and the mixture was stirred at 60°C for 2 hours. The reaction mixture was allowed to room temperature, ethyl acetate and water were added for partitioning, the organic layer was washed with water and dried over anhydrous  
10   magnesium sulfate. The solvent was evaporated, then, the residue was purified by silica gel column chromatography (hexane : ethyl acetate), and 3-benzyloxy-phenol (6.0g, 33%) was obtained as a pale brown oil.

          To a solution of the resulting 3-benzyloxy-phenol (2.6g, 13.0mmol) and a 5-nitrothiophene-2-carbonitrile (2.0g, 13.0mmol) in dimethylsulfoxide (25mL) was  
15   added potassium carbonate (1.98g, 14.0mmol), and the solution was stirred at 70°C for 3 hours. The reaction solution was allowed to room temperature, ethyl acetate and water were added for partitioning, the organic layer was washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 10 : 1),  
20   and 5-(3-benzyloxy-phenoxy)-thiophene-2-carbonitrile (110mg, 2.8%) was obtained as a pale brown solid.

          Next, to a solution of lithium aluminum hydride (27mg, 0.716mmol) in tetrahydrofuran (2.0mL) was added a solution of  
5-(3-benzyloxy-phenoxy)-thiophene-2-carbonitrile obtained above (110mg,  
25   0.358mmol) in tetrahydrofuran (1mL), and the solution was stirred at room

temperature for 3 hours. Water was added to the reaction solution, which was then extracted with ethyl acetate, dried over anhydrous magnesium sulfate, and the solvent was evaporated to obtain the title compound (80mg, 72%) as a red solid. This compound was used in the next reaction without purification.

5

Preparation Example 19. (4-Aminomethylphenyl)-benzyl-amine

To a mixture of sodium *tert*-butoxide (7.44g, 77.4mmol), tris(dibenzylideneacetone)dipalladium(0) (0.38g, 0.415mmol) and rac-2,2-bis(diphenylphosphino)-1,1-binaphthyl (0.172g, 0.277mmol) was added a  
10 solution of 4-bromobenzonitrile (10g, 55.3mmol) and benzylamine (11.8g, 0.11mol) in toluene (100mL) under nitrogen atmosphere, and the solution was stirred at 80°C for 5 hours. The reaction solution was allowed to room temperature, then, filtrated through Celite pad to remove insoluble matter, and the filtrate was concentrated. The residue was purified by silica gel column chromatography (hexane : ethyl acetate),  
15 and 4-benzyl amino-benzonitrile (11.1g, 96%) was obtained as a yellow solid.

Next, to a solution of lithium aluminum hydride (911mg, 24.0mmol) in tetrahydrofuran (60mL) was added a solution of the resulting 4-benzyl amino-benzonitrile (2.0g, 9.61mmol) in tetrahydrofuran (5mL), and the solution was stirred at room temperature for 3 hours. Water was added to the reaction solution,  
20 which was then extracted with ethyl acetate, dried over anhydrous magnesium sulfate, and the solvent was evaporated to obtain the title compound (2.0g, quantitatively) as an orange oil. This compound was used in the next reaction without purification.

25 Preparation Example 20. (4-Aminomethyl-phenyl)-phenyl-amine

To a mixture of sodium *tert*-butoxide (7.44g, 77.4mmol), tris(dibenzylideneacetone)dipalladium(0) (0.38g, 0.415mmol) and rac-2,2-bis(diphenylphosphino)-1,1-binaphthyl (0.172g, 0.277mmol) was added a solution of 4-bromobenzonitrile (10g, 55.3mmol) and benzylamine (6.5mL, 0.11mol) in toluene (100mL) under nitrogen atmosphere, and the solution was stirred at 80°C for 5 hours. The reaction mixture was allowed to room temperature, then, filtered through Celite pad to remove insoluble matter, and the filtrate was concentrated. The residue was purified by silica gel column chromatography (hexane : ethyl acetate), and 4-phenylamino-benzonitrile (6.7g, 63%) was obtained as a yellow solid.

Next, to a solution of lithium aluminum hydride (1.17g, 30.9mmol) in tetrahydrofuran (60mL) was added a solution of the obtained 4-phenylamino-benzonitrile (2.0g, 10.3mmol) in tetrahydrofuran (5mL), and the solution was stirred at room temperature for 22 hours. Water was added to the reaction solution, which was then extracted with ethyl acetate, dried over anhydrous magnesium sulfate, and the solvent was evaporated to obtain the title compound (2.0g, 98%) as an orange oil. This compound was used in the next reaction without purification.

#### Preparation Example 21. (4-Aminomethyl-benzyl)-phenylamine

To a solution of 4-cyanobenzaldehyde (10g, 76.3mmol) and aniline (4.48mL, 76.3mmol) in tetrahydrofuran (370mL) were added acetic acid (21.9mL, 0.383mol) and triacetoxy sodium borohydride (32.3g, 0.153mol), and the solution was stirred at room temperature for 1 hour. Ethyl acetate and water were added to the reaction solution, which was then partitioned, the organic layer was washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue

was purified by silica gel column chromatography (hexane : ethyl acetate), and 4-phenylaminomethyl-benzonitrile (5.1g, 32%) was obtained as a pale yellow solid.

Next, to a solution of lithium aluminum hydride (0.91g, 24mmol) in tetrahydrofuran (60mL) was added a solution of the resulting 4-phenylaminomethyl-benzonitrile (2.0g, 9.61mmol) in tetrahydrofuran (5mL), and the solution was stirred at room temperature for 2 hours. Water was added to the reaction solution, which was then extracted with ethyl acetate, dried over anhydrous magnesium sulfate, and the solvent was evaporated to obtain the title compound (1.98g, 97%) as a yellow oil. This compound was used in the next reaction without purification.

#### Preparation Example 22. 5-(3-Fluorophenoxy)thiophene-2-carbonitrile

5-Nitrothiophene-2-carbonitrile (2g, 13mmol), 3-fluorophenol (1.75g, 15.6mmol) and potassium carbonate (3.6g, 26mmol) were suspended in dimethylsulfoxide (15mL), and the mixture was stirred at room temperature for 16 hours. Water and ethyl acetate were added for partitioning, silica gel was added to the organic layer, and the solvent was evaporated *in vacuo* for adsorption. Purification was carried out by silica gel column chromatography (hexane : ethyl acetate = 10 : 1), and the title compound (670mg, 3.1mmol, 23.5%) was obtained as an oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 6.83 (1H, d, J=4.0Hz), 7.08-7.26 (2H, m), 7.18-7.24 (1H, m), 7.49 (1H, ddd, J=8.0, 8.0, 8.0Hz), 7.81 (1H, d, J=4.0Hz).

#### Preparation Example 23. C-(5-(3-Fluorophenoxy)thiophen-2-yl)methylamine

To a solution of 5-(3-fluorophenoxy)thiophene-2-carbonitrile described in

Preparation Example 22 (670mg, 3mmol) in tetrahydrofuran (30mL) was added lithium aluminum hydride (460mg, 12mmol), and the solution was stirred at room temperature for 16 hours. Water and ethyl acetate were added to the reaction solution, which was then partitioned, the organic layer was filtered with NH silica gel, the filtrate was evaporated *in vacuo*, and the title compound (570mg, 2.42mmol, 80.7%) was obtained as a brown oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.08 (2H, brs), 3.80 (2H, s), 6.54 (1H, d, J=3.6Hz), 6.66-6.70 (1H, m), 6.88-6.99 (3H, m), 7.39 (1H, ddd, J=8.0, 8.0, 8.0Hz).

10 Preparation Example 24. C-(5-Phenoxy thiophen-2-yl)methylamine

5-Nitrothiophene-2-carbonitrile (0.80g, 5.2mmol), phenol (1.0g, 10.4mmol) and potassium carbonate (2.2g, 15.6mmol) were suspended in dimethylsulfoxide (30mL), and the mixture was stirred at room temperature for 15.5 hours. Water and ethyl acetate were added for partitioning, and the organic layer was washed three times with water. NH silica gel was added to the organic layer, the solvent was evaporated *in vacuo* for adsorption, purification was carried out by NH silica gel column chromatography (hexane : ethyl acetate = 8 : 1), and 5-phenoxythiophene-2-carbonitrile (720mg, 3.6mmol, 69.2%) was obtained as a colorless oil.

20 To a solution of this oil in tetrahydrofuran (40mL) was added lithium aluminum hydride (540mg, 14.4mmol), and the mixture was stirred for 30 minutes at room temperature. Water and ethyl acetate were added to the reaction mixture, which was then partitioned, silica gel was added to the organic layer, the solvent was evaporated *in vacuo* for adsorption, purification was carried out by silica gel column chromatography (ethyl acetate, then ethyl acetate : methanol = 4 : 1), and the title

25

compound (570mg, 2.8mmol, 77.2%) was obtained as a light brown oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.34 (2H, brs), 3.78-3.82(2H, m), 6.47 (1H, d, J=4.0Hz), 6.65-6.68 (1H, m), 7.04-7.14 (3H, m), 7.34-7.40(2H, m).

5 Preparation Example 25. 5-Phenoxy thiophene-2-carbonitrile

5-Nitro-thiophene-2-carbonitrile(1.5g, 9.7mmol), phenol (1.8g, 19.4mmol) and potassium carbonate(4.0g, 29.1mmol) were suspended in dimethylsulfoxide (20mL), and the mixture was stirred for 50 minutes at 60°C, followed by further stirring overnight at room temperature. Water and ethyl acetate were added to the reaction mixture, which was then partitioned, the organic layer was washed 4 times with water, then, NH silica gel was added to the organic layer, the solvent was evaporated *in vacuo* for adsorption, purification was carried out by NH silica gel column chromatography (hexane, then hexane : ethyl acetate = 20 : 1, then 10 : 1), and the title compound (1.4g, 7.0mmol, 72.1%) was obtained as a pale yellow oil.

10

15 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 6.75 (1H, d, J=4.0Hz), 7.23-7.31 (3H, m), 7.42-7.49 (2H, m), 7.78 (1H, d, J=4.0Hz).

Preparation Example 26. C-(5-Phenoxy thiophen-2-yl)methylamine

To a solution of 5-phenoxythiophene-2-carbonitrile described in Preparation Example 25 (1.4g, 7.0mmol) in tetrahydrofuran (30mL) was added lithium aluminum hydride (1.1g, 28mmol), and the solution was stirred at room temperature for 25 minutes. Water and ethyl acetate were added to the reaction solution, this mixed solution was filtered through Celite pad, furthermore, the organic layer was partitioned. The solvent was evaporated *in vacuo*, and the title compound (1.29g, 6.3mmol, 89.9%) was obtained as a brown oil.

20

25

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 1.92 (2H, brs), 3.74-3.80(2H, m), 6.46 (1H, d, J=3.6Hz), 6.62-6.66(1H, m), 7.02-7.14 (3H, m), 7.32-7.39 (2H, m).

Preparation Example 27. 5-(4-Fluorophenoxy)thiophene-2-carbonitrile

5            5-Nitrothiophene-2-carbonitrile (2.0g, 13mmol), 4-fluorophenol (2.9g, 26mmol) and potassium carbonate (5.4g, 39mmol) were suspended in dimethylsulfoxide (30mL), and the mixture was stirred for 30 minutes at 60°C. Water and ethyl acetate were added to the reaction mixture, which was then separated, the organic layer was washed with water twice, then, NH silica gel was  
10 added to the organic layer, the solvent was evaporated *in vacuo* for adsorption, purification was carried out by NH silica gel column chromatography (hexane : ethyl acetate = 20 : 1), and the title compound (3.7g, containing 4-fluorophenol) was obtained as a brown oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 6.68-6.76 (2H, m), 7.26-7.38 (3H, m),  
15 7.74-7.80 (1H, m).

Preparation Example 28. C-(5-(4-Fluorophenoxy)thiophen-2-yl)methylamine

To a solution of 5-(4-fluorophenoxy)thiophene-2-carbonitrile described in Preparation Example 27 (containing 4-fluorophenol) (3.7g) in tetrahydrofuran (40mL)  
20 was added lithium aluminum hydride (1.3g, 34mmol), and the solution was stirred at room temperature for 30 minutes. Water and ethyl acetate were added to the reaction solution, which was then partitioned, the solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (ethyl acetate, then ethyl acetate : methanol = 4 : 1), and the title compound (1.2g, 5.4mmol) was  
25 obtained as a brown oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.00 (2H, brs), 3.75-3.80 (2H, m), 6.44-6.48 (1H, m), 6.62-6.67 (1H, m), 7.08-7.14 (2H, m), 7.16-7.24 (2H, m).

Preparation Example 29. 5-*m*-Tolyloxy-thiophene-2-carbonitrile

5        The title compound (960mg, 4.47mmol, 68.7%) was obtained as a yellow oil from 5-nitro-thiophene-2-carbonitrile (1.0g, 6.5mmol) and 3-methylphenol (1.4g, 13mmol) according to an analogous method to Preparation Example 27.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.31 (3H, s), 6.73 (1H, dd, J=0.8, 4.0Hz), 7.03-7.06 (1H, m), 7.07-7.12 (2H, m), 7.33 (1H, dd, J=8.0, 8.0Hz), 7.77 (1H, dd,  
10 J=0.8, 4.0Hz).

Preparation Example 30. C-(5-*m*-Tolyloxy-thiophen-2-yl)-methylamine

The title compound (900mg, 4.10mmol, 91.7%) was obtained as a reddish brown oil from 5-*m*-tolyloxy thiophene-2-carbonitrile described in Preparation  
15 Example 29 (960mg, 4.47mmol) according to an analogous method to Preparation Example 28.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.03 (2H, brs), 2.34 (3H, s), 3.85 (2H, s), 6.51-6.54 (1H, m), 6.71-6.74 (1H, m), 6.90-7.03 (3H, m), 7.31 (1H, dd, J=8.0, 8.0Hz).

20    Preparation Example 31. 5-*p*-Tolyloxy-thiophene-2-carbonitrile

The title compound (1.0g, 4.65mmol, 71.5%) was obtained as a yellow oil from 5-nitro-thiophene-2-carbonitrile (1.0g, 6.5mmol) and 4-methylphenol (1.4g, 13mmol) according to an analogous method to Preparation Example 27.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.30 (3H, s), 6.69-6.71 (1H, m), 7.15-7.18  
25 (2H, m), 6.24-6.28 (2H, m), 7.15-7.78 (1H, m).



Preparation Example 32. C-(5-*p*-Tolyloxy thiophen-2-yl)methylamine

The title compound (780mg, 3.56mmol, 76.5%) was obtained as a reddish brown oil from 5-*p*-tolylloxy thiophene-2-carbonitrile described in Preparation Example 31 (1.0g, 4.65mmol) according to an analogous method to Preparation Example 28 .

<sup>1</sup>H-NMR Spectrum (DMSO-*d*<sub>6</sub>) δ(ppm) : 2.06 (2H, br), 2.22 (3H, s), 3.76 (2H, s), 6.41 (1H, d, J=3.6Hz), 6.62 (1H, d, J=3.6Hz), 6.90-6.98 (2H, m), 7.15-7.18 (2H, m).

Preparation Example 33. 2-(4-(3-Fluoro-phenoxy)-thiophen-2-yl)-[1,3]dioxolane

2-(4-Bromo-thiophen-2-yl)-[1,3]dioxolane (1.0g, 4.3mmol), 3-fluorophenol (0.95g, 8.6mmol), 2,2,6,6-tetramethyl-3,5-heptanedione (0.078g, 0.43mmol), copper(I) chloride (0.21g, 2.7mmol) and cesium carbonate (2.8g, 8.6mmol) were suspended in N-methylpyrrolidone (10mL) under a nitrogen stream, and the mixture was stirred for 4.5 hours at 120°C. To this suspension was added 2,2,6,6-tetramethyl-3,5-heptanedione (0.12g, 0.65mmol), and the solution was further stirred for 8 hours at 140°C. The reaction mixture was filtered through Celite pad, then, water and ethyl acetate were added for partitioning, and the organic layer was washed with water twice. NH silica gel was added to the organic layer, the solvent was evaporated *in vacuo* for adsorption to NH silica gel, purification was carried out by NH silica gel column chromatography (hexane, then hexane : ethyl acetate = 30 : 1), and the title compound (280mg, 1.05mmol, 24.4%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (DMSO-*d*<sub>6</sub>) δ(ppm) : 3.88-3.96 (2H, m), 3.96-4.04 (2H, m), 5.98 (1H, s), 6.82-6.88 (2H, m), 6.91-6.97 (1H, m), 7.04-7.05 (1H, m), 7.09 (1H, d, J=2.0Hz), 7.35-7.42 (1H, m).

Preparation Example 34. 4-(3-Fluorophenoxy)thiophene-2-carbaldehyde

To a solution of 2-(4-(3-fluoro-phenoxy)-thiophen-2-yl)-[1,3]dioxolane described in Preparation Example 33 (280mg, 1.05mmol) in methanol (10mL) was added citric acid aqueous solution (10mL), and the solution was stirred at room temperature for 30 minutes. The reaction solution was neutralized with an aqueous solution of sodium bicarbonate, the solution was extracted with ethyl acetate, the organic layer was dried over anhydrous sodium sulfate, the solvent was evaporated *in vacuo*, and the title compound (210mg, 0.95mmol, 90%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 6.88-7.03 (3H, m), 7.38-7.46 (1H, m), 7.67(1H, d, J=1.6Hz), 7.88 (1H, d, J=1.6Hz), 9.86 (1H, s).

Preparation Example 35. (C-(4-(3-Fluorophenoxy)thiophen-2-yl)methylamine

4-(3-Fluorophenoxy)thiophene-2-carbaldehyde described in Preparation Example 34 (210mg, 0.95mmol) was dissolved in 7N ammonia/methanol (30mL), Raney nickel (500mg) was added thereto, followed by stirring for 19 hours under hydrogen atmosphere at room temperature. The reaction mixture was filtered through Celite pad, Raney nickel was removed, silica gel was added to the filtrate, then, the solvent was evaporated *in vacuo*, for adsorption on silica gel, purification by silica gel chromatography (ethyl acetate, then ethyl acetate : methanol = 4 : 1) was carried out, and the title compound (70mg, 0.32mmol) was obtained as a pale yellow oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.11(2H, brs), 3.82(2H, s), 6.75(1H, s), 6.80-6.95 (4H, m), 7.33-7.41(1H, m).

Preparation Example 36. 2-(5-(4-Fluoro-benzyl)-thiophen-2-yl)-[1,3]dioxolane

*n*-Butyl lithium (2.6N hexane solution, 3.3mL, 8.47mmol) was added dropwise to a solution of 2-(5-bromo-thiophen-2-yl)-[1,3]dioxolane (1.8g, 7.7mmol) in tetrahydrofuran (20mL) that had been cooled to from -75° to -70°C, and the solution  
5 was stirred for 30 minutes. 4-Fluorobenzyl bromide (1.1mL, 8.47mmol) was added dropwise to this reaction solution while keeping it at -70°C or less. After completion of dropwise addition, the reaction solution was gradually allowed to room temperature. Water and ethyl acetate were added to the reaction solution, which was then partitioned, silica gel was added to the organic layer, which was then  
10 evaporated *in vacuo* for adsorption, purification by silica gel column chromatography (hexane, then hexane : ethyl acetate = 20 : 1, then 10 : 1) was carried out, and the title compound (560mg, 2.04mmol, 26.4%) was obtained as a brown oil.

<sup>1</sup>H-NMR Spectrum (DMSO-*d*<sub>6</sub>) δ(ppm) : 3.84-3.98 (4H, m), 4.08 (2H, s), 5.90 (1H, s), 6.75-6.78 (1H, m), 7.00 (1H, d, J=3.6Hz), 7.08-7.15 (2H, m), 7.25-7.30 (2H, m).

15

Preparation Example 37. 5-(4-Fluoro-benzyl)-thiophene-2-carbaldehyde

To a solution of 2-(5-(4-fluoro-benzyl)-thiophen-2-yl)-[1,3]dioxolane described in Preparation Example 36 (560mg, 2.04mmol) in methanol (20mL) was added citric acid aqueous solution (20mL), and the solution was stirred at room temperature for  
20 30 minutes. Ethyl acetate and water were added to the reaction solution, which was then partitioned, the organic layer was washed with water twice, dried over anhydrous sodium sulfate, the solvent was evaporated *in vacuo*, and title compound (460mg, 2.09mmol) was obtained as a brown oil.

<sup>1</sup>H-NMR Spectrum (DMSO-*d*<sub>6</sub>) δ(ppm) : 4.22 (2H, s), 7.08-7.18 (3H, m), 7.29-7.36  
25 (2H, m), 7.83-7.87 (1H, m), 9.79 (1H, s).

Preparation Example 38. C-(5-(4-Fluorobenzyl)thiophen-2-yl)methylamine

7N ammonia/methanol(30mL) and Raney nickel (500mg) were added to 5-(4-fluorobenzyl)thiophene-2-carbaldehyde described in Preparation Example 37 (460mg, 2.09mmol), and the mixture was stirred for 14 hours under hydrogen atmosphere at room temperature. The catalyst was removed by filtering through Celite pad, then, purification by silica gel column chromatography (ethyl acetate, then ethyl acetate : methanol = 4 : 1) was carried out, and the title compound (70mg, 0.316mol, 15.1%) was obtained as an oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.20 (2H, brs), 3.78 (2H, s), 4.03 (2H, s), 6.58-6.80 (2H, m), 7.00-7.38 (4H, m).

Preparation Example 39. 5-Benzyl-furan-2-carbaldehyde

*n*-Butyl lithium (2.44N hexane solution, 15mL, 39.6mmol) was added dropwise to a solution of 2-furan-2-yl-[1,3]dioxolane (5g, 36mmol) in tetrahydrofuran (30mL) that had been cooled to from -75°C to -70°C, and the solution was stirred for 1 hour. Benzyl bromide (4.7mL, 39.6mmol) was added dropwise to this solution at from -75°C to -70°C. After completion of dropwise addition, the cold bath was removed, and the solution was gradually allowed to room temperature. Water and ethyl acetate were added to the reaction mixture for partitioning, silica gel was added to the organic layer, which was then evaporated *in vacuo* to adsorb the reaction mixture, purification by silica gel column chromatography (hexane : ethyl acetate = 50 : 1, then 6 : 1) was carried out, and 2-(5-benzyl-furan-2-yl)-[1,3]dioxolane (3.8g, 16.5mmol, 45.9%) was obtained as a yellow oil.

The resulting 2-(5-benzyl-furan-2-yl)-[1,3]dioxolane (3.8g, 16.5mmol) was

suspended in a mixture solution of methanol (15mL), tetrahydrofuran (10mL) and 2N hydrochloric acid (15mL), and the solution was stirred at room temperature for 2 hours. Ethyl acetate and an aqueous solution of sodium bicarbonate were added to the reaction solution, which was then partitioned, the organic layer was washed with water twice and dried over anhydrous sodium sulfate. The solvent was evaporated, and the title compound (2.5g, 13mmol) was obtained as a brown oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.09 (2H, s), 6.45-6.48 (1H, m), 7.20-7.35 (5H, m), 7.45 (1H, d, J=3.6Hz), 9.46 (1H, s).

10 Preparation Example 40. 2-(5-Benzyl-thiophen-2-yl)-[1,3]dioxolane

The title compound (520mg, 2.1mmol, 41.4%) was obtained as a colorless oil from benzyl bromide according to an analogous method to Preparation Example 36.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.84-3.90 (2H, m), 3.90-3.98 (2H, m), 4.08 (2H, s), 5.90 (1H, s), 6.75-6.78 (1H, m), 7.00 (1H, d, J=3.6Hz), 7.18-7.32 (5H, m).

15

Preparation Example 41. 5-Benzyl-thiophene-2-carbaldehyde

The title compound (containing impurity, 460mg) was obtained as a colorless oil from 2-(5-benzyl-thiophen-2-yl)-[1,3]dioxolane described in Preparation Example 40 (520mg, 2.1mmol) according to an analogous method to Preparation Example 37.

20 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.22 (2H, s), 7.11 (1H, d, J=3.6Hz), 7.20-7.34 (5H, m), 7.85 (1H, d, J=3.6Hz), 9.79 (1H, s).

Preparation Example 42. C-(5-benzyl-thiophen-2-yl)-methylamine

The title compound (270mg) was obtained as a brown oil from 5-benzyl-thiophene-2-carbaldehyde described in Preparation Example 41 (containing

impurity, 460mg, 2.27mmol) according to an analogous method to Preparation Example 38.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 1.85 (2H, brs), 3.75 (2H, s), 4.01 (2H, s), 6.65-6.72 (2H, m), 7.15-7.30 (5H, m).

5

Preparation Example 43. 2-(5-(3-Chloro-benzyl)-thiophen-2-yl)-[1,3]dioxolane

*n*-Butyl lithium (2.6N hexane solution, 15.6mL, 39mmol) was added dropwise to a solution of 2-(5-bromo-thiophen-2-yl)-[1,3]dioxolane (7.0g,30mmol) in tetrahydrofuran (40mL) at from -75° to -68°C, and the solution was stirred for 20 minutes. 3-Chlorobenzyl bromide (4.3mL,33mmol) was added dropwise to this reaction solution at from -75°C to -68°C, and the solution was stirred for 20 minutes. The cold bath was removed, and the reaction solution was gradually allowed to room temperature. Water and ethyl acetate were added to the reaction solution, which was then partitioned, the solvent was evaporated, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 20 : 1), and the title compound (1.6g, 5.7mmol, 19.0%) was obtained as a yellow oil.

15

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.82-4.00 (4H, m), 4.11 (2H, s), 5.91 (1H, s), 6.78-6.80 (1H, m), 7.01 (1H, d, J=3.6Hz), 7.19-7.36 (4H, m).

20 Preparation Example 44. 5-(3-Chloro-benzyl)-thiophene-2-carbaldehyde

To a solution of 2-(5-(3-chloro-benzyl)-thiophen-2-yl)-[1,3]dioxolane described in Preparation Example 43 (1.6g,5.7mmol) in methanol (20mL) was added citric acid aqueous solution (20mL), and the solution was stirred at room temperature for 20 minutes. An aqueous solution of sodium bicarbonate and ethyl acetate were added to the reaction solution, which was then partitioned, the organic layer was washed

25

with water twice and dried over anhydrous magnesium sulfate. The solvent was evaporated, and the title compound (1.2g, 5.08mmol, 89.2%) was obtained as a yellow oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.25 (2H, s), 7.12-7.15 (1H, m), 7.24-7.40 (4H, m), 7.86 (1H, d, J=3.6Hz), 9.80 (1H, s).

Preparation Example 45. C-(5-(3-Chloro-benzyl)-thiophen-2-yl)-methylamine

To a solution of 5-(3-chloro-benzyl)thiophene-2-carbaldehyde described in Preparation Example 44 (1.2g, 5.08mmol) in 7N ammonia/methanol (40mL) was added Raney nickel (2g), and the solution was stirred at room temperature for 17 hours under hydrogen atmosphere. The reaction solution was filtered through Celite pad, the catalyst was removed, then, this filtrate was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (ethyl acetate, then ethyl acetate : methanol = 5 : 1), and the title compound (740mg, 3.12mmol, 61.4%) was obtained as a brown oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 1.94 (2H, brs), 3.76 (2H, s), 4.06 (2H, s), 6.69-6.72 (2H, m), 7.18-7.34 (4H, m).

Preparation Example 46. 5-(4-Chloro-phenoxy)-furan-2-carbaldehyde

To a solution of 4-chlorophenol (4.4g, 33.6mmol) in dimethylsulfoxide (30mL) was added sodium hydride (1.34g, 33.6mmol, 60% in oil), and the solution was stirred at room temperature for 10 minutes. 5-Nitrofuran-2-carbaldehyde (4.0g, 28mmol) was added to this reaction solution, and the solution was stirred at room temperature for 5 minutes. Water and ethyl acetate were added to the reaction solution, which was then partitioned, and the organic layer was washed with water 6 times. The

solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane, then hexane : ethyl acetate = 10 : 1, then 4 : 1), and the title compound (3.3g, 14.9mmol, 53.0%) was obtained as a yellow oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 5.89-5.93 (1H, m), 7.30-7.36 (2H, m),  
5 7.50-7.60 (3H, m), 9.35-9.38 (1H, m).

Preparation Example 47. C-(5-(4-Chloro-phenoxy)-furan-2-yl)-methylaniline

The title compound (200mg, 0.90mmol, 8.7%) was obtained as a brown oil from 5-(4-chloro-phenoxy)-furan-2-carbaldehyde described in Preparation Example  
10 46 (2.3g, 10.3mmol) according to an analogous method to Preparation Example 38.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.56 (2H, s), 5.73 (1H, d, J=3.2Hz), 6.18 (1H, d, J=3.2Hz), 7.03-7.08 (2H, m), 7.40-7.45 (2H, m).

Preparation Example 48. 5-Phenoxy-furan-2-carbaldehyde

15 The title compound (2.3g, 12.2mmol, 43.5%) was obtained as a light brown oil from phenol (3.2g, 33.6mmol) and 5-nitro-furan-2-carbaldehyde (4.0g, 28mmol) according to an analogous method to Preparation Example 46.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 5.85 (1H, d, J=4.0Hz), 7.25-7.33 (3H, m), 7.45-7.50 (2H, m), 7.57 (1H, d, J=4.0Hz), 9.34 (1H, s).

20

Preparation Example 49. C-(5-Phenoxy-furan-2-yl)-methylaniline

The title compound (250mg, 1.32mmol, 24.9%) was obtained as a yellow oil from 5-phenoxy-furan-2-carbaldehyde described in Preparation Example 48 (1.0g, 5.3mmol) according to an analogous method to Preparation Example 38.

25 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 1.98 (2H, brs), 3.56 (2H, s), 5.67 (1H, d,



J=3.2Hz), 6.16-6.18 (1H, m), 6.99-7.04 (2H, m), 7.10-7.16 (1H, m), 7.34-7.40 (2H, m).

Preparation Example 50. 5-(3-Fluoro-phenoxy)-furan-2-carbaldehyde

5           The title compound (1.5g, 7.3mmol, 52.1%) was obtained as a yellow oil from 3-fluorophenol (1.9g, 16.8mmol) and 5-nitro-furan-2-carbaldehyde (2.0g, 14mmol) according to an analogous method to Preparation Example 46.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 5.97-6.00 (1H, m), 7.10-7.20 (2H, m), 7.24-7.30 (1H, m), 7.48-7.55 (1H, m), 7.59 (1H, d, J=3.6Hz) 9.37 (1H, s).

10

Preparation Example 51. (5-(3-Fluoro-phenoxy)-furan-2-yl)-methanol

          To a solution of 5-(3-fluoro-phenoxy)-furan-2-carbaldehyde described in Preparation Example 50 (1.5g, 7.3mmol) in tetrahydrofuran (20mL) was added sodium borohydride (280mg, 7.3mmol), and the solution was stirred at room  
15   temperature for 1 hour. Water and ethyl acetate were added to the reaction solution, which was then partitioned, the organic layer was washed with water twice, and the organic layer was filtered through a glass filter lined with silica gel. The solvent was evaporated *in vacuo*, and the title compound (1.5g, 7.2mmol) was obtained as a pale yellow oil.

20   <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.29 (2H, d, J=6.0Hz), 5.17 (1H, t, J=6.0Hz), 5.78-5.82 (1H, m), 6.78-6.82 (1H, m), 6.85-6.95 (2H, m), 6.98-7.04 (1H, m), 7.40-7.46 (1H, m).

Preparation Example 52. 2-(5-(3-Fluoro-benzyl)-thiophen-2-yl)-[1,3]dioxolane

25           *n*-Butyl lithium (2.44N hexane solution, 6.4mL, 16.9mmol) was added

dropwise to a solution of 2-(5-bromo-thiophen-2-yl)-[1,3]dioxolane (3.0g,13mmol) in tetrahydrofuran (30mL) at from -75°C to -69°C, and the solution was stirred for 17 minutes. 3-Fluorobenzyl bromide (1.7mL, 14.3mmol) was added dropwise to this reaction solution at from -75°C to -69°C. After dropwise addition was completed, the reaction solution was gradually allowed to room temperature. Water and ethyl acetate were added to the reaction solution, which was then partitioned, the organic layer was concentrated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 20 : 1), and the title compound (478mg, 1.81mmol, 13.9%) was obtained as a brown oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.82-4.00 (4H, m), 4.13 (2H, s), 5.91 (1H, s), 6.78-6.80 (1H, m), 7.00-7.10 (4H, m), 7.30-7.37 (1H, m).

#### Preparation Example 53. 5-(3-Fluoro-benzyl)-thiophene-2-carbaldehyde

An aqueous solution of saturated citric acid (20mL) was added to a solution of 2-(5-(3-fluoro-benzyl)thiophen-2-yl)-[1,3]dioxolane described in Preparation Example 52 (670mg, 2.53mmol) in methanol (20mL), and the solution was stirred at room temperature for 30 minutes. Water and ethyl acetate were added to the reaction solution, which was then partitioned, and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, and the title compound (485mg, 2.2mmol, 87.0%) was obtained as a brown oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.25 (2H, s), 7.03-7.18 (4H, m), 7.30-7.40 (1H, m), 7.86 (1H, dd, J=1.6, 3.6Hz), 9.80 (1H, d, J=1.6Hz).

#### Preparation Example 54. 2-(5-(3-Chloro-benzyl)-furan-2-yl)-[1,3]dioxolane

The title compound (1.34g, 5.07mmol, 14.1%) was obtained as a yellow oil

from 2-furan-2-yl-[1,3]dioxolane (5.0g,36mol) and 3-chlorobenzyl bromide (5.2mL,39.6mmol) according to an analogous method to Preparation Example 36.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.84-4.02 (6H, m), 5.76 (1H, s), 6.10-6.12 (1H, m), 6.41 (1H, d, J=3.2Hz), 7.16-7.20 (1H, m), 7.26-7.36 (3H, m).

5

Preparation Example 55. 5-(3-Chloro-benzyl)furan-2-carbaldehyde

The title compound (1.03g, 4.68mmol, 82.1%) was obtained as a yellow oil from 2-(5-(3-chloro-benzyl)-furan-2-yl)-[1,3]dioxolane described in Preparation Example 54 (1.34g, 5.07mmol) according to an analogous method to Preparation

10 Example 37.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.11 (2H, s), 6.48-6.51 (1H, m), 7.20-7.24 (1H, m), 7.28-7.37 (3H, m), 7.46 (1H, d, J=3.2Hz), 9.46-9.49 (1H, m).

Preparation Example 56. C-(5-(3-Chloro-benzyl)-furan-2-yl)-methylamine

15 The title compound (690mg, 3.12mmol, 66.6%) was obtained as a yellow oil from 5-(3-Chloro-benzyl)-furan-2-carbaldehyde described in Preparation Example 55 (1.03g, 4.68mmol) according to an analogous method to Preparation Example 38.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 1.90 (2H, brs), 3.57 (2H, s), 3.92 (2H, s), 6.01 (1H, d, J=2.8Hz), 6.07 (1H, d, J=2.8Hz), 7.16-7.20 (1H, m), 7.24-7.34 (3H, m).

20

Preparation Example 57. 1-Benzyl-1H-pyrrole-3-carbaldehyde

Benzylamine (540mg, 5.00mmol) and acetic acid (10mL) were added to 2,5-dimethoxy-tetrahydrofuran-3-carbaldehyde (1g, 6.25mmol), and the solution was stirred for 20 minutes at 90°C. Water and ethyl acetate were added to the reaction  
25 solution, which was then partitioned, the organic layer was washed with an aqueous

solution 2N sodium hydroxide once and with water twice, filtered with a glass filter lined with silica gel, the filtrate was evaporated *in vacuo*, and title compound (800mg, 4.3mmol, 68.8%) was obtained as a brown oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 5.18 (2H, s), 6.46 (1H, dd, J=2.0, 2.0Hz),  
 5 6.97 (1H, dd, J=2.0, 2.0Hz), 7.24-7.37 (5H, m), 7.71 (1H, dd, J=2.0, 2.0Hz), 9.63 (1H, s).

#### Preparation Example 58. 1-(3-Fluoro-benzyl)-1H-pyrrole-3-carbaldehyde

The title compound (2.33g, 11.4mmol, 71.7%) was obtained as an oil from  
 10 2,5-dimethoxy-tetrahydrofuran-3-carbaldehyde (2.6g, 16mmol) and  
 3-fluorobenzylamine (2.0g, 16mmol) according to an analogous method to  
 Preparation Example 57.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 5.20 (2H, s), 6.44-6.48 (1H, m), 6.98-7.02  
 (1H, m), 7.07-7.17 (3H, m), 7.36-7.42 (1H, m), 7.74 (1H, d, J=1.6Hz), 9.63 (1H, s).

15

#### Preparation Example 59. C-(1-(3-Fluoro-benzyl)-1H-pyrrol-3-yl)-methylaniline

7N Ammonia/methanol(40mL) and Raney nickel(2g) were added to  
 1-(3-fluoro-benzyl)-1H-pyrrole-3-carbaldehyde described in Preparation Example 58  
 (1.0g,4.9mmol), and the solution was stirred for 18 hours under hydrogen  
 20 atmosphere at room temperature. The catalyst was removed by filtering through  
 Celite pad, then, NH silica gel was added to the filtrate, the solvent was evaporated *in vacuo*  
 for adsorption, purification by NH silica gel column chromatography (hexane :  
 ethyl acetate = 2 : 1, then 1 : 1, then ethyl acetate) was carried out, and the title  
 compound (530mg, 2.5mmol, 53.0%) was obtained as a brown oil.

25 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.50 (2H, s), 5.01 (2H, s), 5.95 (1H, d,

J=2.0Hz), 6.64 (1H, d, J=1.2Hz), 6.69-6.74 (1H, m), 6.92-7.10 (3H, m), 7.32-7.38 (1H, m).

**Preparation Example 60. 1-Benzo[1,3]dioxol-5-ylmethyl-1*H*-pyrrole-3-carbaldehyde**

5 The title compound (2.0g, 8.7mmol, 69.8%) was obtained as a brown oil from 2,5-dimethoxy-tetrahydrofuran-3-carbaldehyde (2.0g, 12.5mmol) and C-benzo[1,3]dioxol-5-ylmethylamine) (1.9g, 12.5mmol) according to an analogous method to Preparation Example 57.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 5.04 (2H, s), 5.97 (2H, s), 6.42-6.45 (1H, m),  
10 6.80-7.00 (4H, m), 7.69-7.72 (1H, m), 9.61 (1H, s).

Preparation	Example	61.
-------------	---------	-----

C-(1-benzo[1,3]dioxol-5-ylmethyl-1*H*-pyrrol-3-yl)-methylamine

The title compound (1.5g, 6.5mmol, 74.7%) was obtained as a light green oil from 1-benzo[1,3]dioxol-5-ylmethyl-1*H*-pyrrole-3-carbaldehyde described in Preparation Example 60 (2.0g, 8.7mmol) according to an analogous method to Preparation Example 59.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 1.37 (2H, brs), 3.48 (2H, brs), 4.87 (2H, s), 5.90 (1H, s), 5.95 (2H, s), 6.60 (1H, s), 6.66-6.72 (2H, m), 6.75 (1H, s), 6.83 (1H, d, J=7.6Hz).

### Preparation Example 62. 1-Phenethyl-1*H*-pyrrol-3-carbaldehyde

The title compound (840mg, 4.2mmol, 84%) was obtained as a brown oil from 2,5-dimethoxy-tetrahydrofuran-3-carbaldehyde (1.0g, 6.25mmol) and phenethyl  
amine (600mg, 5.0mmol) according to an analogous method to Preparation Example

57.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.03 (2H, t, J=7.2Hz), 4.19 (2H, t, J=7.2Hz), 6.38-6.42 (1H, m), 6.37-6.82 (1H, m), 7.15-7.30 (5H, m), 7.54-7.60 (1H, m), 9.57 (1H, s).

5

Preparation Example 63. 1-Benzyloxy-1H-pyrrole-3-carbaldehyde

The title compound (500mg, 2.5mmol, 13.1%) was obtained as a yellow oil from 2,5-dimethoxy-tetrahydrofuran-3-carbaldehyde (3.0g, 19mmol) and O-benzyl hydroxylamine (2.3g, 19mmol) according to an analogous method to Preparation

10 Example 57.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 5.24 (2H, s), 6.35-6.48 (1H, m), 7.05-7.08 (1H, m), 7.28-7.43 (5H, m), 7.74-7.77 (1H, m), 9.55 (1H, s).

Preparation Example 64. (1-Benzyloxy-1H-pyrrol-3-yl)-methanol

15 To a solution of 1-benzyloxy-1H-pyrrole-3-carbaldehyde described in Preparation Example 63 (500mg, 2.5mmol) in tetrahydrofuran (10mL) was added lithium aluminum hydride (75mg, 1.97mmol), and the solution was stirred at room temperature for 10 minutes. Water and ethyl acetate were added to the reaction solution, which was then partitioned, silica gel was added to the organic layer, the  
20 solvent was evaporated *in vacuo* for adsorption, purification by silica gel column chromatography (hexane : ethyl acetate = 2 : 1) was carried out, and the title compound (168mg, 0.828mmol, 33.1%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.22 (2H, d, J=5.6Hz), 4.60 (1H, t, J=5.6Hz), 5.10 (2H, s), 5.78-5.81 (1H, m), 6.75-6.78 (1H, m), 6.78-6.81 (1H, m), 7.37-7.42 (5H,  
25 m).

<u>Preparation</u>	<u>Example</u>	<u>65.</u>
--------------------	----------------	------------

(5-[1,3]Dioxolan-2-yl-thiophen-2-yl)-(5-methyl-thiophen-2-yl)-methanol

*n*-Butyl lithium (2.44N hexane solution, 7.4mL, 17.9mmol) was added dropwise to a solution of 2-(5-bromo-thiophen-2-yl)-[1,3]dioxolane (4.0g, 17mmol) in tetrahydrofuran (50mL) at from -75°C to -70°C, and the solution was stirred for 10 minutes. 5-Methylthiophene-2-carbaldehyde (2.4g, 18.7mmol) was further added dropwise to the reaction solution at from -75°C to -70°C. After dropwise addition was completed, the reaction solution was gradually allowed to room temperature.

Water and ethyl acetate were added to the reaction mixture, which was then partitioned, and the solvent was evaporated. The residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 4 : 1, then 2 : 1), and the title compound (2.0g, 7.09mmol, 41.7%) was obtained as a brown oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.38 (3H, s), 3.88-4.04 (4H, m), 5.95 (1H, s), 6.04-6.08 (1H, m), 6.43-6.46 (1H, m), 6.60-6.63 (1H, m), 6.75 (1H, d, J=3.6Hz), 6.83 (1H, d, J=3.6Hz), 7.02 (1H, d, J=3.6Hz).

Preparation Example 66. 5-(5-Methyl-thiophen-2-ylmethyl)-thiophene-2-carbaldehyde

Sodium iodide (6.4g, 42.6mmol) and trimethylsilyl chloride (4.6g, 42.6mmol) were suspended in acetonitrile (100mL) on an ice bath, and (5-[1,3]dioxolane-2-ylthiophen-2-yl)-(5-methyl-thiophen-2-yl)-methanol described in Preparation Example 65 (2.0g, 7.09mmol) was added dropwise. After dropwise addition was completed, the reaction solution was gradually allowed to room temperature. A solution obtained by dissolving an aqueous solution of 2N sodium hydroxide (10.6mL) and sodium thiosulfate pentahydrate (530mg, 2.13mmol) in water

(5mL) was added to the reaction solution that had been cooled on an ice bath, and the solution was stirred. Water and ethyl acetate were added to the reaction solution, which was then partitioned, and the organic layer was concentrated *in vacuo*. The residue was purified by silica gel column chromatography (hexane : ethyl acetate = 20 : 1), and the title compound (790mg, 3.56mmol, 50.2%) was obtained as a brown oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.38 (3H, s), 4.38 (2H, s), 6.62-6.67 (1H, m), 6.76-6.80 (1H, m), 7.12-7.16 (1H, m), 7.85-7.90 (1H, m), 9.83 (1H, s).

#### 10 Preparation Example 67. (5-(5-Methyl-thiophen-2-ylmethyl)-thiophen-2-yl)-methanol

To a solution of 5-(5-methyl-thiophen-2-ylmethyl)-thiophene-2-carbaldehyde described in Preparation Example 66 (790mg, 3.56mmol) in tetrahydrofuran (5mL) was added lithium aluminum hydride (41mg, 1.06mmol), and the mixture was stirred at room temperature for 10 minutes. Water and ethyl acetate were added to the reaction mixture, which was then partitioned, the organic layer was filtered with a glass filter lined with silica gel, the filtrate was evaporated *in vacuo*, and the title compound (640mg, 2.86mmol, 80.3%) was obtained as a light brown oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.36 (3H, s), 4.20 (2H, s), 4.52 (2H, d, J=6.0Hz), 5.33 (1H, t, J=6.0Hz), 6.56-6.63 (1H, m), 6.66-6.76 (3H, m).

20

#### Preparation Example 68. (5-[1,3]Dioxolan-2-yl-thiophen-2-yl)-(5-methyl-furan-2-yl)-methanol

The title compound (4.2g, 16mmol) was obtained as a reddish brown oil from 2-(5-bromo-thiophen-2-yl)-[1,3]dioxolane (4.0g, 17mmol) and 5-methyl-furan-2-carbaldehyde (1.9g, 17mmol) according to an analogous method to

25



Preparation Examples 65.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.22 (3H, s), 3.89-3.96 (2H, m), 3.98-4.06 (2H, m), 5.83 (1H, d, J=4.8Hz), 5.96-6.02 (2H, m), 6.11-6.13 (1H, m), 6.22-6.24 (1H, m), 6.82-6.84 (1H, m), 7.02-7.05 (1H, m).

5

Preparation Example 69. 5-(5-Methyl-furan-2-ylmethyl)-thiophene-2-carbaldehyde

The title compound (400mg, 1.9mmol, 11.8%) was obtained as a brown oil from 5-[1,3]dioxolan-2-yl-thiophen-2-yl)-(5-methyl-furan-2-yl)-methanol described in Preparation Example 68 (4.2g,16mmol) according to an analogous method to

10 Preparation Example 66.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.21 (3H, s), 4.24 (2H, s), 5.98-6.02 (1H, m), 6.12-6.14 (1H, m), 7.13 (1H, d, J=3.6Hz), 7.89 (1H, d, J=3.6Hz), 9.82 (1H, s).

Preparation Example 70. (5-(5-Methyl-furan-2-ylmethyl)-thiophen-2-yl)-methanol

15 The title compound (210mg, 1.0mmol, 52.6%) was obtained as a brown oil from 5-(5-methyl-furan-2-ylmethyl)-thiophene-2-carbaldehyde described in Preparation Example 69 (400mg, 1.9mmol) according to an analogous method to Preparation Example 67.

20 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.20 (3H, s), 4.05 (2H, s), 4.53 (2H, d, J=5.6Hz), 5.33 (1H, t, J=5.6Hz), 5.94-5.98 (1H, m), 6.02 (1H, d, J=2.8Hz), 6.70-6.77 (2H, m).

Preparation Example 71. Benzofuran-2-yl-(5-[1,3]dioxolan-2-yl-thiophen-2-yl)-methanol

25 The title compound (7.2g, 23.8mmol, 91.5%) was obtained as a yellow oil

from 2-(5-bromo-thiophen-2-yl)-[1,3]dioxolane (6.0g, 26mmol) and benzofuran-2-carbaldehyde (3.8g, 26mmol) according to an analogous method to Preparation Example 65.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.88-4.06 (4H, m), 5.98 (1H, s), 6.08-6.11 (1H, m), 6.59 (1H, d, J=5.2Hz), 6.78 (1H, s), 6.95 (1H, d, 3.6Hz), 7.07 (1H, d, J=3.6Hz), 7.20-7.30 (2H, m), 7.52 (1H, d, J=8.0Hz), 7.61 (1H, d, J=7.2Hz).

Preparation Example 72. (5-Benzofuran-2-ylmethyl-thiophene)-methanol

(Benzofuran-2-ylmethyl)-thiophene-2-carbaldehyde (1.3g, 5.4mmol, 54.5%) was obtained as a brown oil from benzofuran-2-yl-(5-[1,3]dioxolan-2-yl-thiophen-2-yl)-methanol described in Preparation Example 71 (3.0g, 9.9mmol) according to an analogous method to Preparation Example 66. Using this oil (1.2g) according to an analogous method to Preparation Example 67, the title compound (900mg, 3.7mmol, 68.5%) was obtained as a brown oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.32 (2H, s), 4.55 (2H, d, J=5.6Hz), 5.36 (1H, t, J=5.6Hz), 6.67 (1H, s), 6.80 (1H, d, J=3.2Hz), 6.83 (1H, d, J=3.2Hz), 7.17-7.27 (2H, m), 7.50 (1H, d, J=8.0Hz), 7.56 (1H, d, J=7.2Hz).

Preparation Example 73. 1-Phenyl-1*H*-pyrrole-3-carbaldehyde

The title compound (1.2, 7.0mmol, 70%) was obtained as a brown oil from 2,5-dimethoxy-tetrahydrofuran-3-carbaldehyde (2.0g, 12.5mmol) and aniline (930mg, 10mmol) according to an analogous method to Preparation Example 57.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 6.67-6.70 (1H, m), 7.32-7.39 (1H, m), 7.48-7.55 (3H, m), 7.65-7.70 (2H, m), 8.22-8.26 (1H, m), 9.77 (1H, s).

Preparation Example 74. C-(1-Phenyl-1*H*-pyrrol-3-yl)-methanamine

The title compound (580mg, 3.37mmol, 48.1%) was obtained as a light green oil from 1-phenyl-1*H*-pyrrole-3-carbaldehyde described in Preparation Example 73 (1.2g, 7.0mmol) according to an analogous method to Preparation Example 59.

<sup>1</sup>H-NMR Spectrum (DMSO-*d*<sub>6</sub>) δ(ppm) : 3.58 (2H, s), 6.19-6.22 (1H, m), 7.16-7.22 (2H, m), 7.24-7.28 (1H, m), 7.38-7.44 (2H, m), 7.47-7.52 (2H, m).

Preparation Example 75. (3-Phenoxy-benzyl)-carbamic acid phenyl ester

To a solution of phenyl chloroformate (0.29mL, 2.3mmol) in tetrahydrofuran (10mL) were added 3-phenoxybenzylamine described in Preparation Example 4 (0.5g, 2.5mmol) and triethylamine (0.35mL, 2.5mmol) dropwise on an ice bath, and then, the solution was stirred at room temperature for 4 hours. The reaction solution was poured into brine, the solution was extracted with ethyl acetate and concentrated, and the title compound (0.7g, 2.2mmol, 88%) was obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-*d*<sub>6</sub>) δ(ppm) : 4.24 (2H, d, J=6.0 Hz), 6.87-7.49 (14H, m), 8.29 (1H, t, J=6.0 Hz).

Preparation Example 76. 5-(2,4-Difluoro-phenoxy)-furan-2-carbaldehyde

To a solution of 2,4-difluorophenol (6.54mL, 68.0mmol) in dimethylsulfoxide (70mL) was sodium hydride (4534mg, 68.0mmol, 60% in oil), which was then stirred for 40 minutes. 5-Nitro-2-furaldehyde (8000mg, 56.7mmol) was added to the reaction solution, and the solution was stirred at room temperature for 2 hours. Water was added to the reaction mixture, which was then extracted with ethyl acetate. The organic layer was washed with water and brine, dried over anhydrous

magnesium sulfate, and then, concentrated *in vacuo*. The resulting residue was purified by silica gel chromatography (hexane : ethyl acetate), and the title compound (4134mg, 18.44mmol, 33%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 5.54 (1H, d, J=3.6Hz), 6.90-7.03 (2H, m), 7.21  
5 (1H, d, J=3.6Hz), 7.24-7.31 (1H, m), 9.40 (1H, s).

Preparation Example 77. C-(5-(2,4-Difluoro-phenoxy)-furan-2-yl)-methylaniline

To a solution of 5-(2,4-difluoro-phenoxy)-furan-2-carbaldehyde (2060mg, 9.19mmol) in 7N ammonia/methanol solution (100mL) was added Raney nickel  
10 (5.9g), and the mixture was stirred for 24 hours at room temperature under hydrogen atmosphere. The reaction mixture was filtered through Celite pad, and the filtrate was concentrated *in vacuo*. The residue was purified by NH silica gel column chromatography (hexane : ethyl acetate), and the title compound (1814mg, 88.06mmol, 87.7%) was obtained as a yellow oil.

15 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 1.70 (2H, brs), 3.55 (2H, s), 5.55 (1H, d, J=3.2Hz), 6.13 (1H, d, J=3.2Hz), 7.05-7.22 (1H, m), 7.21-7.28(1H, m), 7.43-7.50(1H, m).

Preparation Example 78. 5-(2,5-Difluoro-phenoxy)-furan-2-carbaldehyde

20 To a solution of 2,5-difluorophenol (3360mg, 25.83mmol) in N,N-dimethylformamide (60mL) was sodium hydride (1032mg, 25.83mmol, 60% in oil), which was then stirred for 1 hour. 5-Bromo-2-furaldehyde (3826mg, 21.52mmol) was added to the reaction solution, and the solution was stirred at 60°C for 12 hours. Water was added to the reaction mixture, which was then extracted with ethyl acetate.  
25 The organic layer was washed with water and brine, dried over anhydrous

magnesium sulfate, and then, concentrated *in vacuo*. The resulting residue was purified by silica gel chromatography (hexane-ethyl acetate), and the title compound (1104mg, 4.92mmol, 22.9%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 5.67 (1H, d, J=3.6Hz), 6.93-7.04 (2H, m),  
5 7.15-7.22(1H, m), 7.24 (1H, d, J=3.6Hz), 9.43 (1H, s).

Preparation Example 79. C-(5-(2,5-Difluoro-phenoxy)-furan-2-yl)-methylaniline

The title compound (2353mg, 10.50mmol, 97%) was obtained from 5-(2,5-difluoro-phenoxy)-furan-2-carbaldehyde (2402mg, 11.65mmol) according to an  
10 analogous method to Preparation Example 77.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) :1.72 (2H, brs), 3.56 (2H, s), 5.71 (1H, d, J=3.2Hz), 6.17 (1H, d, J=3.2Hz), 7.01-7.12 (2H, m), 7.41-7.50 (1H, m).

Preparation Example 80. 2-Benzyloxy-thiophene

15 To a solution of benzyl alcohol (3.45mL, 33.3mmol) in 1,2-dimethoxyethane (80mL) was added *n*-butyl lithium (2.6M hexane solution, 13.5mL, 33.3mmol) dropwise, which was then stirred for 10 minutes. Copper(I) chloride (5210mg, 49.45mmol) was added thereto, followed by stirring for 10 minutes on an ice bath, then, it was stirred for 2.5 hours at room temperature. 2-Iodothiophene (4995mg,  
20 23.78mmol) and pyridine 320mL were further added, and the solution was stirred for 13 hours under reflux. Ethyl acetate was added to the reaction mixture, which was sequentially washed with 1N hydrochloric acid, an aqueous solution of sodium hydrogen sulfate and brine, dried over anhydrous magnesium sulfate, and then, concentrated *in vacuo*. The resulting residue was purified by silica gel  
25 chromatography (hexane-ethyl acetate), and the title compound (420mg, 2.21mmol,

9.5%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 5.08 (2H, s), 6.28 (1H, dd, J=1.6, 4.0Hz), 6.57 (1H, dd, J=1.6, 5.6Hz), 6.71 (1H, dd, J=4.0, 5.6Hz), 7.30-7.47 (5H, m).

5 Preparation Example 81. 5-Benzyloxy-thiophene-2-carbonitrile

To a solution of 2-benzyloxy-thiophene (184mg, 0.967mmol) in diethyl ether (4mL) was added *n*-butyl lithium (2.47M hexane solution, 0.47mL, 1.16mmol) at -78°C under nitrogen atmosphere, then, the solution was stirred for 1.5 hours on an ice bath. The solution was cooled again to -78°C, N,N-dimethylformamide (487μl, 4.84mmol) was added, and the solution was stirred for 45 minutes while warming to room temperature. An aqueous solution of saturated ammonium chloride was added to the reaction mixture, which was then extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, then, filtered with silica gel, the filtrate was concentrated *in vacuo*, and 5-benzyloxy-thiophene-2-carbaldehyde (171mg) was obtained.

The resulting 5-benzyloxy-thiophene-2-carbaldehyde (171mg) was dissolved in N,N-dimethylformamide (4mL), pyridine (82μl, 1.02mmol) and hydroxylamine hydrochloride (65mg, 0.94mmol) were added thereto, followed by stirring for 30 minutes at 60°C, then, cooled on an ice bath. 1,1'-Carbonyldiimidazole (635mg, 3.92mmol) was added, the solution was warmed again to 60°C and stirred for 35 minutes, triethylamine (272μl, 1.96mmol) was added, and the solution was further stirred for 30 minutes. The reaction mixture was cooled to room temperature, water was added, and the solution was extracted with ethyl acetate. The organic layer was sequentially washed with an aqueous solution of oxalic acid, an aqueous solution of saturated sodium bicarbonate and brine, dried

over anhydrous sodium sulfate, and then, concentrated *in vacuo*. The resulting residue was purified by silica gel chromatography (hexane-ethyl acetate), and the title compound (30mg, 0.14mmol, 14%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 5.15 (2H, s), 6.28 (1H, d, J=4.0Hz), 7.32 (1H, d, J=4.0Hz), 7.38-7.48 (5H, m).

Preparation Example 82. 2-(5-(3-Fluoro-benzyl)-furan-2-yl)-[1,3]dioxolane

*n*-Butyl lithium (2.66M hexane solution, 25.5mL, 67.88mmol) was added dropwise to a solution of 2-(1,3-dioxolan-2-yl)furan (8272mg, 59.03mmol) in tetrahydrofuran (160mL) that had been cooled to -78°C solution under nitrogen atmosphere, which was then stirred for 10 minutes. To this solution was added a solution of 3-fluorobenzyl bromide (14.50g, 76.73mmol) in tetrahydrofuran (60mL) dropwise, which was then stirred for 1 hour at -78°C, and for 1.25 hours at room temperature. An aqueous solution of saturated ammonium chloride was added to the reaction mixture, which was extracted with diethyl ether. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and then, concentrated *in vacuo*. The obtained residue was purified by silica gel chromatography (hexane-ethyl acetate), and a mixture of the title compound and 2-(1,3-dioxolane-2-yl)furan (8033mg), which is the starting material, was obtained.

Preparation Example 83. 5-(3-Fluoro-benzyl)-furan-2-carbaldehyde

To a solution of a mixture of 2-(5-(3-fluoro-benzyl)-furan-2-yl)-[1,3]dioxolane and 2-(1,3-dioxolan-2-yl)furan (8033mg) in methanol (80mL) was added an aqueous solution of oxalic acid (22g, 115mmol) (80mL), and the solution was stirred at room temperature for 1 hour. Water was added to the reaction mixture, which extracted

with diethyl ether, the organic layer was sequentially washed with an aqueous solution of saturated sodium bicarbonate, water and brine, dried over anhydrous sodium sulfate, and then, concentrated *in vacuo*. The resulting residue was purified by silica gel chromatography (hexane-ethyl acetate), and the title compound (4084mg, 20.0mmol, 34%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) :4.07 (2H, s), 6.23 (1H, d, J=4.0Hz), 6.90-7.10 (3H, m), 7.18 (1H, d, J=4.0Hz), 7.25-7.37 (1H, m), 9.56 (1H, s).

Preparation Example 84. C-(5-(3-Fluoro-benzyl)-furan-2-yl)-methylamine

The title compound (4104mg, 20.0mmol, 100%) was obtained from 5-(3-fluoro-benzyl)-furan-2-carbaldehyde (4084mg, 20.0mmol) according to an analogous method to Preparation Example 77.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) :1.60 (2H, brs), 3.31 (2H, s), 3.93 (2H, s), 5.97-6.11 (2H, m), 6.82-7.15 (3H, m), 7.20-7.41 (1H, m).

Preparation Example 85. 2-(3-[1,2,3] Triazol-2-yl-propyl)-isoindol-1,3-dione

To a solution of 1H-1,2,3-triazole (2000mg, 28.96mmol) in N,N-dimethylformamide (60mL) was added sodium hydride (1159mg, 28.96mmol, 60% in oil), which was stirred for 30 minutes. N-(3-Bromopropyl)phthalimide (7057mg, 26.32mmol) and potassium iodide (431mg, 2.63mmol) were added thereto, and the solution was stirred for 3 hours at 70°. Water was added to the reaction mixture, which was then extracted with ethyl acetate. The organic layer was washed with water and brine, dried over anhydrous magnesium sulfate, and then, concentrated *in vacuo*. The resulting residue was purified by silica gel chromatography (dichloromethane-ethyl acetate), and the title compound (3526mg,



13.75mmol, 47%) was obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.19-2.28 (2H, m), 3.65 (2H, t, J=6.8Hz), 4.51 (2H, t, J=6.8Hz), 7.74 (2H, s), 7.80-7.89 (4H, m).

5 Preparation Example 86. 3-[1,2,3] Triazol-2-yl-propyl amine

To a mixture solution of 2-(3-[1,2,3]triazol-2-yl-propyl)-isoindol-1,3-dione (1782mg, 6.95mmol) in methanol-tetrahydrofuran (5:4, 27mL) was added hydrazine monohydrate (371μl, 7.65mmol), and the solution was stirred for 5 days at room temperature. Methanol (8mL) was added thereto, followed by further stirring for  
10 3.25 hours under reflux. The reaction mixture was filtered, and the filtrate was concentrated. The resulting residue was dissolved in tetrahydrofuran, adsorbed onto NH silica gel, purified by NH silica gel chromatography (ethyl acetate-methanol), and the title compound (491mg, 1.36mmol, 19.6%) was obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) :1.85-1.93 (2H, m), 2.46-2.51 (2H, m),  
15 4.42-4.52 (2H, m), 7.75 (2H, s).

Preparation Example 87. 3-Benzyl amino-benzonitrile

3-Bromo-benzonitrile (500mg, 2.75mmol), benzylamine (360μl, 3.30mmol), 2,2-bis(diphenylphosphino)-1,1'-binaphthyl (8.6mg, 14μmol),  
20 tris(dibenzylideneacetone)dipalladium(0) (19mg, 21μmol) and sodium *tert*-butoxide (370mg, 3.85mmol) were suspended in toluene (10mL), and the mixture was stirred at 80°C for 22 hours under nitrogen atmosphere. The reaction mixture was filtered through Celite pad, then, the filtrate was evaporated *in vacuo*, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 15 : 1),  
25 and the title compound (331mg, 1.59mmol, 58%) was obtained as a pale yellow oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.33 (2H, m), 4.34 (1H, s), 6.79-6.83 (2H, m), 6.97 (1H, dt, J=1.2, 7.7Hz), 7.22 (1H, t, J=8.1Hz), 7.29-7.39 (5H, m).

Preparation Example 88. 4-Phenylamino-benzonitrile

5           The title compound (460mg, 2.37mmol, 86%) was obtained as a white solid from 4-bromo-benzonitrile (500mg, 2.75mmol) and phenylamine (300μl, 3.30mmol) according to an analogous method to Preparation Example 87.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>)δ(ppm) : 6.04 (1H, s), 6.97 (2H, d, J=8.8Hz), 7.12 (1H, t, J=7.3Hz), 7.17 (2H, d, J=7.7Hz), 7.36 (2H, t, J=7.5Hz), 7.48 (2H, d, J=8.8Hz).

10

Preparation Example 89. 4-Benzylamino-benzonitrile

          The title compound (472mg, 2.27mmol, 83%) was obtained as a pale yellow solid from 4-bromo-benzonitrile (500mg, 2.75mmol) and benzylamine (360μl, 3.30mmol) according to an analogous method to Preparation Example 87.

15       <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>)δ(ppm) : 4.38 (2H, d, J=5.5Hz), 4.58 (1H, s), 6.59 (2H, d, J=8.8Hz), 7.29-7.39 (5H, m), 7.42 (2H, d, J=8.8Hz).

Preparation Example 90. 2-(3-Bromo-phenyl)-[1,3]dioxolane

          3-Bromobenzaldehyde (4.00g, 21.6mmol), ethane-1,2-diol (6.03mL, 20   108mmol) and toluene-4-sulfonic acid monohydrate (186mg,1.08mmol) were dissolved in toluene (80mL), and the solution was stirred under reflux for 4 hours. An aqueous solution of saturated sodium bicarbonate was added to the reaction solution at 0°C, which was then extracted with ethyl acetate, the organic layer was washed with brine, and the organic layer was dried over anhydrous magnesium 25   sulfate. The solvent was evaporated *in vacuo*, the residue was purified by NH silica

gel column chromatography (hexane : ethyl acetate = 20 : 1), and the title compound (4.79g, 20.9mmol, 97%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.02-4.05 (2H, m), 4.07-4.13 (2H, m), 5.79 (1H, s), 7.23-7.27 (1H, m), 7.40 (1H, d, J=7.7Hz), 7.49 (1H, dt, J=1.1, 7.1Hz), 7.64 (1H, s).

5

Preparation Example 91. 2-(3-Phenylsulfanyl-phenyl)-[1,3]dioxolane

2-(3-Bromo-phenyl)-[1,3]dioxolane described in Preparation Example 90 (515mg, 2.25mmol) was dissolved in tetrahydrofuran (10mL) under nitrogen atmosphere, *n*-butyl lithium (2.47M hexane solution, 1.64mL, 4.05mmol) was added at 10 -78°C, the solution was stirred for 15 minutes, then, diphenyldisulphide (540mg, 2.48mmol) was added thereto, followed by stirring for 3 hours. The reaction solution was cooled to 0°C, water was added, the solution was extracted with ethyl acetate, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by silica gel 15 column chromatography (hexane : ethyl acetate = 20 : 1), and the title compound (402mg, 1.56mmol, 69%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.01-4.06 (2H, m), 4.09-4.14 (2H, m), 5.76 (1H, s), 7.22-7.39 (8H, m), 7.48 (1H, s).

20 Preparation Example 92. 3-Phenylsulfanyl-benzaldehyde

2-(3-Phenylsulfanyl-phenyl)-[1,3]dioxolane described in Preparation Example 91 (396mg, 1.53mmol) was dissolved in a mixture solution of ethanol (5mL), water (5mL), tetrahydrofuran (5mL) and sulfuric acid (1mL), and the solution was stirred for 2.5 hours under reflux. The reaction solution was cooled to 0°C, an aqueous 25 solution of saturated sodium bicarbonate was added thereto, the solution was

extracted with ethyl acetate, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 20 : 1), and the title compound (323mg, 1.51mmol, 98%) was obtained as a colorless oil.

5 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 7.31-7.39 (3H, m), 7.41-7.50 (3H, m), 7.52 (1H, d, J=7.9Hz), 7.71 (1H, d, J=7.3Hz), 7.76 (1H, s), 9.94 (1H, s).

Preparation Example 93. (3-Phenylsulfanyl-phenyl)-methanol

3-Phenylsulfanyl-benzaldehyde described in Preparation Example 92 (321mg, 1.49mmol) was dissolved in ethanol (6mL), sodium borohydride (113mg, 2.98mmol) was added thereto at 0°C, followed by stirring at room temperature for 3 hours. Water was added to the reaction solution, which was then extracted with ethyl acetate, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 5 : 1), and the title compound (220mg, 1.02mmol, 68%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.73 (2H, s), 4.66 (1H, s), 7.18-7.37 (9H, m).

Preparation Example 94. 2-(3-Phenylsulfanyl-benzyl)-isoindol-1,3-dione

20 (3-Phenylsulfanyl-phenyl)-methanol described in Preparation Example 93 (212mg, 0.980mmol), phthalimide (144mg, 0.980mmol), diethylazodicarboxylate (170μl, 1.08mmol) and triphenylphosphine (308mg, 1.18mmol) were dissolved in tetrahydrofuran (4mL), and the solution was stirred overnight at room temperature. Water was added to the reaction solution, which was then extracted with ethyl acetate, the organic layer was washed with brine and dried over anhydrous

25

magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 10 : 1), and the title compound (124mg, 0.359mmol, 37%) was obtained as a solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.80 (2H, s), 7.17-7.36 (9H, m), 7.73 (2H, dd, J=2.9, 5.3Hz), 7.86 (2H, dd, J=2.9, 5.3Hz).

#### Preparation Example 95. 3-Phenylsulfanyl-benzylamine

2-(3-Phenylsulfanyl-benzyl)-isoindole-1,3-dione described in Preparation Example 94 (123mg, 0.356mmol) was dissolved in ethanol (3mL), hydrazine monohydrate (518μl, 10.7mmol) was added at 0°C, and the solution was stirred under reflux for 2 hours. Water was added to the reaction solution at 0°C, which was then extracted with ethyl acetate, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by NH silica gel column chromatography (ethyl acetate), and the title compound (75mg, 0.35mmol, 98%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.83 (2H, s), 7.19-7.36 (9H, m).

#### Preparation Example 96. 2-(4-Bromo-phenyl)-[1,3]dioxolane

4-Bromo-benzaldehyde (4.00g, 21.6mmol), ethane-1,2-diol (6.03mL, 108mmol) and toluene-4-sulfonic acid monohydrate (186mg, 1.08mmol) were dissolved in toluene (80mL), and the solution was stirred under reflux for 4 hours. An aqueous solution of saturated sodium bicarbonate was added to the reaction solution at 0°C, which was then extracted with ethyl acetate, the organic layer was washed with brine, and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by NH silica

gel column chromatography (hexane : ethyl acetate = 20 : 1), and the title compound (4.66g, 20.3mmol, 94%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.01-4.05 (2H, m), 4.07-4.13 (2H, m), 5.77 (1H, s), 7.35 (2H, d, J=8.2Hz), 7.51 (2H, d, J=8.4Hz).

5

Preparation Example 97. 2-(4-Benzylsulfanyl-phenyl)-[1,3]dioxolane

The title compound (568mg, 2.09mmol, 48%) was obtained as a solid from 2-(4-bromo-phenyl)-[1,3]dioxolane described in Preparation Example 96 (1.00g, 4.37mmol) and benzyl disulphide (1.18g, 4.81mmol) according to an analogous method to Preparation Example 91.

10

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.02-4.04 (2H, m), 4.10-4.13 (2H, m), 4.13 (2H, s), 5.76 (1H, s), 7.23-7.27 (1H, m), 7.28-7.32 (6H, m), 7.37 (2H, d, J=8.2Hz).

Preparation Example 98. 4-Benzylsulfanyl-benzaldehyde

The title compound (462mg, 2.02mmol, 97%) was obtained as a white solid from 2-(4-benzyl sulfanyl-phenyl)-[1,3]dioxolane described in Preparation Example 97 (568mg, 2.09mmol) according to an analogous method to Preparation Example 92.

15

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.24 (2H, s), 7.26-7.40 (7H, m), 7.75 (2H, d, J=8.6Hz), 9.92 (1H, s).

20

Preparation Example 99. (4-Benzylsulfanyl-phenyl)-methanol

The title compound (406mg, 1.76mmol, 87%) was obtained as a white solid from 4-benzylsulfanyl-benzaldehyde described in Preparation Example 98 (462mg, 2.02mmol) according to an analogous method to Preparation Example 93.

25

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.11 (2H, s), 4.65 (2H, d, J=4.4Hz), 7.20-7.35

(8H, m), 7.37 (1H, d, J=4.4Hz).

Preparation Example 100. 2-(4-Benzylsulfanyl-benzyl)-isoindol-1,3-dione

The title compound (563mg, 1.57mmol, 89%) was obtained as a white solid  
5 from (4-benzylsulfanyl)-methanol described in Preparation Example 99 (406mg, 1.76mmol) according to an analogous method to Preparation Example 94.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.09 (2H, s), 4.79 (2H, s), 7.20-7.35 (5H, m), 7.24 (2H, d, J=8.2Hz), 7.32 (2H, d, J=8.1Hz), 7.71 (2H, dd, J=2.9, 5.3Hz), 7.84 (2H, dd, J=2.9, 5.3Hz).

10

Preparation Example 101. 4-Benzylsulfanyl-benzylamine

The title compound (260mg, 1.13mmol, 72%) was obtained as a white solid  
from 2-(4-benzylsulfanyl-benzyl)-isoindol-1,3-dione described in Preparation Example  
100 (563mg, 1.57mmol) according to an analogous method to Preparation Example  
15 95.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.83 (2H, s), 4.10 (2H, s), 7.20 (2H, d, J=8.6Hz), 7.19-7.32 (7H, m).

Preparation Example 102. (5-Phenylaminomethyl-furan-2-yl)-methanol

20 Acetic acid 5-formyl-furan-2-ylmethyl ester (2.00g, 11.9mmol), aniline (1.63mL, 17.9mmol) and triacetoxy sodium borohydride (5.04g, 23.8mmol) were suspended in a mixture solution of tetrahydrofuran (40mL) and acetic acid (1mL) at 0°C, and the mixture was stirred at room temperature for 7 hours. An aqueous solution of saturated sodium bicarbonate was added to the reaction solution at 0°C,  
25 which was then extracted with ethyl acetate and tetrahydrofuran, the organic layer

was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 1 : 1), and a mixture of acetic acid 5-phenylaminomethyl-furan-2-ylmethyl ester and aniline (2.91g) was obtained as a  
 5 pale yellow oil.

Then, the resulting mixture of acetic acid 5-phenylaminomethyl-furan-2-ylmethyl ester and aniline (2.91g) as well as potassium carbonate (3.28g, 23.7mmol) were suspended in methanol (60mL), and the solution was stirred overnight at room temperature. The reaction solution was evaporated *in*  
 10 *vacuo*, water and ethyl acetate were added to the residue, the organic layer was partitioned, washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 1 : 1), and the title compound (1.99g, 9.79mmol, 82%) was obtained as a colorless oil.

15 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.04 (1H, brs), 4.31 (2H, s), 4.59 (2H, d, J=5.9Hz), 6.19 (1H, d, J=3.1Hz), 6.23 (1H, d, J=3.1Hz), 6.68 (2H, d, J=7.5Hz), 6.75 (1H, t, J=7.3Hz), 7.19 (2H, t, J=7.3Hz).

Preparation Example 103. 2-(5-phenylaminomethyl-furan-2-ylmethyl)-

20 isoindol-1,3-dione

The title compound (603mg, 1.81mmol, 23%) was obtained as a pale yellow solid from (5-phenylaminomethyl-furan-2-yl)-methanol described in Preparation Example 102 (1.58g, 7.77mmol) according to an analogous method to Preparation Example 94.

25 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.25 (2H, s), 4.82 (2H, s), 6.14 (1H, d, J=3.1Hz),



6.28 (1H, d, J=3.1Hz), 6.63 (2H, d, J=7.5Hz), 6.70 (1H, t, J=7.3Hz), 7.14 (2H, t, J=7.3Hz), 7.72 (2H, dd, J=3.1,5.3Hz), 7.87 (2H, dd, J=3.1,5.3Hz).

Preparation Example 104. (5-Aminomethyl-furan-2-ylmethyl)-phenyl-amine

5           The title compound (92mg, 0.46mmol, 60%) was obtained as a pale yellow oil from 2-(5-phenylaminomethyl-furan-2-ylmethyl)-isoindol-1,3-dione described in Preparation Example 103 (251mg, 0.755mmol) according to an analogous method to Preparation Example 95.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.79 (2H, s), 4.28 (2H, s), 6.06 (1H, d, J=3.1Hz),  
10   6.15 (1H, d, J=3.1Hz), 6.68 (2H, d, J=7.7Hz), 6.74 (1H, t, J=7.3Hz), 7.19 (2H, t, J=7.3Hz).

Preparation Example 105. (2-(5-[1,3]Dioxolan-2-yl-furan-2-yl)-ethyl)-phenylamine

5-[1,3]Dioxolan-2-yl-furan-2-carbaldehyde           (2.03g,           12.1mmol),  
15   trimethylsulfonium bromide (1.90g, 12.1mmol) and potassium hydroxide (779mg, 13.9mmol) were suspended in acetonitrile (75mL), and the solution was stirred overnight at room temperature. Water was added to the reaction solution, which was then extracted with ethyl acetate, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*,  
20   2-(5-oxiranyl-furan-2-yl)-[1,3]dioxolane (2.25g) was obtained as a pale yellow oil.

          The resulting 2-(5-oxiranyl-furan-2-yl)-[1,3]dioxolane (2.25g) and silica gel (5.00g) were suspended in ethyl acetate (40mL), and the solution was stirred for 6.5 hours at room temperature. The reaction solution was filtered, then, the filtrate was evaporated *in vacuo*, and (5-[1,3]dioxolan-2-yl-furan-2-yl)-acetaldehyde (1.57g) was  
25   obtained as a yellow oil.

Next, the resulting (5-[1,3]dioxolan-2-yl-furan-2-yl)-acetaldehyde (1.57g), aniline (0.94mL, 10.3mmol) and triacetoxysodium borohydride (3.76g, 17.2mmol) were suspended in a mixture solution of tetrahydrofuran (30mL) and acetic acid (1mL) at 0°C, and the mixture was stirred at room temperature for 19 hours. An aqueous solution of saturated sodium bicarbonate was added to the reaction mixture at 0°C, which was then extracted with ethyl acetate, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 5 : 1), and the title compound (453mg, 1.75mmol, 14%) was obtained as a brown oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 2.95 (2H, t, J=6.8Hz), 3.43 (2H, t, J=6.8Hz), 3.99-4.06 (2H, m), 4.10-4.15 (2H, m), 5.88 (1H, s), 6.06 (1H, d, J=3.3Hz), 6.37 (1H, d, J=3.1Hz), 6.62 (2H, dd, J=1.1, 8.6Hz), 6.71 (1H, tt, J=1.1, 7.3Hz), 7.18 (2H, dd, J=7.3, 8.6Hz).

#### Preparation Example 106. 5-(2-Phenylamino-ethyl)-furan-2-carbaldehyde

The title compound (314mg, 1.46mmol) was obtained as a light brown oil from (2-(5-[1,3]dioxolan-2-yl-furan-2-yl)-ethyl)-phenylamine described in Preparation Example 105 (453mg, 1.75mmol) according to an analogous method to Preparation Example 44.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.05 (2H, t, J=6.8Hz), 3.53 (2H, t, J=6.8Hz), 6.33 (1H, d, J=3.5Hz), 6.62 (2H, dd, J=1.1, 8.6Hz), 6.73 (1H, t, J=7.3Hz), 7.17-7.21 (3H, m), 9.55 (1H, s).

#### Preparation Example 107. (2-(5-Aminomethyl-furan-2-yl)-ethyl)-phenylamine

The title compound (117mg,0.541mmol,78%) was obtained as a pale yellow oil from 5-(2-phenylamino-ethyl)-furan-2-carbaldehyde described in Preparation Example 106 (150mg,0.697mmol) according to an analogous method to Preparation Example 45.

5 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 2.92 (2H, t, J=6.8Hz), 3.41 (2H, t, J=6.8Hz), 3.78 (2H, s), 6.00 (1H, d, J=3.1Hz), 6.04 (1H, d, J=2.9Hz), 6.62 (2H, dd, J=1.1, 8.6Hz), 6.71 (1H, t, J=7.3Hz), 7.18 (2H, dd, J=7.3, 8.6Hz).

Preparation Example 108. 2-(4-Bromo-thiophen-2-yl)-[1,3]dioxolane

10 4-Bromo-thiophene-2-carbaldehyde (9.24g,48.4mmol), ethane-1,2-diol (13.5mL,242mmol), toluene-4-sulfonic acid monohydrate (416mg,2.42mmol) were dissolved in toluene (100mL), and the solution was stirred for 1.5 hours under reflux. An aqueous solution of saturated sodium bicarbonate was added to the reaction solution at 0°C, which was then extracted with ethyl acetate, the organic layer was  
15 washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 20 : 1), and the title compound (11.8g, quantitatively) was obtained as a white solid.

20 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.02-4.04 (2H, m), 4.09-4.11 (2H, m), 6.07 (1H, s), 7.08 (1H, dd, J=0.73, 1.5Hz), 7.22 (1H, d, J=1.5Hz).

Preparation Method 109. 2-(4-Phenoxy-thiophen-2-yl)-[1,3]dioxolane

The title compound (5.40g, 21.7mmol, 73%) was obtained as a pale yellow oil from 2-(4-bromo-thiophen-2-yl)-[1,3]dioxolane described in Preparation Example 108  
25 (6.96g, 29.6mmol) and phenol (6.60g, 71.0mmol) according to an analogous method

to Preparation Example 33.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.00-4.04 (2H, m), 4.12-4.17 (2H, m), 6.04 (1H, s), 6.60 (1H, d, J=1.7Hz), 6.94 (1H, d, J=1.7Hz), 7.04 (2H, dd, J=1.1, 8.6Hz), 7.09 (1H, tt, J=1.1, 7.3Hz), 7.32 (2H, dd, J=7.3, 8.6Hz).

5

Preparation Example 110. 4-Phenoxy-thiophene-2-carbaldehyde

The title compound (183mg, 0.896mmol, 44%) was obtained as a colorless oil from 2-(4-phenoxy-thiophen-2-yl)-[1,3]dioxolane described in Preparation Example 109 (500mg, 2.01mmol) according to an analogous method to Preparation Example

10 34.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 7.04 (1H, dd, J=1.3, 1.6Hz), 7.07 (2H, dd, J=1.1, 8.8Hz), 7.16 (1H, tt, J=1.1, 7.3Hz), 7.35 (2H, dd, J=7.3, 8.6Hz), 7.51 (1H, d, J=1.7Hz), 9.84 (1H, s).

15 Preparation Example 111. C-(4-Phenoxy-thiophen-2-yl)-methylamine

The title compound (94mg, 0.458mmol, 51%) was obtained as a pale yellow oil from 4-phenoxy-thiophene-2-carbaldehyde described in Preparation Example 110 (183mg, 0.896mmol) according to an analogous method to Preparation Example 35.

20 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.99 (2H, s), 6.46 (1H, d, J=1.7Hz), 6.69-6.70 (1H, m), 7.05 (2H, dd, J=1.1, 7.7Hz), 7.09 (1H, t, J=7.5Hz), 7.33 (2H, dd, J=7.5, 8.6Hz).

Preparation Example 112. 5-Oxo-2,5-dihydro-isoxazole-4-carboxylic acid ethyl ester

25 2-Ethoxymethylene-malonic acid diethyl ester (5.00g, 23.1mmol), hydroxylamine hydrochloride (4.01g, 57.8mmol) and triethylamine (8.06mL,

57.8mmol) were dissolved in ethanol (100mL), the solution was stirred at room temperature for 17 hours, then, the solution was stirred for 4.5 hours under reflux. The reaction solution was cooled to room temperature, water was added, and 1N hydrochloric acid was added until a salt precipitated. The precipitated salt was collected by filtration, and a hydrochloride salt of the title compound (2.39g, 15.2mmol, 66%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) :1.19 (3H, t, J=7.1Hz), 4.08 (2H, q, J=7.1Hz), 4.90 (1H, brs), 8.51 (1H, d, J=6.8Hz).

10 Preparation Example 113. 5-Oxo-2-phenoxy thiocarbonyl-2,5-dihydro-isoxazole-4-carboxylic acid ethyl ester

5-Oxo-2,5-dihydro-isoxazole-4-carboxylic acid ethyl ester described in Preparation Example 112 (1.00g, 5.17mmol), phenyl chlorothionoformate (786μl, 5.69mmol) and pyridine (919μl, 11.4mmol) were dissolved at 0°C in toluene (20mL), and the solution was stirred at room temperature for 17 hours under nitrogen atmosphere. Water was added to the reaction solution at 0°C, which was then extracted with ethyl acetate, and the organic layer was washed with brine. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 1 : 1), and the title compound (1.66g, 5.66mmol, quantitatively) was obtained as a pale yellow solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) :1.39 (3H, t, J=7.1Hz), 4.39 (2H, q, J=7.1Hz), 7.16 (2H, dd, J=1.3, 8.6Hz), 7.38 (1H, t, J=7.1Hz), 7.49 (2H, dd, J=7.1, 8.8Hz), 9.30 (1H, s).

25 Preparation Example 114. 2-Phenoxy-thiazole-5-carboxylic acid ethyl ester

5-Oxo-2-phenoxythiocarbonyl-2,5-dihydro-isoxazole-4-carboxylic acid ethyl ester described in Preparation Example 113 (500mg, 2.01mmol) was dissolved in acetone (500mL), and the solution was irradiated with light (300nm) for 30 minutes at room temperature. The reaction solution was evaporated *in vacuo*, the residue was  
5 purified by silica gel column chromatography (hexane : ethyl acetate = 2 : 1), and the title compound (384mg, 1.54mmol, 90%) was obtained as a pale yellow oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 1.35 (3H, t, J=7.1Hz), 4.33 (2H, q, J=7.1Hz), 7.26-7.35 (3H, m), 7.47 (2H, dd, J=7.5, 8.4Hz), 7.90 (1H, s).

10 Preparation Example 115. (2-Phenoxy-thiazol-5-yl)-methanol

2-Phenoxy-thiazole-5-carboxylic acid ethyl ester described in Preparation Example 114 (384mg, 1.54mmol) was dissolved in tetrahydrofuran (5mL), and lithium aluminum hydride (292mg, 7.70mmol) was added at 0°C. The solution was stirred at room temperature for 1 hour, then, at 0°C, water (292μl), an aqueous solution of  
15 5N sodium hydroxide (292μl) and water (876μl) were sequentially added thereto. The reaction solution was filtered through Celite pad, then, the filtrate was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 1 : 4), and the title compound (270mg, 1.30mmol, 85%) was obtained as a white solid.

20 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.73 (2H, d, J=5.9Hz), 7.13 (1H, s), 7.25-7.29 (3H, m), 7.41-7.45 (2H, m).

Preparation Example 116. 2-(2-Phenoxy-thiazol-5-ylmethyl)-isoindol-1,3-dione

The title compound (131mg, 0.389mmol, 30%) was obtained as a colorless  
25 oil from (2-phenoxy-thiazole-5-yl)-methanol described in Preparation Example 115

(270mg, 1.30mmol) according to an analogous method to Preparation Example 94.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.89 (2H, s), 7.21-7.28 (4H, m), 7.40 (2H, t, J=8.0Hz), 7.73 (2H, dd, J=3.1, 5.3Hz), 7.86 (2H, dd, J=2.9, 5.5Hz).

5 Preparation Example 117. C-(2-Phenoxy-thiazol-5-yl)-methylamine

The title compound (63mg, 0.31mmol, 78%) was obtained as a colorless oil from 2-(2-phenoxy-thiazol-5-ylmethyl)-isoindol-1,3-dione described in Preparation Example 116 (131mg, 0.389mmol) according to an analogous method to Preparation Example 95.

10 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.95 (2H, s), 7.03 (1H, t, J=1.1Hz), 7.25-7.28 (3H, m), 7.39-7.43 (2H, m).

Preparation Example 118. 4-Benzyloxy-2-fluoro-benzonitrile

To a solution of 4-hydroxy-2-fluoro-benzonitrile (1.0g, 7.3mmol) in  
15 N,N-dimethylformamide (15mL) was added potassium carbonate (2.0g, 15mmol) and benzyl bromide (0.87mL, 7.3mmol), and the solution was stirred at room temperature for 5 hours. Water and ethyl acetate were added to the reaction solution, which was then extracted, washed with brine, then, the solvent was evaporated *in vacuo*. The residue was purified by silica gel column chromatography (hexane : ethyl acetate =  
20 5 : 1), and the title compound (1.5g, 6.7mmol, 92%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 5.11 (2H, s), 6.78 (1H, dd, J=2.4, 11.0Hz), 6.83 (1H, ddd, J=0.6, 2.4, 8.8Hz), 7.37-7.43 (5H, m), 7.52 (1H, dd, J=7.5, 8.6Hz).

Preparation Example 119. 4-[1,3]Dioxolane-2-yl-benzonitrile

25 4-Formyl-benzonitrile (3.00g, 22.9mmol), ethane-1,2-diol (6.38mL, 115mmol)

and toluene-4-sulfonic acid monohydrate (197mg, 1.15mmol) were dissolved in toluene (60mL), and the solution was stirred under reflux for 10 hours. An aqueous solution of saturated sodium bicarbonate was added to the reaction solution at 0°C, which was then extracted with ethyl acetate, the organic layer was washed with brine, and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 10 : 1), and the title compound (3.78g, 21.6mmol, 94%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.04-4.13 (4H, m), 5.85 (1H, s), 7.59 (2H, d, J=8.1Hz), 7.68 (2H, d, J=8.4Hz).

#### Preparation Example 120. 4-[1,3]Dioxolan-2-yl-benzylamine

4-[1,3]Dioxolan-2-yl-benzonitrile described in Preparation Example 119 (3.78g, 21.6mmol) was dissolved in tetrahydrofuran (76mL), lithium aluminum hydride (4.09g, 108mmol) was added at 0°C, and the solution was stirred overnight at room temperature. Water (4.09mL), an aqueous solution of 5N sodium hydroxide (4.09mL) and water (12.3mL) were sequentially added to the reaction solution. The reaction solution was filtered through Celite pad, then, the filtrate was evaporated *in vacuo*, and the title compound (3.92g, quantitatively) was obtained as a pale yellow oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.89 (2H, s), 4.03-4.16 (4H, m), 5.81 (1H, s), 7.34 (2H, d, J=7.9Hz), 7.46 (2H, d, J=8.1Hz).

#### Preparation Example 121. 5-(3-Chloro-phenoxy)-thiophene-2-carbonitrile

5-Nitro-thiophene-2-carbonitrile (5.00g, 32.5mmol), 3-chloro-phenol (6.90mL,



65.0mmol) and potassium carbonate (13.4g, 97.5mmol) were suspended in dimethylsulfoxide (50mL), and the mixture was stirred at 60°C for 4 hours. Water was added to the reaction mixture at 0°C, which was then extracted with ethyl acetate, the organic layer was washed with water twice, and further washed with  
5 brine. The organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by NH silica gel silica gel column chromatography (hexane : ethyl acetate = 20 : 1), and the title compound (5.56g, 23.6mmol, 73%) was obtained as a pale yellow oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 6.49 (1H, d, J=4.2Hz), 7.04 (1H, ddd, J=0.92,  
10 2.4, 8.2Hz), 7.15 (1H, t, J=2.2Hz), 7.22 (1H, ddd, J=0.92, 2.0, 8.1Hz), 7.33 (1H, t, J=8.2Hz), 7.4 (1H, d, J=4.2Hz).

Preparation Example 122. 2-(5-(2-Fluoro-benzyl)-thiophen-2-yl)-[1,3]dioxolane

The title compound (4.33g, 16.4mmol, 54%) was obtained as a pale yellow oil  
15 from 2-(5-bromo-thiophen-2-yl)-[1,3]dioxolane (8.00g, 30.4mmol) and 1-bromo methyl-2-fluoro-benzene (4.48mL, 36.5mmol) according to an analogous method to Preparation Example 36.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.97-4.03 (2H, m), 4.06-4.13 (2H, m), 4.14 (2H, s), 6.01 (1H, s), 6.71 (1H, d, J=3.7Hz), 6.98 (1H, d, J=3.9Hz), 7.01-7.08 (2H, m),  
20 7.19-7.23 (2H, m).

Preparation Example 123. 5-(2-Fluoro-benzyl)-thiophene-2-carbaldehyde

2-(5-(2-Fluoro-benzyl)-thiophen-2-yl)-[1,3]dioxolane described in Preparation Example 122 (4.33g, 16.4mmol) was dissolved in a mixture solvent of methanol  
25 (40mL) and water (10mL), 1N hydrochloric acid (20mL) was added thereto, followed

by stirring at room temperature for 1 hour. An aqueous solution of saturated sodium bicarbonate was added to the reaction solution at 0°C, which was then extracted with ethyl acetate, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified  
 5 by silica gel column chromatography (hexane : ethyl acetate = 10 : 1), and the title compound (3.54g, 16.1mmol, 98%) was obtained as a light yellow oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.22 (2H, s), 6.94 (1H, d, J=3.8Hz), 7.05-7.13 (2H, m), 7.21-7.28 (2H, m), 7.60 (1H, d, J=3.8Hz), 9.81 (1H, s).

10 Preparation Example 124. 5-(2-fluoro-benzyl)-thiophene-2-yl)-methanol

5-(2-Fluoro-benzyl)-thiophene-2-carbaldehyde described in Preparation Example 123 (2.81g, 12.7mmol) was dissolved in ethanol (40mL), sodium borohydride (964mg, 25.4mmol) was added at 0°C, and the solution was stirred at room temperature for 1 hour. Water was added to the reaction solution at 0°C,  
 15 which was then extracted with ethyl acetate, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 5 : 1), and the title compound (2.10g, 9.45mmol, 74%) was obtained as a colorless oil.

20 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.14 (2H, s), 4.73 (2H, s), 6.69 (1H, d, J=3.5Hz), 6.82 (1H, d, J=3.7Hz), 7.02-7.10 (2H, m), 7.19-7.26 (2H, m).

Preparation Example 125. 2-(5-(2-fluoro-benzyl)-thiophen-2-ylmethyl)-isoindol-1,3-dione

25 The title compound (1.49g, 4.24mmol, 45%) was obtained as a pale yellow

solid from (5-(2-fluoro-benzyl)-thiophen-2-yl)-methanol described in Preparation Example 124 (2.10g, 9.44mmol) according to an analogous method to Preparation Example 94.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.08 (2H, s), 4.92 (2H, s), 6.63 (1H, d, J=3.5Hz),  
5 6.94 (1H, d, J=3.5Hz), 6.99-7.08 (2H, m), 7.12-7.23 (2H, m), 7.70 (2H, dd, J=3.1,5.5Hz), 7.84 (2H, dd, J=3.1,5.5Hz).

Preparation Example 126. C-(5-(2-Fluoro-benzyl)thiophen-2-yl)-methylamine

The title compound (901mg, 4.07mmol, 96%) was obtained as a pale yellow  
10 oil from 2-(5-(2-fluoro-benzyl)-thiophen-2-ylmethyl)-isoindol-1,3-dione described in Preparation Example 125 (1.49g, 4.24mmol) according to an analogous method to Preparation Example 95.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.95 (2H, s), 4.12 (2H, s), 6.65 (1H, d, J=3.5Hz),  
15 6.71 (1H, d, J=3.5Hz), 7.01-7.09 (2H, m), 7.18-7.25 (2H, m).

Preparation Example 127. (5-Bromo-4-phenoxy-thiophen-2-yl)-methanol

2-(4-Phenoxy-thiophen-2-yl)-[1,3]dioxolane described in Preparation Example  
109 (4.88g, 19.7mmol) and N-bromosuccinimide (3.85g, 21.7mmol) were dissolved in tetrahydrofuran (100mL), and the solution was stirred for 4.5 hours at room  
20 temperature. An aqueous solution of saturated sodium bicarbonate was added to the reaction solution at 0°C, which was then extracted with ethyl acetate, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, and the residue was silica gel-filtered to obtain 2-(5-bromo-4-phenoxy-thiophen-2-yl)-[1,3]dioxolane (5.48g) as a pale yellow oil.

25 Then, 5-bromo-4-phenoxy-thiophene-2-carbaldehyde (3.11g) was obtained

as a colorless oil from 2-(5-bromo-4-phenoxy-thiophen-2-yl)-[1,3]dioxolane (5.48g) according to an analogous method to Preparation Example 34.

Then, the title compound (2.76g, 9.68mmol, 88%) was obtained as a colorless oil from 5-bromo-4-phenoxy-thiophene-2-carbaldehyde (3.11g, 11.0mmol) according to an analogous method to Preparation Example 93.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.70 (2H, s), 6.62 (1H, s), 6.98 (2H, dd, J=1.1, 8.8Hz), 7.09 (1H, tt, J=1.1, 7.5Hz), 7.29-7.34 (2H, m).

Preparation Example 128. 2-(5-Bromo-4-phenoxy-thiophen-2-ylmethyl)-

10 isoindol-1,3-dione

The title compound (2.66g, 6.42mmol, 68%) was obtained as a white solid from (5-bromo-4-phenoxy-thiophen-2-yl)-methanol described in Preparation Example 127 (2.71g, 9.50mmol) according to an analogous method to Preparation Example 94.

15 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.86 (2H, s), 6.76 (1H, s), 6.95 (2H, dd, J=1.1, 8.8Hz), 7.08 (1H, t, J=7.5Hz), 7.30 (2H, dd, J=7.3, 8.8Hz), 7.70-7.76 (2H, m), 7.83-7.88 (2H, m).

Preparation Example 129. C-(5-Bromo-4-phenoxy-thiophen-2-yl)-methylamine

20 The title compound (1.62g, 5.70mmol, 89%) was obtained as a colorless oil from 2-(5-bromo-4-phenoxy-thiophen-2-ylmethyl)-isoindole-1,3-diene described in Preparation Example 128 (2.66g, 6.42mmol) according to an analogous method to Preparation Example 95.

25 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.94 (2H, s), 6.53 (1H, t, J=1.1Hz), 6.97 (2H, dd, J=1.1, 8.6Hz), 7.08 (1H, tt, J=1.1, 7.5Hz), 7.31 (2H, dd, J=7.5, 8.8Hz).

Preparation Example 130. 3-Aminomethylphenol

The title compound (2.9g, 24mmol, 97%) was obtained as a white solid from 3-hydroxybenzaldehyde (3.0g, 24mmol) according to an analogous method to

5 Preparation Example 38.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.60 (2H, s), 6.55 (1H, d, J=7.5Hz), 6.68-6.70 (2H, m), 7.03-7.07 (1H, m).

Preparation Example 131. 1-Quinolin-6-yl-ethanone

10 6-Bromoquinoline (4.4g, 21.3mmol), 1-ethoxyvinyl (tri-*n*-butyl)tin (10g, 27.69mmol) and dichlorobis(triphenylphosphine)palladium(II) (1.2g, 1.7 mmol) were dissolved in toluene (120mL) under nitrogen atmosphere, and the solution was stirred at 80°C for 7 hours. At room temperature, 5M hydrochloric acid (30mL) and tetrahydrofuran (150 mL) were added to the reaction solution, which was then stirred  
15 for 15 hours. The reaction solution was poured into an aqueous solution of saturated sodium bicarbonate containing finely ground ice (200mL), the solution was adjusted to pH 8-9, and filtered through Celite pad. The filtrate was extracted with ethyl acetate twice, washed with brine twice, dried over anhydrous magnesium sulfate, and then, filtered. The organic layer was evaporated *in vacuo*, then, residue  
20 of reddish brown oil (16.8g) was obtained. Ethyl acetate (20mL) was added thereto, which was dissolved, then, silica gel(80mL) was added, the solvent was evaporated *in vacuo* for adsorption, purification was carried out by silica gel column chromatography (hexane : ethyl acetate = 95 : 5, then hexane : ethyl acetate = 60 : 40), and a yellow solid (4.16 g) was obtained. This yellow solid was dissolved in  
25 ethyl acetate (100mL), then, the solution was extracted with 1M hydrochloric acid (70,

50mL) twice, the aqueous layer was adjusted to pH 8 with sodium bicarbonate, then, the solution was extracted with ethyl acetate twice. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, then, filtered, organic layer was evaporated *in vacuo*, then, the title compound (2.98g, 17.4mmol, 77%) was obtained as a yellow solid.

<sup>1</sup>H-NMR Spectrum (Acetone-d<sub>6</sub>) δ(ppm) : 2.73 (3H, s), 7.07 (1H, dd, J=8.4, 4.4 Hz), 8.11 (1H, d, J=9.2 Hz), 8.27 (1H, dd, J=8.8, 2.0 Hz), 8.50-8.53 (1H, m), 8.70 (1H, m), 9.02 (1H, m).

#### 10 Preparation Example 132. 4-Benzyloxy-3-methoxymethoxy-benzonitrile

To a solution of 3,4-dihydroxy-benzonitrile (1.36g, 10mmol) and potassium *tert*-butoxide (1.5g, 13mmol) in dimethylsulfoxide (15mL) was added benzyl chloride (1.5mL, 13mmol) under nitrogen atmosphere at room temperature, and the solution was stirred for 24 hours. Silica gel (100mL) was added portionwise to the reaction solution for adsorption, a pale yellow oil of 4-benzyloxy-3-hydroxy-benzonitrile (2.38g) was obtained by silica gel column chromatography (hexane : ethyl acetate = 7 : 3), in addition, this was purified by NH silica gel column chromatography (hexane : ethyl acetate = 1 : 1, then ethyl acetate, then ethyl acetate : methanol = 90 : 10, then 85 : 15), and 4-benzyloxy-3-hydroxy-benzonitrile (0.588g, 2.61mmol, 24.8%) was obtained as a pale yellow solid.

To a solution of the resulting 4-benzyloxy-3-hydroxy-benzonitrile (300mg, 1.33mmol) and potassium *tert*-butoxide (300mg, 2.66mmol) in dimethylsulfoxide (4mL) was added chloromethyl methyl ether (0.204mL, 2.66mmol) portionwise under nitrogen atmosphere stirring on an ice bath, and the solution was stirred for 2 days. NH silica gel (25mL) was added to the reaction solution, purification was carried out

by NH silica gel column chromatography (hexane : ethyl acetate = 7 : 3) then silica gel column chromatography (hexane : ethyl acetate = 7 : 3), and the title compound (306mg, 0.829mmol, 85.3%) was obtained as a pale yellow solid.

<sup>1</sup>H-NMR Spectrum (Acetone-d<sub>6</sub>) δ(ppm) :3.46(3H, s), 5.26(2H, s), 5.27(2H,s),  
5 7.26(1H, d, J=8.4 Hz), 7.36-7.44(4H, m), 7.45(1H, d, J=2.4 Hz), 7.50-7.53(2H, m).

Preparation Example 133. 3-(3-Methyl-2-butenyloxy)-benzonitrile

3-Hydroxybenzonitrile (1.19g, 10mmol) and 4-bromo-2-methyl-2-butene (1.66g, 10mmol) were dissolved in N,N-dimethylformamide (5mL), potassium carbonate (1.66g, 12mmol) was added, and the solution was stirred at room  
10 temperature for 2 hours. Water (50mL) was added to the reaction solution, which was then extracted with ethyl acetate (50mL). The extract was washed, then, dried over anhydrous magnesium sulfate, and the solvent was evaporated *in vacuo*. The residue was purified by silica gel column chromatography (hexane : ethyl acetate =  
15 95 : 5), and the title compound (1.71g, 10mmol, 99.4%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) :1.73(3H, s), 1.78(3H, s), 4.51(2H, d, J=6.8Hz),  
5.48(1H, t, J=6.8Hz), 7.13(1H, dd, J=2.4, 8.4Hz), 7.15(1H, d, J=2.4Hz), 7.23(1H, d,  
J=8.4Hz), 7.28(1H, t, J=8.4Hz).

20

Preparation Example 134. 3-(3-Methyl-2-butenyloxy)-benzylamine

3-(3-Methyl-2-butenyloxy)-benzonitrile described in Preparation Example 133 (1.71g,10mmol) was dissolved in tetrahydrofuran (20mL), lithium aluminum hydride (0.57g,15mmol) was added under stirring at room temperature, the solution was  
25 heated to 70°C, then stirring continued for 2 hours. The reaction solution was

cooled on an ice bath, then, water (0.6mL), an aqueous solution of 15% sodium hydroxide (0.6mL) and water (1.8mL) were added in this order, then, the solid was filtered, and washed with tetrahydrofuran (20mL). The filtrate and the washings were combined, and dried over anhydrous sodium sulfate. The solvent was evaporated *in vacuo*, and the title compound (1.50g, 8.52mmol, 85.2%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 1.73(3H, s), 1.78(3H, s), 4.51(2H, d, J=6.8Hz), 4.68(2H, d, J=5.6Hz), 5.48(1H, t, J=6.8Hz), 6.77(1H, dd, J=2.4, 8.4Hz), 6.83(1H, d, J=2.4Hz), 6.87(1H, s), 7.23(1H, t, J=8.4Hz).

#### Preparation Example 135. (3-Cyanobenzyl)phosphonic acid diethyl ester

3-Cyanobenzyl bromide (9.8g, 50mmol) and triethylphosphite (9.97g, 60mmol) were stirred at 140°C for 3 hours. The resulting residue was evaporated, the distillate which has 145°C/1mmHg was collected, and the title compound (10g, 39.5mmol, 79.1%) was obtained as a colorless oil.

#### Preparation Example 136. 3-(2-Methylpropenyl) benzonitrile

Sodium hydride (0.40g, 10mmol, 60% in oil) was suspended in tetrahydrofuran (5mL), (3-cyanobenzyl)phosphonic acid diethyl ester obtained in Preparation Example 135 (2.53g, 10mmol) was added dropwise under stirring at room temperature. After stirring for 1 hour at 60°C, the solution was allowed to room temperature, acetone (0.92g, 20mmol) was added dropwise, and the solution was further stirred for 30 minutes at room temperature. Water (100mL) was added to the reaction solution, which was then extracted with ethyl acetate (50mL). The organic layer was washed with water, then, dried over anhydrous magnesium sulfate,



and the solvent was evaporated *in vacuo*. The residue was purified by silica gel column chromatography (hexane : ethyl acetate = 97 : 3), and the title compound (0.44g, 2.80mmol, 28%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) :1.85(3H, s), 1.93(3H, s), 6.23(1H, s),  
5 7.40-7.50(4H, m).

Preparation Example 137. (3-(2-Methylpropenyl)-benzylamine)

3-(2-Methylpropenyl)benzonitrile described in Preparation Example 136 (0.44g, 2.8mmol) was dissolved in tetrahydrofuran (5mL), lithium aluminum hydride  
10 (0.16g, 4.2mmol) was added under stirring at room temperature, the solution was heated to 70°C, followed by stirring for 2 hours. The reaction solution was cooled on an ice bath, then, water (0.16mL), an aqueous solution of 15% sodium hydroxide (0.16mL) and water (0.48mL) were sequentially added, then, the solid was filtered, and washed with tetrahydrofuran (10mL). The filtrate and the washings were  
15 combined, and dried over anhydrous sodium sulfate. The solvent was evaporated *in vacuo*, and the title compound (0.40g, 2.48mmol, 88.7%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) :1.87(3H, s), 1.90(3H, s), 3.85(2H, s), 6.27(1H, s),  
20 7.11-7.28(4H, m).

Preparation Example 138. 3-Cyclopentylidenemethylbenzonitrile

Potassium *tert*-butoxide (1.12g, 10mmol) was suspended in N,N-dimethylformamide (10mL), and (3-cyanobenzyl)phosphonic acid diethyl ester (2.53g, 10mmol) was added dropwise under stirring at room temperature. The  
25 solution was stirred at room temperature for 1 hour, then, cyclopentanone

(0.84g, 10mmol) was added, and the solution was stirred at room temperature for 2 hours. Water (100mL) was added to the reaction solution, which was then extracted with ethyl acetate (50mL). The organic layer was washed with water, then, dried over anhydrous magnesium sulfate, and the solvent was evaporated *in vacuo*. The residue was purified by silica gel column chromatography (hexane : ethyl acetate = 97 : 3), and the title compound (1.32g, 7.21mmol, 72.3%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 1.65-1.79(4H, m), 2.47-2.58(4H, m), 6.33(1H, s), 7.40-7.57(4H, m).

10

#### Preparation Example 139. 3-Cyclopentylidenemethyl-benzylamine

3-Cyclopentylidenemethylbenzonitrile described in Preparation Example 138 (1.32g, 7.21mmol) was dissolved in tetrahydrofuran (10mL), lithium aluminum hydride (0.41g, 10.8mmol) was added under stirring at room temperature, the solution was heated to 70°C, followed by stirring for 2 hours. The reaction solution was cooled on an ice bath, then, water (0.41mL), an aqueous solution of 15% sodium hydroxide (0.41mL) and water (1.23mL) were sequentially added, then, the solid was filtered, and washed with tetrahydrofuran (20mL). The filtrate and the washings were combined, and dried over anhydrous sodium sulfate. The solvent was evaporated *in vacuo*, and the title compound (1.30g, 6.95mmol, 96.4%) was obtained as a colorless oil.

20

#### Preparation Example 140. (5-Bromothiophen-2-yl)methanol

5-Bromo-2-thiophene carboxy aldehyde (25g, 131mmol) was dissolved in a mixture solvent of ethanol-tetrahydrofuran (1:1) (200mL), sodium borohydride (1.86g,

25

49mmol) was added in small amounts under ice-cold stirring, and stirring continued for 30 minutes. An aqueous solution of saturated ammonium chloride (20mL) was added to the reaction solution, which was then stirred for 30 minutes. The reaction solution was extracted with ethyl acetate (100mL), washed with water, dried over anhydrous magnesium sulfate, then, the solvent was evaporated *in vacuo*. The residue was purified by silica gel column chromatography (hexane : ethyl acetate = 90 : 10), and the title compound (6.7g, 34.7mmol, 26.5%) was obtained as a colorless oil.

10 Preparation Example 141. 2-Bromo-5-chloromethylthiophene

(5-Bromothiophen-2-yl)methanol described in Preparation Example 140 (6.7g, 34.7mmol) was dissolved in diethyl ether (40mL), 10mL of concentrated hydrochloric acid was added thereto, and the solution was vigorously stirred at room temperature for 8 hours. An ice water (200mL) was added to the reaction solution, an aqueous solution of sodium bicarbonate was further added for neutralization, then, the solution was extracted with ethyl acetate (100mL). The organic layer was washed with water, dried over anhydrous magnesium sulfate, then, the solvent was evaporated *in vacuo*, and the title compound (7.3g, 34.5mmol, 99.4%) was obtained as a colorless oil.

20

Preparation Example 142. (5-Bromothiophen-2-ylmethyl)phosphonic acid diethyl ester

2-Bromo-5-chloromethylthiophene described in Preparation Example 141 (7.3g, 34.5mmol) and triethylphosphite (6.35g, 38.2mmol) were stirred at 140°C for 3 hours. The resulting residue was purified by silica gel column chromatography

25

(hexane : ethyl acetate = 50 : 50), and the title compound (8.82g, 31.2mmol, 90.3%) was obtained as a reddish brown oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) :1.27-1.36(6H, m), 3.25(2H, d, J=24Hz), 4.05-4.16(4H, m), 6.72(1H, d, J=3.6Hz), 6.92(1H, d, J=3.6Hz).

5

Preparation Example 143. 2-Bromo-5-(2-methylpropenyl)-thiophene

(5-Bromothiophen-2-ylmethyl)phosphonic acid diethyl ester described in Preparation Example 142 (3.13g, 10mmol) was dissolved in tetrahydrofuran (20mL), sodium hydride (0.40g, 10mmol, 60% in oil) was added to this solution under stirring at room temperature. The solution was stirred for 30 minutes at 60°C, then, acetone (1g, 17.2mmol) was added, and the solution was further stirred for 30 minutes. Water (100mL) was added to the reaction solution, which was then extracted with hexane (50mL), the organic layer was washed with water, dried over anhydrous magnesium sulfate, then, the solvent was evaporated *in vacuo*. The residue was purified by silica gel column chromatography (hexane), and the title compound (60mg, 0.27mmol, 2.7%) was obtained as a colorless oil.

10

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) :1.90(3H, s), 1.93(3H, s), 6.27(1H, s), 6.62(1H, d, J=3.6Hz), 6.93(1H, d, J=3.6Hz).

20 Preparation Example 144. 5-(2-Methylpropenyl)-thiophene-2-carbonitrile

2-Bromo-5-(2-methylpropenyl)-thiophene described in Preparation Example 143 (60mg,0.27mmol) was dissolved in N,N-dimethylformamide (1mL), copper cyanide (62mg,0.69mmol) was added at 160°C, and the solution was stirred for 2 hours. The reaction solution was cooled, then, concentrated aqueous ammonia (5mL) was added, and the solution was extracted with diethyl ether (10mL). The

25

organic layer was washed, dried over anhydrous magnesium sulfate, then, the solvent was evaporated *in vacuo*. The residue was purified by silica gel column chromatography (hexane : ethyl acetate = 95 : 5), and the title compound (15mg, 0.092mmol ,34%) was obtained as a colorless oil.

- 5 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) :1.97(3H, s), 2.00(3H, s), 6.39(1H, s), 6.85(1H, d, J=4.0Hz), 7.50(1H, d, J=4.0Hz).

Preparation Example 145. C-(5-(2-Methylpropenyl)-thiophen-2-yl)-methylamine

- 5-(2-Methylpropenyl)-thiophene-2-carbonitrile described in Preparation  
10 Example 144 (15mg, 0.092mmol) was dissolved in tetrahydrofuran (2mL), lithium aluminum hydride (10mg,0.26mmol) was added under stirring at room temperature, the solution was heated to 70°C, followed by stirring for 2 hours. The reaction solution was cooled on an ice bath, then, water (0.01mL), an aqueous solution of 15% sodium hydroxide (0.01mL) and water (0.03mL) were sequentially added, then,  
15 the solid was filtered, and washed with tetrahydrofuran (5mL). The filtrate and the washings were combined, and dried over anhydrous sodium sulfate. The solvent was evaporated *in vacuo*, and the title compound (14mg, 0.083mmol, 91.1%) was obtained as a colorless oil.

- <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) :1.91(3H, s), 1.96(3H, s), 4.09(2H, s), 6.33(1H, s),  
20 6.75(1H, d, J=3.6Hz), 6.95(1H, d, J=3.6Hz).

Preparation Example 146. 3-Isobutylbenzylamine

- 3-(2-Methylpropenyl)-benzylamine described in Preparation Example 137  
(100mg, 0.621mmol) was dissolved in ethanol (5mL), 10% palladium-carbon (50%  
25 water wet, 20mg) was added thereto, and the mixture was stirred for 2 hours at room

temperature under hydrogen atmosphere. The catalyst was filtered, then, the solvent was evaporated, and the title compound (58mg, 56.6%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.90(6H, d, J=6.8Hz), 1.87(1H, dq, J=7.6, 6.8Hz),  
 5 2.48(2H, d, J=7.6Hz), 3.84(2H, s), 7.02-7.28(4H, m).

#### Preparation Example 147. 2-(2-cyclopropyl vinyl)thiophene

Potassium *tert*-butoxide (1.81g, 16.2mmol) was suspended in N,N-dimethylformamide (50mL), thiophen-2-ylmethyl triphenylphosphonium chloride  
 10 (6.38g,16.2mmol) was added while stirring at room temperature under a nitrogen stream, and the mixture was stirred at room temperature for 30 minutes. Then, cyclopropanecarboxyaldehyde (1.13g, 16.2mmol) was added dropwise to the reaction solution, and the solution was stirred for 1 hour. Water (100mL) was added to the reaction solution, which was then extracted with hexane (50mL). The organic  
 15 layer was passed through silica gel (10g) for filtration, the filtrate was evaporated *in vacuo*, and the title compound (1.27g, 8.47mmol, 52.3%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.44-0.52(2H, m), 0.76-0.84(2H, m), 1.451.55(1H, m), 5.60(1H, dd, J=8.8, 15.6Hz), 6.83(1H, d, J=3.2Hz), 6.92(1H, dd,  
 20 J=3.2, 5.2Hz), 7.00(1H, dd, J=3.2, 5.2Hz), 7.05(1H, J=5.2Hz).

#### Preparation Example 148. 5-(2-Cyclopropylvinyl)thiophene-2-carboxy aldehyde

2-(2-Cyclopropylvinyl)thiophene described in Preparation Example 147 (1.27g,8.47mmol) was dissolved in anhydrous diethyl ether (20mL), *n*-butyl lithium  
 25 (2.47M hexane solution, 4.1mL, 10.2mmol) was added dropwise under stirring on an

ice bath, and the solution was stirred for 30 minutes. The reaction solution was cooled on a dry ice-acetone bath, N,N-dimethylformamide (2g, 27.4mmol) was added thereto, and the solution was stirred as is for 30 minutes. Acetic acid (1mL) and water (10mL) were sequentially added to the reaction solution, which was then  
5 allowed to room temperature. The reaction solution was extracted with ethyl acetate (50mL), then, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 95 : 5), and the title compound (960mg, 5.39mmol, 63.7%) was obtained as a colorless oil.

10 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.57-0.63(2H, m), 0.88-0.94(2H, m), 1.53-1.60(1H, m), 5.84(1H, dd, J=9.2, 15.6Hz), 6.59(1H, d, J=15.6Hz), 6.91(1H, d, J=3.6Hz), 7.59(1H, d, J=3.6Hz), 9.80(1H, s).

Preparation Example 149. (5-(2-Cyclopropylvinyl)thiophen-2-yl)methanol

15 5-(2-Cyclopropylvinyl)thiophen-2-aldehyde described in Preparation Example 148 (960mg, 5.39mmol) was dissolved in a mixture solvent of tetrahydrofuran-ethanol (2:1) (30mL), sodium borohydride (100mg, 2.64mmol) was added under stirring on an ice bath, and the solution was stirred for 30 minutes. Acetic acid (0.5mL) and water (10mL) were added sequentially to the reaction  
20 solution, which was then extracted with ethyl acetate (50mL). The organic layer was washed with an aqueous solution of saturated sodium bicarbonate and brine, and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, and the title compound (930mg, 5.19mmol, 96.2%) was obtained as a colorless oil.

25 Preparation Example 150. 2-(5-(2-cyclopropylvinyl)thiophen-2-yl)methyl)

isoindol-1,3-dione

(5-(2-Cyclopropylvinyl)thiophen-2-yl)methanol described in Preparation Example 149 (930mg, 5.19mmol), triphenylphosphine (2040mg, 7.78mmol) and phthalimide (1140mg, 7.78mmol) were dissolved in tetrahydrofuran (50mL), azodicarboxylic acid dimethyl ester (1140mg, 7.78mmol) was added under stirring at room temperature, and the solution was stirred for 1 hour. Water (50mL) was added to the reaction solution, which was then extracted with ethyl acetate (50mL). The organic layer was washed with water, dried over anhydrous magnesium sulfate, and the solvent was evaporated *in vacuo*. The residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 9 : 1), and the title compound (330mg, 1.07mmol, 20.6%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.44-0.48(2H, m), 0.76-0.80(2H, m), 1.42-1.50(1H, m), 4.92(2H, s), 5.52(1H, dd, J=8.8, 15.6Hz), 6.47(1H, d, J=15.6Hz), 6.62(1H, d, J=3.6Hz), 6.93(1H, d, J=3.6Hz), 7.67-7.73(2H, m), 7.82-7.86(2H, m).

Preparation Example 151. C-(5-(2-Cyclopropylvinyl)thiophen-2-yl)methylamine

2-(5-(2-Cyclopropylvinyl)thiophen-2-ylmethyl)isoindol-1,3-dione described in Preparation Example 150 (330mg, 1.02mmol) was dissolved in ethanol (50mL), hydrazine monohydrate (500mg, 10mmol) was added thereto, and the solution was stirred under reflux for 2 hours. After cooling to room temperature, 2N sodium hydroxide solution (10mL) and water (100mL) were added to the reaction solution, which was then washed twice with hexane (50mL). The organic layer was dried over anhydrous sodium sulfate, then, the solvent was evaporated *in vacuo*, and the title compound (180mg, 1.01mmol, 98.6%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.44-0.48(2H, m), 0.76-0.80(2H, m),



1.42-1.50(1H, m), 3.96(2H, s), 5.52(1H, dd, J=8.8, 15.6Hz), 6.52(1H, d, J=15.6Hz), 6.65(1H, d, J=3.6Hz), 6.71(1H, d, J=3.6Hz).

Preparation Example 152. C-(5-(2,2-Dicyclopropylvinyl)thiophen-2-yl)methylamine

5           The title compound was synthesized according to an analogous method to Preparation Example 147 to 151.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.42-0.46(2H, m), 0.61-0.66(2H, m), 0.83-0.88(4H, m), 1.16-1.23(1H, m), 1.96-2.03(1H, m), 3.96(2H, s), 6.30(1H, s), 6.85(1H, d, J=3.2Hz), 6.90(1H, d, J=3.2Hz).

10

Preparation Example 153. Methanesulfonic acid 2-fluoro-benzyl ester

To a solution of 2-fluorobenzyl alcohol (4.40g, 34.9mmol) in dichloromethane (40mL) was added methanesulfonyl chloride (3.24mL, 41.9mmol) and triethylamine (5.84mL, 41.9mmol) on an ice bath, the solution was warmed to room temperature and stirred overnight. The reaction solution was diluted with dichloromethane, washed with an aqueous solution of 5% sodium bicarbonate, then, dried over anhydrous magnesium sulfate, the solvent was evaporated, and the title compound (4.62g, 65%) was obtained as a brown oil. This was used in the next reaction without purification.

20

Preparation Example 154. 4-(2-Fluorobenzyloxy)-benzylamine

The title compound (710mg, 14%) was obtained as a yellow oil from *p*-cyanophenol (2.70g, 22.7mmol) and methanesulfonic acid 2-fluoro-benzyl ester described in Preparation Example 153 (4.63g, 22.7mmol) according to an analogous method to Preparation Example 6.

25

Preparation Example 155. 4-(4-Fluorobenzyloxy)-benzylamine

4-(4-Fluorobenzyloxy)-benzonitrile (5.89g, quantitatively) was obtained from *p*-cyanophenol (3.00g, 25.2mmol) and 4-fluorobenzyl bromide (4.76g, 25.2mmol)

5 according to an analogous method to Preparation Example 6.

Next, the title compound (1.02g, 67%) was obtained as a yellow solid from the resulting 4-(4-fluorobenzyloxy)-benzonitrile (1.5g, 6.6mmol) according to an analogous method to Preparation Example 6.

10 Preparation Example 156. 5-(4-chloro-phenoxy)-thiophene-2-carbonitrile

The title compound (770mg, 3.27mmol, 65%) was obtained as a pale yellow oil from 5-nitrothiophene-2-carbonitrile (771mg, 5mmol) and 4-chlorophenol (643mg, 5mmol) according to an analogous method to Preparation Example 22.

<sup>1</sup>H-NMR Spectrum (Acetone-d<sub>6</sub>) δ(ppm) :6.72(1H, d, J=4.4Hz), 7.30-7.32(2H, m),  
15 7.50-7.52(2H, m), 7.67(1H, d, J=4.4Hz)

Preparation Example 157. C-(5-(4-Chloro-phenoxy)-thiophen-2-yl)-methylamine

The title compound (307mg, 1.28mmol, 86%) was obtained as an orange oil from 5-(4-chloro-phenoxy)-thiophene-2-carbonitrile described in Preparation Example  
20 156 (350mg, 1.49mmol) according to an analogous method to Preparation Example 23.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) :2.19(2H, brs), 3.81-3.82(2H, m), 6.53-6.54(1H, m), 6.69-6.70(1H, m), 7.10-7.13(2H, m), 7.42-7.45(2H, m).

25 Preparation Example 158. 5-(2-Chloro-phenoxy)-thiophene-2-carbonitrile

The title compound (516mg, 2.19mmol, 44%) was obtained as a white solid from 5-nitrothiophene-2-carbonitrile (771mg, 5mmol) and 2-chlorophenol (643mg, 5mmol) according to an analogous method to Preparation Example 22.

<sup>1</sup>H-NMR Spectrum (Acetone-d<sub>6</sub>) δ(ppm) : 6.63(1H, d, J=4.0Hz), 7.35-7.40(1H, m),  
5 7.42-7.50(2H, m), 7.61-7.65(2H, m).

Preparation Example 159. C-(5-(2-Chloro-phenoxy)-thiophen-2-yl)-methylamine

The title compound (305mg, 1.27mmol, 72%) was obtained as an orange oil from 5-(2-chloro-phenoxy)-thiophene-2-carbonitrile described in Preparation Example  
10 158 (356mg, 1.51mmol) according to an analogous method to Preparation Example 23.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) :2.10(2H, brs), 3.80(2H, s), 6.48-6.50(1H, m),  
6.66-6.72(1H, m), 7.15-7.23(2H, m), 7.34-7.38(1H, m), 7.56-7.59(1H, m).

15 Preparation Example 160. 5-(2-Fluoro-phenoxy)-thiophene-2-carbonitrile

The title compound (684mg, 3.12mmol, 77%) was obtained as a colorless oil from 5-nitrothiophene-2-carbonitrile (771mg, 5mmol) and 2-fluorophenol (673mg, 6mmol) according to an analogous method to Preparation Example 22.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) :6.76(1H, d, J=4.4Hz), 7.29-7.33(1H, m),  
20 7.35-7.41(1H, m), 7.43-7.53(2H, m), 7.79(1H, d, J=4.4Hz).

Preparation Example 161. C-(5-(2-Fluoro-phenoxy)-thiophen-2-yl)-methylamine

The title compound (298mg, 1.33mmol, 84%) was obtained as a brown oil from 5-(2-fluoro-phenoxy)-thiophene-2-carbonitrile described in Preparation Example  
25 160 (350mg, 1.60mmol) according to an analogous method to Preparation Example

23.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.79(2H, s), 6.45-6.47(1H, m), 6.64-6.70(1H, m), 7.15-7.27(3H, m), 7.34-7.41(1H, m).

5 Preparation Example 162. 3-Bromo-quinoline-6-carboxylic acid methyl ester

To a mixture of quinoline-6-carboxylic acid methyl ester (0.50g, 2.7mmol) and tetrahydrofuran(10mL) was added 1,3-dibromo-5,5-dimethyl hydantoin (0.76g, 2.7mmol) on an ice bath, and the solution was stirred at 50°C for 4 hours. Water, an aqueous solution of saturated sodium bicarbonate and ethyl acetate were added to  
10 the reaction solution, which was then partitioned, the organic layer was washed with brine, the solvent was then evaporated *in vacuo*. The residue was purified by silica gel column chromatography (hexane : ethyl acetate = 4 : 1), and the title compound (69mg, 0.26mmol, 10%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.01 (3H, s), 8.13 (1H, d, J=8.8Hz), 8.32 (1H, dd, J=2.0, 8.8Hz), 8.42 (1H, d, J=2.4Hz), 8.52 (1H, d, J=1.8Hz), 9.00 (1H, d, J=2.4Hz).  
15

Preparation Example 163. 3-Bromo-quinoline-6-carboxylic acid lithium salt

To a mixture of 3-bromo-quinoline-6-carboxylic acid methyl ester described in Preparation Example 162 (26mg, 0.098mmol) and tetrahydrofuran (2mL) were added  
20 methanol (0.2mL), lithium hydroxide monohydrate (4.1mg, 0.098mmol) and water (0.2mL), and the solution was stirred overnight at room temperature. The solvent was evaporated *in vacuo*, and the title compound (27mg) was obtained.

Preparation Example 164. Quinoline-6-carboxylic acid methyl ester N-oxide

25 To a mixture of quinoline-6-carboxylic acid methyl ester (4.7g, 25mmol) and

chloroform (80mL) was added 3-chloro-peroxyl benzoic acid (purity 65%, 8.6g, 33mmol) on an ice bath, and the solution was stirred at room temperature for 75 minutes. Water and an aqueous solution of 1N sodium hydroxide were added to the reaction solution, which was then partitioned, the organic layer was sequentially  
5 washed with an aqueous solution of saturated sodium thiosulfate and brine, and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, and the title compound (3.8g, 19mmol, 75%) was obtained as a pale yellow solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.03 (3H, s), 7.38 (1H, dd, J=6.0, 8.4Hz), 7.84 (1H, d, J=8.4Hz), 8.34 (1H, dd, J=1.8, 9.2Hz), 8.60 (1H, dd, J=0.9, 6.0Hz), 8.63 (1H,  
10 d, J=1.8Hz) , 8.81 (1H, d, J=9.2Hz).

Preparation Example 165. 2-Chloro-quinoline-6-carboxylic acid methyl ester

Phosphorus oxychloride (10mL) was added to quinoline-6-carboxylic acid methyl ester N-oxide (1.5g, 7.6mmol), and the solution was refluxed for 2 hours.  
15 The reaction solution was poured onto an ice, and warmed gradually to room temperature. Ethyl acetate was added to the reaction solution, which was then extracted, and sequentially washed with an aqueous solution of saturated sodium bicarbonate and brine. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 2 : 1), and the  
20 title compound (0.47g, 2.1mmol, 28%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.01 (3H, s), 7.47 (1H, dd, J=0.6, 8.6Hz), 8.07 (1H, d, J=8.8Hz), 8.22 (1H, d, J=8.6Hz), 8.32-8.35 (1H, m), 8.59 (1H, s).

Preparation Example 166. 2-Chloro-quinoline-6-carboxylic acid lithium salt

25 The title compound (54mg) was obtained as a crude compound from

2-chloro-quinoline-6-carboxylic acid methyl ester described in Preparation Example 165 (40mg, 0.18mmol) according to an analogous method to Preparation Example 163.

5 Preparation Example 167. C-(5-(2-Chloro-phenoxy)-thiophen-2-yl)-methylamine

To a solution of lithium aluminum hydride (1.79g, 47.1mmol) in tetrahydrofuran was added aluminum chloride (7.54g, 56.5mmol) on an ice bath, which was then stirred for 10 minutes. To the suspension was 5-(2-chloro-phenoxy)-thiophene-2-carbonitrile described in Preparation Example 158 (2.22g, 9.42mmol) was added on an ice bath, and the solution was stirred for 1 hour. An aqueous ammonia was added to the reaction solution, then, anhydrous magnesium sulfate was added for drying, and the solution was filtered through Celite pad. The filtrate was concentrated *in vacuo*, the residue was purified by NH silica gel column chromatography (heptane / ethyl acetate = 1 / 2), and the title compound (2.26g, 9.42mmol, 100%) was obtained as a light brown oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.81(2H, s), 6.56(1H, d, J=3.6Hz), 6.69(1H, dd, J=1.2, 3.6Hz), 7.05(1H, dd, J=2.4, 8.4Hz), 7.11(1H, t, J=2.0Hz), 7.19(1H, dt, J=1.2, 8.0Hz), 7.40(1H, t, J=8.0Hz).

20 Preparation Example 168. 5-(4-Fluoro-phenoxy)-furan-2-carbaldehyde

To a solution of 4-fluorophenol (2.39g, 21.3mmol) in dimethylsulfoxide (20mL) was added sodium hydride (785mg, 19.6-23.6mmol, 60-72% in oil), and the solution was stirred at room temperature for 20 minutes. Next, a solution of 5-nitro-furan-2-carbaldehyde (3g, 21.3mmol) in dimethylsulfoxide (10mL) was added dropwise, the solution was then stirred for 2 hours at room temperature. The

reaction mixture was poured into brine, and the mixture was extracted with ethyl acetate. The fractionated organic layer was dried over anhydrous magnesium sulfate, then concentrated, and the title compound (4.3g, 20.8mmol, 98%) was obtained.

- 5 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 5.23 (1H, d, J=4.0 Hz), 7.07-7.24 (5H, m), 9.40 (1H, s).

Preparation Example 169. C-(5-(4-Fluoro-phenoxy)-furan-2-yl)methylamine

- A suspension of 5-(4-fluoro-phenoxy)-furan-2-carbaldehyde described in  
10 Preparation Example 168 (4.3g, 20.9mmol), Raney nickel (1.5g) and 7N ammonia-methanol solution (40mL) was stirred at room temperature for 15 hours under hydrogen atmosphere (1 atm). The reaction solution was filtered through Celite pad to remove the catalyst, and the filtrate was concentrated. The residue was dissolved in ethyl acetate, filtered by NH silica gel lined over a glass filter, and  
15 then, this filtrate was concentrated to obtain the title compound (3.5g, 16.9mmol, 81%).

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.55 (2H, s), 5.64 (1H, d, J=3.2 Hz), 6.13-6.16 (1H, m), 7.04-7.10 (2H, m), 7.17-7.24 (2H, m).

20 Preparation Example 170. 4-(Pyridin-2-ylmethoxy)-benzonitrile

- To a solution of 4-cyanophenol (5g, 42mmol) in N,N-dimethyl formamide (40mL) were added potassium carbonate (17.4g, 126mmol) and 2-picolyl bromide hydrobromide (10.6g, 42mmol), and the solution was stirred at room temperature for 15 hours. The reaction solution was poured into brine, and the solution was  
25 extracted with ethyl acetate. The fractionated organic layer was dried over

anhydrous magnesium sulfate, then, filtered by NH silica gel lined over a glass filter, and the filtrate was concentrated to obtain the title compound (4.7g, 22.4mmol, 53%).

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 5.25 (2H, s), 7.03-7.07 (2H, m), 7.25-7.29 (1H, m), 7.48 (1H, d, J=8.0 Hz), 7.57-7.61 (2H, m), 7.74 (1H, dt, J=1.6 Hz, 8.0Hz),  
5 8.61-8.63 (1H, m).

Preparation Example 171. 4-(Pyridin-2-ylmethoxy)-benzylamine

To a solution of 4-(pyridine-2-ylmethoxy)-benzonitrile described in Preparation Example 170 (1.2g, 5.70mmol) in tetrahydrofuran (30mL) was added lithium aluminum hydride (0.22g, 5.80mmol), and the solution was stirred for 2 hours  
10 30 minutes at room temperature. Ice water was added to the reaction solution, which was then stirred for 30 minutes. This mixture was filtered through Celite pad, and washed with ethyl acetate. The filtrate was partitioned, this organic layer was dried over anhydrous magnesium sulfate, then, concentrated, and the title compound (1.1g, 5.13mmol, 90%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.81 (2H, s), 5.21 (2H, s), 6.94-6.98 (2H, m),  
15 7.20-7.26 (3H, m), 7.52 (1H, d, J=7.8 Hz), 7.74 (1H, dt, J=1.6 Hz, 7.8 Hz), 8.59-8.62 (1H, m).

Preparation Example 172. 4-(Pyridin-2-yloxymethyl)-benzylamine

To a solution of 2-bromopyridine (2.35g, 15.0mmol) and  
20 4-(hydroxymethyl)benzonitrile (3.00g, 22.5mmol) in N,N-dimethylformamide (20mL) was added sodium hydride (0.90g, 22.5mmol; 60% in oil), and the solution was stirred at 70°C for 30 minutes. The reaction solution was allowed to room temperature, then, partitioned in ethyl acetate and water, the organic layer was  
25 washed with water, and then, dried over anhydrous magnesium sulfate. The solvent



was evaporated, the residue was purified by NH silica gel column chromatography (hexane/ethyl acetate), and a white solid (581mg, 18%) was obtained.

To a solution of the obtained white solid (100mg, 0.476mmol) in tetrahydrofuran (3mL) was added lithium aluminum hydride (45mg, 1.19mmol), and the solution was stirred at room temperature for 1 hour. The reaction solution was partitioned in ethyl acetate and water, and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated, and the title compound (71mg, 70%) was obtained as a colorless solid, which is a crudely purified product.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.70(2H, s), 5.31(2H, s), 6.84-6.87(1H, m), 6.97-7.00(1H, m), 7.32(2H, d, J=8.0Hz), 7.37(2H, d, J=8.0Hz), 7.69-7.74(1H, m), 8.16-8.18(1H, m).

#### Preparation Example 173. 5-Bromo-2,3-dihydro-benzofuran

To a solution of 2,3-dihydrobenzofuran (15.0g, 125mmol) in tetrahydrofuran (300mL) was added N-bromosuccinimide (24.5g, 138mmol) at 0°C. The reaction solution was stirred at room temperature for 50 minutes, then, water was added, the solution was extracted with ethyl acetate, and the organic layer was washed with brine. Anhydrous magnesium sulfate was added to the organic layer for drying, which was then filtered, the filtrate was concentrated *in vacuo*, the residue was purified by NH silica gel column chromatography (hexane), and the title compound (24.0g, 97%) was obtained as a colorless oily substance.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.19(2H, t, J=8.6Hz), 4.54(2H, t, J=8.6Hz), 6.72(1H, d, J=8.2Hz), 7.22(1H, dd, J=2.2. 8.4Hz), 7.40(1H, s).

Preparation Example 174. 2,3-Dihydro-benzofuran-5-carbaldehyde

To a solution of 5-bromo-2,3-dihydro-benzofuran described in Preparation Example 173 (15.0g, 75.4mmol) in tetrahydrofuran (300mL) was added *n*-butyl lithium (31.2mL, 82.9mmol) at -78°C. The reaction solution was stirred for 85  
5 minutes at -78°C, then, N,N-dimethylformamide (6.42mL, 82.9mmol) was added thereto, followed by stirring at room temperature for 1 hour. 1N Hydrochloric acid was added to the reaction solution, which was then extracted with ethyl acetate, and the organic layer was washed with brine. Anhydrous magnesium sulfate was added to the organic layer for drying, which was then filtered, the filtrate was concentrated *in*  
10 *vacuo*, the residue was purified by silica gel column chromatography (hexane / ethyl acetate = 5 / 1), and the title compound (10.1g, 90%) was obtained as a pale yellow oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.26(2H, t, J=8.6Hz), 4.67(2H, t, J=8.6Hz), 6.96(1H, d, J=7.9Hz), 7.72(1H, d, J=8.2Hz), 7.77(1H, s), 9.82(1H, s).

15

Preparation Example 175. Benzofuran-5-carbaldehyde

To a solution of 2,3-dihydro-benzofuran-5-carbaldehyde described in Preparation Example 174 (6.00g, 40.5mmol) in toluene (120mL) was added 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (18.3g, 81mmol), and the solution was  
20 refluxed for 4 hours 30 minutes. The reaction solution was cooled to room temperature, water was added, the solution was extracted with ethyl acetate and tetrahydrofuran, and the organic layer was washed with brine. Anhydrous magnesium sulfate was added to the organic layer for drying, which was then filtered, the filtrate was concentrated *in vacuo*, the residue was purified by silica gel column  
25 chromatography (hexane / ethyl acetate = 5 / 1), and the title compound (1.24g, 21%)

was obtained as a pale yellow oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 6.91(1H, dd, J=0.92, 2.2Hz), 7.63(1H, d, J=8.6Hz), 7.74(1H, d, J=2.4Hz), 7.89(1H, dd, J=1.7, 8.6Hz), 8.16(1H, s), 10.08(1H, s).

5 Preparation Example 176. Benzofuran-5-yl-(5-bromo-thiophen-2-yl)-methanol

To a solution of 2,5-dibromothiophene (2.05g, 8.48mmol) in tetrahydrofuran (25mL) was added *n*-butyl lithium (3.48mL, 8.48mmol) at -78°C, and the solution was stirred for 40 minutes. Then, benzofuran-5-carbaldehyde described in Preparation Example 175 (1.24g, 8.48mmol) was added to this reaction solution at -78°C, and the  
 10 solution was stirred at room temperature for 75 minutes. Water was added to the reaction solution, which was then extracted with ethyl acetate, and the organic layer was washed with brine. Anhydrous magnesium sulfate was added to the organic layer for drying, which was then filtered, the filtrate was concentrated *in vacuo*, the residue was purified by NH silica gel column chromatography (heptane / ethyl acetate  
 15 = 3 / 1), and the title compound (2.11g, 81%) was obtained as a light brown oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 5.98(1H, s), 6.40(1H, d, J=2.9Hz), 6.68(1H, dd, J=0.92, 3.8Hz), 6.96(1H, dd, J=0.92, 2.2Hz), 7.01(1H, d, J=3.8Hz), 7.34(1H, dd, J=1.5, 8.4Hz), 7.55(1H, d, J=8.4Hz), 7.68(1H, s), 7.99(1H, d, J=2.2Hz).

20 Preparation Example 177. 5-(Benzofuran-5-yl-hydroxy-methyl)-thiophene-2-carbonitrile

To a solution of benzofuran-5-yl-(5-bromo-thiophen-2-yl)-methanol described in Preparation Example 176 (755mg, 2.44mmol) in 1-methyl-2-pyrrolidinone (15mL) were added zinc cyanide (344mg, 2.93mmol) and  
 25 tetrakis(triphenylphosphine)palladium (282mg, 0.244mmol), and the mixture was

stirred at 120°C for 3 hours. The reaction mixture was cooled to room temperature, then, aqueous ammonia was added, and the solution was filtered through Celite pad. The mother liquor was extracted with ethyl acetate, and the organic layer was washed with brine. Anhydrous magnesium sulfate was added to the organic layer for drying, which was then filtered, the filtrate was concentrated *in vacuo*, the residue was purified by silica gel column chromatography (heptane / ethyl acetate = 3 / 1), and the title compound (364mg, 58%) was obtained as a pale yellow oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 6.13(1H, s), 6.73(1H, s), 6.97(1H, d, J=2.2Hz), 6.99(1H, d, J=3.8Hz), 7.37(1H, dd, J=1.5, 8.4Hz), 7.58(1H, d, J=8.6Hz), 7.72(1H, s), 7.78(1H, d, J=3.8Hz), 8.01(1H, d, J=2.2Hz).

Preparation Example 178. C-(5-Benzofuran-5-ylmethyl-thiophen-2-yl)-methylamine

To a solution of lithium aluminum hydride (488mg, 12.9mmol) in tetrahydrofuran (10mL) was added aluminum chloride (1.72g, 12.9mmol) on an ice bath, and the solution was stirred for 10 minutes. 5-(Benzofuran-5-yl-hydroxy-methyl)-thiophene-2-carbonitrile described in Preparation Example 177 (364mg, 1.43mmol) was added to this suspension on an ice bath, and the solution was stirred for 3 hours. Aqueous ammonia was added to the reaction solution, then, anhydrous magnesium sulfate was added for drying, and the solution was filtered through Celite pad. The filtrate was concentrated *in vacuo*, the residue was purified by NH silica gel column chromatography (heptane / ethyl acetate = 1 / 1), and the title compound (285mg, 82%) was obtained as a light brown oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.77(2H, s), 4.15(2H, s), 6.70(2H, s), 6.91(1H, d, J=2.2Hz), 7.19(1H, dd, J=1.7, 8.4Hz), 7.50-7.52(2H, m), 7.96(1H, d, J=2.2Hz).

Preparation Example 179. C-(6-Benzyloxypyridin-3-yl)-methylaniline

To a solution of 2,5-dibromopyridine (5.0g, 21.1mmol) and benzyl alcohol (3.28mL, 31.7mmol) in N,N-dimethylformamide (50mL) was added sodium hydride (1.27g, 31.7mmol; 60% in oil), and the solution was stirred at 70°C for 2 hours. The reaction solution was allowed to room temperature, then, partitioned in ethyl acetate and water, the organic layer was washed with water, and then, dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by silica gel column chromatography (hexane / ethyl acetate), and a colorless oil (4.60g, 83%) was obtained.

To a solution of the resulting colorless oil (2.0g, 7.60mmol) in N,N-dimethylformamide (20mL) were added zinc cyanide (1.78g, 15.2mmol) and tetrakis(triphenylphosphine)palladium (0) (878mg, 0.760mmol) under nitrogen atmosphere, and the solution was stirred at 140°C for 4 hours. The reaction solution was allowed to room temperature, then, partitioned in ethyl acetate and water, the organic layer was washed with water, and then, dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by silica gel column chromatography (hexane/ethyl acetate), and a pale yellow solid (1.15g, 72%) was obtained.

To a solution of the pale yellow solid (100mg, 0.476mmol) in tetrahydrofuran (3mL) was added lithium aluminum hydride (45mg, 0.120mmol), and the solution was stirred at room temperature for 1 hour. The reaction solution was partitioned in ethyl acetate and water, and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated, and the title compound (75mg, 74%) was obtained as a yellow oil, which was a crudely purified product.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.37-4.38(2H, br d), 5.36(2H, s), 6.83-6.87(1H,

m), 7.30-7.46(4H, m), 7.74-7.76(1H, m), 8.17(1H, s), 8.80-8.83(1H, m).

Preparation Example 180. C-(6-Benzyl-pyridin-3-yl)-methylamine

A solution of 2,5-dibromopyridine (10g, 42.2mmol) in diethyl ether (260mL)  
5 was cooled to -78°C under nitrogen atmosphere, and *n*-butyl lithium (17.8mL, 46.4mmol; 2.6M hexane solution) was added dropwise. This solution was stirred at -78°C for 15 minutes, then, a solution of N,N-dimethylformamide (4.94mL) in diethyl ether solution (10mL) was added dropwise at -78°C. This solution was warmed to 0°C, and further stirred for 2 hours at this temperature. After the reaction was  
10 completed, this solution was partitioned in diethyl ether and water. The organic layer was separated, washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by silica gel column chromatography (hexane : ethyl acetate), and an aldehyde derivative (5.82g, 74%) was obtained.

15 To a solution of the resulting aldehyde derivative (5.82g, 31.3mmol) in toluene (120mL) were added ethyleneglycol (17.5mL, 0.313mol) and D-10-camphor-sulfonic acid (73mg, 0.313mmol), and the solution was refluxed for 7 hours. The reaction mixture was cooled to room temperature, then, washed with an aqueous solution of saturated sodium hydrogen carbonate and brine, and the organic  
20 layer was separated. The organic layer was dried over anhydrous magnesium sulfate, then, solvent was evaporated, the residue was purified by silica gel column chromatography (hexane : ethyl acetate), and an acetal (6.65g, 93%) was obtained.

Next, to a suspension of zinc (1.53g, 23.4mmol) in tetrahydrofuran (120mL) was added benzyl bromide (2.1mL, 17.6mmol) dropwise over 15 minutes, at 0°C,  
25 and the solution was stirred at this temperature for 4 hours. After 4 hours,

bis(triphenylphosphine)nickel(II) chloride (1.58g, 2.42mmol) and a solution of the acetal (3.0g, 13.1mmol) in tetrahydrofuran (90mL) were added to this suspension, and the solution was stirred at room temperature for 12 hours. After the reaction was completed, this suspension was partitioned in ethyl acetate and an aqueous solution of saturated ammonium chloride. The organic layer was separated, washed with water and brine, and then, dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by silica gel column chromatography (hexane : ethyl acetate), and a 2-benzyl pyridine derivative (1.08g, 34%) was obtained.

To a mixture solution of the resulting 2-benzyl pyridine derivative (1.08g, 4.48mmol) in methanol and tetrahydrofuran (5mL:4mL) was added 2N hydrochloric acid (5mL), and the solution was stirred at room temperature for 2 hours. Furthermore, to this solution was added 5N hydrochloric acid (8mL) was added in three times, the solution was stirred at room temperature for 24 hours, then, refluxed for 30 minutes. The reaction solution was cooled to room temperature, then, partitioned in ethyl acetate and an aqueous solution of saturated sodium bicarbonate. The organic layer was separated, washed with water, and then, dried over anhydrous magnesium sulfate. The solvent was evaporated, and a formyl derivative (950mg, quantitatively) was obtained.

To a solution of the resulting formyl derivative (950mg, 4.48mmol) in methanol (25mL) was added sodium borohydride (176mg, 4.66mmol), and the solution was stirred at room temperature for 1 hour. The reaction solution was partitioned in ethyl acetate and water. This organic layer was separated, washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated, and a benzyl alcohol derivative (810mg, 91%) was obtained.

Methanesulfonyl chloride (0.37mL, 4.83mmol) and triethylamine (0.67mL, 4.84mL) were added to a solution of the resulting benzyl alcohol derivative (810mg, 4.07mmol) in dichloromethane (8mL) that had been cooled to 0°C, and the solution was stirred at room temperature for 14 hours. The reaction mixture was partitioned  
5 in dichloromethane and an aqueous solution of saturated sodium bicarbonate. The organic layer was separated, washed with brine, and then, dried over anhydrous magnesium sulfate. The solvent was evaporated, and a methanesulfonate ester derivative (1.09g, 97%) was obtained.

To a solution of the resulting methanesulfonate ester derivative (1.09g, 3.93mmol) in N,N-dimethylformamide (10mL) was added phthalimide potassium salt (757mg, 4.09mmol), and the solution was stirred for 2 hours under reflux. This reaction mixture was cooled to room temperature, and partitioned in ethyl acetate and water. This organic layer was separated, washed with water and brine, and then, dried over anhydrous magnesium sulfate. The solvent was evaporated, the  
10 residue was purified by silica gel column chromatography (hexane : ethyl acetate), and a phthalimide derivative (910mg, 71%) was obtained.

Then, to a solution of the resulting phthalimide derivative (910mg, 2.77mmol) in ethanol (23mL) was added hydrazine monohydrate (144mg, 2.88mmol), and the solution was stirred for 2 hours under reflux. The reaction mixture was cooled to  
20 room temperature, then, water was added to this mixture. This mixture was concentrated *in vacuo* until the liquid volume of this mixture became half. This concentrated solution was partitioned in ethyl acetate and water. This organic layer was separated, washed with 2N sodium hydroxide and water, and dried over anhydrous magnesium sulfate. The solvent was evaporated, and the title  
25 compound (360mg, 66%) was obtained. This was used in the next reaction without



further purification.

Preparation Example 181. 2-Amino-methyl-5-phenoxy-pyridine

2-Cyano-5-phenoxy-pyridine (150mg, 0.76mmol) and lithium aluminum  
5 hydride (58mg, 1.53mmol) were suspended in tetrahydrofuran (5mL) and diethyl  
ether (5mL), and the solution was stirred for 10 minutes under reflux. The reaction  
solution was partitioned in water and ethyl acetate. The organic layer was  
separated, the solvent was evaporated, and the title compound (140mg, brown oil)  
was obtained as a crude product.  
10 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.80 (2H, s), 6.70-6.80 (2H, m), 7.00-7.12  
(1H, m), 7.12-7.22 (2H, m), 7.40-7.50 (2H, m), 8.30 (1H, s).

Preparation Example 182. 5-Benzyloxy-2-methyl-pyridine

To a solution of 3-hydroxy-6-methylpyridine (5.00g, 45.8mmol) in  
15 N,N-dimethylformamide (50mL) was added sodium hydride (2.02g, 50.4mmol, 60% in  
oil) at 0°C, and the solution was stirred for 15 minutes at 0°C. Then, benzyl bromide  
(5.99mL, 50.4mmol) was added at 0°C, and the solution was stirred for 3.5 hours at  
room temperature. The reaction solution was partitioned in water and ethyl acetate.  
This organic layer was separated, washed with water and brine and dried over  
20 anhydrous magnesium sulfate. The organic layer was filtered, then, the solvent was  
evaporated *in vacuo*, the residue was purified by silica gel column chromatography  
(heptane / ethyl acetate = 2 / 1), and the title compound (5.99g, 66%) was obtained  
as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 2.49(3H, s), 5.08(2H, s), 7.05(1H, d, J=8.6Hz),  
25 7.17(1H, dd, J=2.9Hz, 8.4Hz), 7.31-7.44(5H, m), 8.27(1H, d, J=2.9Hz).

Preparation Example 183. (5-Benzyloxy-pyridin-2-yl)-methanol

To a solution of 5-benzyloxy-2-methyl pyridine described in Preparation Example 182 (5.99g, 30.1mmol) in methylene chloride (100mL) was added  
5 3-chloroperoxybenzoic acid (8.79g, 33.1mmol, 65%) at 0°C, and the solution was stirred at room temperature for 2 hours. An aqueous solution of saturated sodium bicarbonate was added to the reaction solution at 0°C, which was partitioned with methylene chloride. This organic layer was separated, washed with an aqueous solution of saturated sodium bicarbonate and brine, and dried over anhydrous  
10 magnesium sulfate. The organic layer was filtered, then, the solvent was evaporated *in vacuo*, and 5-benzyloxy-2-methyl-pyridine-1-oxide (7.71g) was obtained as a crude product of a white solid. Next, acetic anhydride (77mL) was added to 5-benzyloxy-2-methyl-pyridine-1-oxide (7.71g), and the solution was stirred for 80 minutes at 120°C. This reaction mixture was cooled to room temperature,  
15 then, the solvent was evaporated *in vacuo*. Ethanol (50mL) and an aqueous solution of 5N sodium hydroxide (7mL) were added to the resulting residue, and the solution was stirred at room temperature for 50 minutes. The solvent was evaporated *in vacuo*, this residue was partitioned in brine and ethyl acetate. The organic layer was separated, washed with brine and dried over anhydrous  
20 magnesium sulfate. The organic layer was filtered, then, the solvent was evaporated *in vacuo*, the residue was purified by NH silica gel column chromatography (heptane / ethyl acetate = 1 / 1), and the title compound (4.17g, 54%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.46(2H, d, J=5.9Hz), 5.15(2H, s), 5.26(1H, t,  
25 J=5.9Hz), 7.29-7.40(4H, m), 7.42-7.45(3H, m), 8.22(1H, d, J=2.9Hz).

Preparation Example 184. 2-(5-Benzyloxy-pyridin-2-ylmethyl)-isoindol-1,3-dione

To a solution of (5-benzyloxy-pyridin-2-yl)-methanol described in Preparation Example 183 (2.00g, 9.29mmol) in tetrahydrofuran (40mL) were added phthalimide (1.50g, 10.2mmol), triphenylphosphine (2.92g, 11.1mmol) and diethyl azodicarboxylate (5.08mL, 11.1mmol, 40% toluene solution) at 0°C, and the solution was stirred at room temperature for 2 hours. The reaction solution was partitioned in water and ethyl acetate. The organic layer was separated, washed with brine and dried over anhydrous magnesium sulfate. The organic layer was filtered, then, the solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (heptane / ethyl acetate = 2 / 1), and the title compound (4.1g, quantitatively) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.84(2H, s), 5.15(2H, s), 7.31-7.45(7H, m), 7.86-7.92(4H, m), 8.20(1H, d, J=2.9Hz).

Preparation Example 185. C-(5-Benzyloxy-pyridin-2-yl)-methylamine

2-(5-Benzyloxy-pyridin-2-ylmethyl)-isoindol-1,3-dione described in Preparation Example 184 (4.10g, 11.9mmol) was dissolved in ethanol (40mL) and tetrahydrofuran (40mL). Hydrazine monohydrate (5.77mL, 119mmol) was added to this solution at room temperature, and the solution was stirred under reflux for 50 minutes. The reaction solution was partitioned in water and ethyl acetate. This organic layer was separated, washed with brine and dried over anhydrous magnesium sulfate. The organic layer was filtered, then, the solvent was evaporated *in vacuo*, the residue was purified by NH silica gel column chromatography (ethyl acetate), and the title compound (2.8g, quantitatively) was

obtained as a pale yellow solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.70(2H, s), 5.15(2H, s), 7.30-7.45(7H, m), 8.23(1H, d, J=3.2Hz).

5 Preparation Example 186. 2-Methyl-5-phenoxyethyl-pyridine

Thionyl chloride (1mL) was added to (6-methyl-pyridin-3-yl)-methanol (300mg, 2.44mmol) at 0°C, and the solution was stirred at room temperature for 20 minutes. The reaction solution was partitioned in an aqueous solution of saturated sodium bicarbonate and ethyl acetate. This organic layer was separated, washed with brine, and then, dried over anhydrous magnesium sulfate. The organic layer was filtered, then, the solvent was evaporated *in vacuo*. N,N-dimethylformamide (5mL), phenol (230mg, 2.44mmol) and potassium carbonate (674mg, 4.88mmol) were added to this residue at room temperature. This reaction mixture was stirred at room temperature for 40 minutes, then, further stirred for at 60°C 40 minutes. The reaction solution was partitioned in water and ethyl acetate. The organic layer was separated, washed with water and brine, and then, dried over anhydrous magnesium sulfate. The organic layer was filtered, then, the solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (heptane / ethyl acetate = 2 / 1), and the title compound (323mg, 66%) was obtained as a white solid.

20 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 2.58(3H, s), 5.04(2H, s), 6.96-7.00(3H, m), 7.18(1H, d, J=8.2Hz), 7.30(2H, t, J=8.8Hz), 7.67(1H, dd, J=2.0Hz, 8.0Hz), 8.56(1H, s).

Preparation Example 187. (5-Phenoxyethyl-pyridin-2-yl)-methanol

25 To a solution of 2-methyl-5-phenoxyethyl-pyridine described in Preparation

Example 186 (323mg, 1.62mmol) in methylene chloride was added 3-chloroperoxybenzoic acid (473mg, 1.78mmol, 65%) at 0°C, and the solution was stirred at room temperature for 7 hours. An aqueous solution of saturated sodium bicarbonate was added to the reaction solution at 0°C, which was then partitioned with ethyl acetate. The organic layer was separated, washed with an aqueous solution of saturated sodium bicarbonate and brine, and then, dried over anhydrous magnesium sulfate. The organic layer was filtered, then, the solvent was evaporated *in vacuo*. Acetic anhydride (4mL) was added to this residue, which was then stirred for 30 minutes at 120°C. This reaction solution was cooled to room temperature, then, the solvent was evaporated *in vacuo*. Ethanol (5mL), and an aqueous solution of 5N sodium hydroxide (2mL) were added to this residue, and the solution was stirred at room temperature for 45 minutes. This reaction solution was concentrated *in vacuo*, the residue was partitioned in water and ethyl acetate. The organic layer was separated, washed with brine, and then, dried over anhydrous magnesium sulfate. The organic layer was filtered, then, the solvent was evaporated *in vacuo*, the residue was purified by NH silica gel column chromatography (heptane / ethyl acetate = 1 / 1), and the title compound (167mg, 48%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.63(1H, t, J=5.2Hz), 4.78(2H, d, J=4.8Hz), 5.09(2H, s), 6.97-7.02(3H, m), 7.27-7.33(3H, m), 7.78(1H, dd, J=2.2Hz, 8.0Hz), 8.63(1H, d, J=1.7Hz).

Preparation Example 188. 2-(5-Phenoxymethyl-pyridin-2-yl)methyl)-isoindol-1,3-dione

To a solution of (5-phenoxymethyl-pyridin-2-yl)-methanol described in Preparation Example 187 (167mg, 0.776mmol) in tetrahydrofuran (4mL) were added

phthalimide (126mg, 0.856mmol), triphenylphosphine (244mg, 0.930mmol) and diethyl azodicarboxylate (424 $\mu$ L, 0.931mmol, 40% toluene solution) at 0°C, and the solution was stirred at room temperature for 30 minutes. The reaction solution was partitioned with water and ethyl acetate. The organic layer was separated, washed  
5 with brine, and then, dried over anhydrous magnesium sulfate. The organic layer was filtered, then, the solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (heptane / ethyl acetate = 1 / 1), and the title compound (383mg, quantitatively) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>)  $\delta$ (ppm) : 4.92(2H, s), 5.10(2H, s), 6.94(1H, t, J=7.3Hz),  
10 7.00(2H, dd, J=0.92Hz, 8.8Hz), 7.29(2H, dd, J=7.2Hz, 8.8Hz), 7.44(1H, d, J=7.9Hz), 7.85(1H, dd, J=2.0Hz, 8.0Hz), 7.87-7.94(4H, m), 8.52(1H, d, J=1.8Hz).

Preparation Example 189. C-(5-Phenoxymethyl-pyridin-2-yl)-methylamine

2-(5-Phenoxymethyl-pyridin-2-ylmethyl)-isoindol-1,3-dione described in  
15 Preparation Example 188 (383mg, 1.11mmol) was dissolved in ethanol (3mL) and tetrahydrofuran (3mL). Hydrazine monohydrate (538 $\mu$ L, 11.1mmol) was added to this solution at room temperature, and the solution was stirred under reflux for 1 hour. The reaction solution was partitioned in water and ethyl acetate. The organic layer was separated, washed with brine, and then, dried over anhydrous magnesium  
20 sulfate. The organic layer was filtered, then, the solvent was evaporated *in vacuo*, the residue was purified by NH silica gel column chromatography (ethyl acetate / methanol = 10 / 1), and the title compound (122mg, 51%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>)  $\delta$ (ppm) : 4.00(2H, s), 5.07(2H, s), 6.67-7.01(3H, m),  
25 7.29-7.33(3H, m), 7.75(1H, dd, J=2.4Hz, 8.0Hz), 8.63(1H, d, J=1.6Hz).

Preparation Example 190. 4-(6-Fluoro-pyridin-2-yloxymethyl)-benzylamine

To a solution of 2,6-difluoropyridine (500mg, 4.34mmol) and 4-(hydroxymethyl)benzonitrile (867mg, 6.51mmol) in N,N-dimethylformamide (3mL) was added sodium hydride (0.26g, 6.51mmol; 60% in oil), and the solution was stirred at 70°C for 7 hours. The reaction solution was allowed to room temperature, then, partitioned in ethyl acetate and water, the organic layer was washed with water, and then, dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by silica gel column chromatography (hexane / ethyl acetate), and a white solid (734mg, 74%) was obtained.

To a solution of the resulting white solid (734mg, 3.22mmol) in tetrahydrofuran (5mL) was added lithium aluminum hydride (244mg, 6.44mmol), and the solution was stirred at room temperature for 30 minutes. The reaction solution was partitioned in ethyl acetate and water, and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated, and the title compound (662mg, 89%) was obtained as a crude product.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.71(2H, s), 5.27(2H, s), 6.71-6.73(1H, m), 6.79-6.81(1H, m), 7.33-7.39(4H, m), 7.86-7.90(1H, m).

Preparation Example 191. 4-(3-Chloro-benzyloxy)-benzylamine

To a solution of 4-cyanophenol (2.28g, 19.1mmol) and 3-chlorobenzyl bromide (2.2mL, 16.8mmol) in N,N-dimethylformamide (100mL) was added potassium carbonate (5.88g, 42.5mmol), and the solution was stirred at 50°C for 9 hours. The reaction mixture was partitioned in ethyl acetate and water. This organic layer was separated, washed with 2N sodium hydroxide, water and then

brine, and dried over anhydrous magnesium sulfate. The solvent was evaporated, and a crude product (4.20g, quantitatively) was obtained.

To a suspension of aluminum chloride in tetrahydrofuran (40mL) was added lithium aluminum hydride (4.70g, 35.2mmol) while cooling on a water bath. A solution of the crude product (1.15g, 4.71mmol) in tetrahydrofuran (10mL) was added to this suspension, which was then stirred at 0-1°C for 50 minutes. Concentrated aqueous ammonia (8mL) was added to the reaction mixture, ultrasound was applied, concentrated aqueous ammonia (8mL) was further added, and the solution was stirred for 1 hour at room temperature. This mixture was filtered through Celite pad, and this filtrate was separated. This filtrate was partitioned in tetrahydrofuran, ethyl acetate and water. This organic layer was separated, washed with water and brine, and dried over anhydrous sodium sulfate and anhydrous magnesium sulfate. The solvent was evaporated, and the title compound (1.15g, 99%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.03(2H, brs), 3.64(2H, s), 5.11(2H, s), 6.94(2H, d, J=8.0Hz), 7.24(2H, d, J=8.0Hz), 7.37-7.45(3H, m), 7.50(1H, s).

#### Preparation Example 192. 4-(3-Methoxy-benzyloxy)-benzylamine

To a solution of 4-cyanophenol (3.31g, 27.8mmol) and 3-methoxybenzyl bromide (3.7mL, 26.4mmol) in N,N-dimethylformamide (100mL) was added potassium carbonate (8.50g, 61.5mmol), and the solution was stirred at 50°C for 5 hours. The reaction mixture was partitioned in diethyl ether and water. This organic layer was separated, washed with 2N sodium hydroxide, water and brine, and dried over anhydrous magnesium sulfate. The solvent was evaporated, and a crude product (6.52g, 98%) was obtained.

To a solution of the resulting crude product (3.75g, 15.7mmol) in



tetrahydrofuran (80mL) was added lithium aluminum hydride (596mg, 15.7mmol), and the solution was stirred at room temperature for 23 hours. Sodium fluoride (6.6g) was added to the reaction mixture solution at room temperature, which was then cooled with an ice water bath, then, a mixture solution of water (2mL) and  
5 tetrahydrofuran (18mL) was added, followed by stirring. This mixture solution was filtered through Celite pad, and what was on Celite was washed with tetrahydrofuran and ethyl acetate. This filtrate was separated, the solvent was evaporated to obtain the title compound (3.84g, quantitatively) as a crude product.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.33(2H, brs), 3.63(2H, s), 3.76(3H, s), 5.06(2H, s),  
10 6.87-6.90(2H, d, J=8.4Hz), 6.93(2H, d, J=8.4Hz), 7.00-7.01(2H, m), 7.23(2H, d, J=8.4Hz), 7.28-7.32(1H, m).

Preparation Example 193. 4-(4-Methyl-pyridin-2-yloxymethyl)-benzylamine

To a solution of 2-fluoro-4-methylpyridine (500mg, 4.50mmol) and  
15 4-(hydroxymethyl)benzonitrile (899mg, 6.75mmol) in N,N-dimethylformamide (3mL) was added sodium hydride (0.27g, 6.75mmol; 60% in oil), and the solution was stirred at 70°C for 1 hour. The reaction solution was allowed to room temperature, then, partitioned in ethyl acetate and water, the organic layer was washed with water, and then, dried over anhydrous magnesium sulfate. The solvent was evaporated,  
20 the residue was purified by silica gel column chromatography (hexane / ethyl acetate), and a white solid (833mg, 83%) was obtained.

To a solution of the resulting white solid (200mg, 0.891mmol) in tetrahydrofuran (3mL) was lithium aluminum hydride (68mg, 1.78mmol), and the solution was stirred at room temperature for 1 hour. The reaction solution was  
25 partitioned in ethyl acetate and water, and the organic layer was dried over

anhydrous magnesium sulfate. The solvent was evaporated, and the title compound (181mg, 89%) was obtained as a crude product.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.77(3H, s), 3.69(2H, s), 5.29(2H, s), 6.68(1H, s), 6.82(1H, d, J=4.4Hz), 7.30-7.39(4H, m), 8.02(1H, d, J=5.6Hz).

5

Preparation Example 194. 4-(5-Methyl-pyridin-2-yloxymethyl)-benzylamine

To a solution of 2-fluoro-5-methylpyridine (1.0g, 9.0mmol) and 4-(hydroxymethyl)benzonitrile (1.8g, 13.5mmol) in N,N-dimethylformamide (5mL) was added sodium hydride (0.54mg, 13.5mmol; 60% in oil), and the solution was stirred  
10 for 30 minutes at 70°C. The reaction solution was allowed to room temperature, then, partitioned in ethyl acetate and water, the organic layer was washed with water, and then, dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by silica gel column chromatography (hexane / ethyl acetate), and a white solid (1.46g, 72%) was obtained.

15 To a solution of the resulting white solid (500mg, 2.23mmol) in tetrahydrofuran (5mL) was added lithium aluminum hydride (169mg, 4.46mmol), and the solution was stirred at room temperature for 30 minutes. The reaction solution was partitioned in ethyl acetate and water, and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated, and the title  
20 compound (457mg, 90%) was obtained as a crude product.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.20(3H, s), 3.69(2H, s), 5.27(2H, s), 6.76(1H, d, J=8.4Hz), 7.30-7.39(4H, m), 7.54(1H, d, J=7.2Hz), 7.97(1H, s).

Preparation Example 195. 1-Bromo-4-(2-propoxy-ethyl)-benzene

25 To a mixture of sodium hydride (66%, 360mg, 15mmol) and tetrahydrofuran

(10mL) was added 2-(4-bromophenyl)ethanol (1.5g, 7.5mmol) on an ice bath, the solution was stirred at room temperature for 1 hour. 1-Iodopropane (1.5mL, 15mmol) and N,N-dimethylformamide (10mL) were added to the reaction solution on an ice bath, and the solution was stirred overnight at 45°C. The reaction solution  
5 was partitioned with water (100mL) and heptane (200mL). The organic layer was separated, and washed with brine. this solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (heptane : ethyl acetate = 30 : 1), and the title compound (0.80g, 3.3mmol, 44%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.88-0.92(3H, m), 1.55-1.61(2H, m), 2.83(2H, t, J=7.0Hz), 3.36-3.40(2H, m), 3.60(2H, dt, J=1.5, 7.0Hz), 7.10(2H, d, J=8.1Hz),  
10 7.40(2H, d, J=8.2Hz).

Preparation Example 196. 4-(2-Propoxy-ethyl)-benzonitrile

1-Bromo-4-(2-propoxy-ethyl)-benzene described in Preparation Example 195  
15 (790mg, 3.2mmol), zinc cyanide (380mg, 3.2mmol) and tetrakis(triphenylphosphine)palladium (190mg, 0.16mmol) were added to N-methylpyrrolidinone (10mL), and this mixture was stirred at 125°C for 4 hours. This reaction mixture was allowed to cool, and water (50mL) and ethyl acetate (50mL) were added thereto. This mixture solution was filtered through Celite pad.  
20 The organic layer was separated, then, washed with water (3 times) and brine, and then, concentrated *in vacuo*. The residue was purified by silica gel column chromatography (heptane : ethyl acetate = 8 : 1), and the title compound (120mg, 0.62mmol, 19%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.89(3H, t, J=7.3Hz), 1.57(2H, tq, J=7.3, 7.3Hz),  
25 2.93(2H, t, J=6.6Hz), 3.38(2H, t, J=6.7Hz), 3.64(2H, t, J=6.6Hz), 7.34(2H, d,

J=8.1Hz), 7.58(2H, d, J=8.2Hz).

Preparation Example 197. 4-(2-Propoxy-ethyl)-benzylamine

To a mixture of lithium aluminum hydride (120mg, 2.5mmol) and  
 5 tetrahydrofuran (3mL) was added 4-(2-propoxy-ethyl)-benzonitrile described in  
 Preparation Example 196 (120mg, 0.62mmol), the solution was stirred overnight at  
 room temperature. The reaction solution was cooled to 0°C, tetrahydrofuran (30mL),  
 water (0.12mL), an aqueous solution of 5N sodium hydroxide (0.12mL) and water  
 (0.36mL) were sequentially added dropwise. After stirring for 1 hour at room  
 10 temperature, this reaction mixture was filtered through a filter paper. This filtrate  
 was concentrated *in vacuo*, the residue was filtered using NH-silica gel, and the title  
 compound (123mg, 0.64mmol, 103%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.91(3H, t, J=7.4Hz), 1.55-1.62(2H, m), 2.88(2H,  
 t, J=7.3Hz), 3.40(2H, t, J=6.7Hz), 3.63(2H, t, J=7.2Hz), 3.84(2H, s), 7.20(2H, d,  
 15 J=8.1Hz), 7.23(2H, d, J=8.2Hz).

Example A-1. 2,6-Diamino-N-(5-(4-fluoro-phenoxy)-furan-2-ylmethyl)-nicotinamide

2,6-Diamino-nicotinic acid described in Preparation Example A-15 (0.15g,  
 0.98mmol), triethylamine (0.41mL, 2.94mmol) and  
 20 benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (0.65g,  
 1.47mmol) were dissolved in N,N-dimethylformamide (5mL), and the solution was  
 stirred at room temperature for 10 minutes. Then, a solution of  
 C-(5-(4-fluoro-phenoxy)-furan-2-yl)methylamine described in Preparation Example  
 169 (304mg, 1.47mmol) in N,N-dimethylformamide (1mL) was added thereto,  
 25 followed by stirring at room temperature for 14 hours 50 minutes. After the reaction

was completed, reaction solution was poured into brine, the solution was extracted with ethyl acetate, the fractionated organic layer was dried over anhydrous magnesium sulfate and then concentrated. The obtained residue was subjected to silica gel column chromatography, eluted with solvent (ethyl acetate), and the title compound (0.12g, 0.35mmol, 36%) was obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.28 (2H, d, J=5.2 Hz), 5.64-5.69 (2H, m), 6.10 (2H, br s), 6.22 (1H, d, J=3.2 Hz), 6.96 (2H, br s), 7.08-7.14 (2H, m), 7.19-7.26 (2H, m), 7.63 (1H, d, J=8.8 Hz), 8.22 (1H, t, J=5.2 Hz)

10 Example A-2. 2,6-Diamino-N-(5-benzofuran-2-ylmethyl-furan-2-ylmethyl)-nicotinamide

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.18(2H, s), 4.31 (2H, d, J=5.6Hz), 5.65 (1H, dd, J=1.2, J=8.4Hz), 6.08 (2H, brs), 6.13-6.20 (2H, m), 6.65 (1H, s), 6.95 (2H, brs), 7.18-7.28 (2H, m), 7.48-7.53 (1H, m), 7.55 (1H, dd, J=0.8, J=5.6Hz), 7.63 (1H, d, J=8.4Hz), 8.23 (1H, t, J=5.6Hz).

Example A-3. 2-Amino-N-(5-(4-chloro-phenoxy)-furan-2-ylmethyl)nicotinamide

The title compound (55mg, 0.160mmol, 72.9%) was obtained as a brown oil from 2-aminonicotinic acid (34mg,0.24mmol) and C-(5-(4-chlorophenoxy)furan-2-yl)methylamine described in Preparation Example 47 (50mg,0.22mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.33 (2H, d, J=5.6Hz), 5.73-5.77 (1H, m), 6.29 (1H, d, J=3.2Hz), 6.54-6.58 (1H, m), 7.00-7.10 (4H, m), 7.38-7.45 (2H, m), 7.88 (1H, d, J=7.6Hz), 8.03-8.07 (1H, m), 8.85 (1H, t, J=5.6Hz).

Example A-4. 2-Amino-N-(5-(3-chloro-benzyl)-furan-2-ylmethyl)-nicotinamide

The title compound (110mg, 0.322mmol, 89.4%) was obtained as a white solid from 2-aminonicotinic acid (55mg, 0.39mmol) and C-(5-(3-chloro-benzyl)-furan-2-yl)-methylamine described in Preparation Example 56 (80mg, 0.36mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.94 (2H, s), 4.34 (2H, d, J=5.6Hz), 6.05 (1H, d, J=3.2Hz), 6.15 (1H, d, J=3.2Hz), 6.54 (1H, dd, J=4.0, 8.0Hz), 7.03 (2H, brs), 7.16-7.20 (1H, m), 7.24-7.35 (3H, m), 7.87 (1H, dd, J=1.6, 8.0Hz), 8.05 (1H, dd, J=1.6, 4.0Hz), 8.48 (1H, t, J=5.6Hz).

Example A-5. 2-Amino-N-(5-benzyl-furan-2-ylmethyl)-nicotinamide

The title compound (118mg, 0.384mmol, 24%) was obtained from C-(5-benzyl-furan-2-yl)-methylamine (360mg, 1.92mmol) prepared from 5-benzyl-furan-2-carbaldehyde according to an analogous method to Example Q-1, and 2-aminonicotinic acid (221mg, 1.60mmol) according to an analogous method to Example H-1.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.91(2H, s), 4.34 (2H, d, J=6.0Hz), 6.00 (1H, d, J=2.8Hz), 6.14 (1H, d, J=2.8Hz), 6.54 (1H, dd, J=4.8, 7.6Hz), 7.04 (2H, brs), 7.13-7.32 (5H, m), 7.87 (1H, dd, J=1.6, 7.6Hz), 8.05 (1H, d, J=1.6, 4.8Hz), 8.84 (1H, t, J=6.0Hz).

Example A-6. 2-Amino-N-(5-(3-fluoro-benzyl)-furan-2-ylmethyl)-nicotinamide

The title compound (252mg, 0.775mmol, 65%) was obtained from 2-aminonicotinic acid (164mg, 1.19mmol) and C-(5-(3-fluoro-benzyl)-furan-2-yl)-methylamine described in Preparation Example 84

(269mg, 1.31mmol) according to an analogous method to Example H-1.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.95 (2H, s), 4.34 (2H, d, J=5.6Hz), 6.04 (1H, d, J=3.2Hz), 6.15 (1H, d, J=3.2Hz), 6.54 (1H, dd, J=4.8, 7.6Hz), 6.97-7.12 (5H, m), 7.28-7.36 (1H, m), 7.87 (1H, dd, J=2.0, 7.6Hz), 8.04(1H, dd, J=2.0, 4.8Hz), 8.84 (1H, t, J=5.6Hz).

Example A-7. 2-Amino-N-(5-phenylaminomethyl-furan-2-ylmethyl)-nicotinamide

The title compound (49mg, 0.15mmol, 90%) was obtained as a white solid from (5-aminomethyl-furan-2-ylmethyl)-phenyl-amine described in Preparation Example 104 (34mg, 0.17mmol) and 2-amino-nicotinic acid (26mg, 0.19mmol) according to an analogous method to Example A-26.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.19 (2H, d, J=6.0Hz), 4.39 (2H, d, J=5.5Hz), 6.02 (1H, t, J=6.0Hz), 6.19 (2H, dd, J=3.1, 11Hz), 6.52 (1H, t, J=7.3Hz), 6.57 (1H, dd, J=4.8, 7.7Hz), 6.62 (2H, d, J=7.7Hz), 7.03-7.07 (4H, m), 7.91 (1H, dd, J=1.7, 7.7Hz), 8.07 (1H, dd, J=1.7, 4.8Hz), 8.88 (1H, t, J=5.5Hz).

Example A-8. 2-Amino-N-(5-(2-phenylamino-ethyl)-furan-2-ylmethyl)-nicotinamide

The title compound (29mg, 86μmol, 89%) was obtained as a white solid from 2-(5-aminomethyl-furan-2-yl)-ethyl)-phenylamine described in Preparation Example 107 (21mg, 97μmol) and 2-amino-nicotinic acid (16mg, 0.12mmol) according to an analogous method to Example A-26.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.83 (2H, t, J=7.3Hz), 3.26 (2H, m), 4.38 (2H, d, J=5.5Hz), 5.61 (1H, t, J=5.7Hz), 6.12 (1H, d, J=3.1Hz), 6.17 (1H, d, J=3.1Hz), 6.52 (1H, t, J=7.3Hz), 6.54-6.59 (3H, m), 7.04-7.08 (4H, m), 7.92 (1H, dd, J=1.8, 7.7Hz), 8.07 (1H, dd, J=1.8, 4.8Hz), 8.87 (1H, t, J=5.7Hz).

Example A-9. 6-Amino-N-(5-(3-fluoro-benzyl)-furan-2-ylmethyl)-nicotinamide

The title compound (265mg, 0.814mmol, 63%) was obtained from 6-aminonicotinic acid (180mg, 1.30mmol) and C-(5-(3-fluoro-benzyl)-furan-2-yl)-methylamine described in Preparation Example 84 (293mg, 1.43mmol) according to an analogous method to Example H-1.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.95 (2H, s), 4.33(2H, d, J=5.6Hz), 6.03 (1H, d, J=3.2Hz), 6.12 (1H, d, J=3.2Hz), 6.38 (1H, d, J=8.8Hz), 6.45 (2H, brs), 7.00-7.09 (3H, m), 7.28-7.36 (1H, m), 7.78 (1H, dd, J=2.4, 8.4Hz), 8.43 (1H, d, J=2.4Hz), 8.56 (1H, t, J=5.6Hz).

Example A-10. 2,6-Diamino-N-(4-benzyloxy-benzyl)-nicotinamide

2,6-Diamino-nicotinic acid described in Preparation Example A-15 (0.6g, 3.92mmol), triethylamine (1.64mL, 11.8mmol) and benzotriazole-1-yloxy tris(dimethylamino)phosphonium hexafluorophosphate (2.6g, 5.9mmol) were dissolved in N,N-dimethylformamide (200mL), 4-benzyloxy-benzylamine described in Preparation Example 1 (1.25g, 5.9mmol) was added thereto, and the solution was stirred at room temperature for 16 hours. After the reaction was completed, the reaction solution was poured into brine, and the solution was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, then concentrated, the obtained residue was purified by NH silica gel column chromatography (ethyl acetate : hexane = 2 : 1, then ethyl acetate), the resulting solid was washed with solvent (chloroform : ethyl acetate = 2 : 1), and the title compound (0.37g, 1.1mmol, 27%) was obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.29 (2H, d, J=6.0Hz), 5.07 (2H, s), 5.66 (1H,



d, J=8.4Hz), 6.06 (2H, br s), 6.91-6.97 (4H, m), 7.19 (2H, d, J=8.8Hz), 7.29-7.45 (5H, m), 7.65 (1H, d, J=8.4Hz), 8.27 (1H, t, J=6.0Hz).

**Example A-11. 2,6-Diamino-N-(4-(2-fluoro-benzyloxy)-benzyl)-nicotinamide**

5           2,6-Diamino-nicotinic acid described in Preparation Example A-15 (200mg, 1.3mmol), triethylamine (0.54mL, 3.87mmol) and benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (862mg, 1.95mmol) were added to N,N-dimethylformamide (20mL), and the solution was stirred at room temperature for 20 minutes. Next, 10 4-(2-fluoro-benzyloxy)-benzylamine described in Preparation Example 154 (453mg, 1.96mmol) was added thereto, and the solution was stirred at room temperature for 14 hours. After the reaction was completed, the reaction solution was poured into brine, the solution was extracted with ethyl acetate, the fractionated organic layer was dried over anhydrous magnesium sulfate and then concentrated. The obtained 15 residue was subjected to silica gel column chromatography, eluted with solvent (ethyl acetate) and the title compound (147mg, 0.40mmol, 31%) was obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.30 (2H, d, J=6.0 Hz), 5.11 (2H, s), 5.67 (1H, d, J=8.8 Hz), 6.06 (2H, br s), 6.89-7.02 (4H, m), 7.16-7.28 (4H, m), 7.37-7.44 (1H, m), 7.51-7.56 (1H, m), 7.66 (1H, d, J=8.8 Hz), 8.28 (1H, t, J=6.0 Hz).

20

**Example A-12. 2,6-Diamino-N-(4-(pyridin-2-ylmethoxy)-benzyl)-nicotinamide**

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.29 (2H, d, J=6.0Hz), 5.15 (2H, s), 5.66 (1H, d, J=8.8 Hz), 6.06 (2H, br s), 6.91-6.98 (4H, m), 7.17-7.23 (2H, m), 7.31-7.35 (1H, m), 7.49 (1H, d, J=7.6 Hz), 7.65 (1H, d, J=8.8 Hz), 7.82 (1H, dt, J=2.0, 7.6 Hz), 8.28 (1H, 25 t, J=6.0 Hz), 8.55-8.58 (1H, m).

Example A-13. 2,6-Diamino-N-(4-phenoxyethyl-benzyl)-nicotinamide

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.51(2H, brs), 4.57(2H, d, J=5.6Hz), 5.06(2H, s),  
 5.77(1H, d, J=8.4Hz), 6.04(1H, brs), 6.45(2H, brs), 6.96-6.98(3H, m), 7.28-7.31(3H,  
 5 m), 7.34-7.43(4H, m).

Example A-14. 2,6-Diamino-N-(4-(thiophen-3-ylmethoxy)-benzyl)-nicotinamide

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.28 (2H, d, J=6.0 Hz), 5.04 (2H, s), 5.65 (1H,  
 d, J=8.4 Hz), 6.05 (2H, br s), 6.89-6.99 (4H, m), 7.13-7.21 (3H, m), 7.51-7.55 (2H, m),  
 10 7.64 (1H, d, J=8.4 Hz), 8.26 (1H, t, J=6.0 Hz).

Example A-15. 2-amino-N-(4-(2-nitro benzyloxy)-benzyl)-nicotinamide

To a solution of sodium  
 4-(((2-aminopyridin-3-carbonyl)-amino)-methyl)-phenolate described in Preparation  
 15 Example A+-1 (100mg, 0.377mmol) in N,N-dimethylformamide (2.5mL) was added  
 O-Nitrobenzyl chloride (65mg, 0.379mmol), and the solution was stirred at room  
 temperature for 2 hours. Ethyl acetate and water were added to the reaction  
 solution, which was then partitioned, the organic layer was washed with water and  
 dried over anhydrous magnesium sulfate. The residue was purified by silica gel  
 20 column chromatography (hexane : ethyl acetate), and the title compound (51mg,  
 37%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.55(2H, d, J=5.6Hz), 5.50(2H, s), 6.23(1H, brs),  
 6.39(2H, brs), 6.57-6.60(1H, m), 6.98(2H, d, J=8.6Hz), 7.30(2H, d, J=8.6Hz),  
 7.50-7.52(1H, m), 7.58-7.60(1H, m), 7.67-7.71(1H, m), 7.87-7.89(1H, m),  
 25 8.15-8.18(2H, m).

Example A-16. 2-Amino-N-(4-(2-amino-benzyloxy)-benzyl)-nicotinamide

2-Amino-N-(4-(2-nitrobenzyloxy)-benzyl)-nicotinamide described in Example A-15 was dissolved in a mixture solvent of ethanol-tetrahydrofuran-water (3:1:1.5), iron powder (4mg) and ammonium chloride (85mg) were added thereto, and the solution was stirred overnight under reflux. In addition, iron powder (10mg) and ammonium chloride (20mg) were added, and the solution was stirred under reflux for 2 hours. The reaction solution was allowed to room temperature, filtered through Celite pad to remove insoluble matter, and the filtrate was concentrated *in vacuo*.

The residue was purified by NH silica gel chromatography (hexane : ethyl acetate), and the title compound (9mg, 98%) was obtained as a pale brown solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.53(2H, d, J=5.6Hz), 5.04(2H, s), 6.24(1H, brs), 6.51(2H, brs), 6.57-6.60(1H, m), 6.72-6.79(2H, m), 6.99(2H, d, J=8.4Hz), 7.16-7.20(2H, m), 7.28(2H, d, J=8.4Hz), 7.58-7.60(1H, m), 8.13-8.14(1H, m).

Example A-17. 2-Amino-N-(4-benzyloxy-benzyl)-nicotinamide

The title compound (257mg, 0.771mmol, 72%) was obtained from 2-aminonicotinic acid (148mg, 1.07mmol) and 4-benzyloxy-benzylamine described in Preparation Example 1 (251mg, 1.18mmol) according to an analogous method to Example H-1.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.34 (2H, d, J=6.0Hz), 5.06 (2H, s), 6.55 (1H, dd, J=4.8, 7.6Hz), 6.94 (2H, d, J=8.0Hz), 7.03 (2H, brs), 7.20 (2H, d, J=8.0Hz), 7.29 (1H, t, J=6.4Hz), 7.36 (2H, dd, J=6.4, 6.8Hz), 7.41 (2H, d, J=6.8Hz), 7.90 (1H, d, J=7.6Hz), 8.05 (1H, d, J=4.8Hz), 8.88 (1H, t, J=6.0Hz).

Example A-18. 2-Amino-N-(3-phenoxy-benzyl)-nicotinamide

The title compound (87mg, 0.27mmol, 26%) was obtained from 2-aminonicotinic acid (144mg, 1.04mmol) and 3-phenoxy-benzylamine described in Preparation Example 4 (228mg, 1.15mmol) according to an analogous method to Example H-1.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.40 (2H, d, J=5.6Hz), 6.56 (1H, dd, J=4.8, 7.6Hz), 6.85 (1H, dd, J=1.2, 8.0Hz), 6.92-7.05 (5H, m), 7.06 (1H, d, J=7.6Hz), 7.11 (1H, dd, J=7.6, 8.0Hz), 7.29-7.40 (3H, m), 7.89 (1H, dd, J=2.0, 7.6Hz), 8.06 (1H, dd, J=2.0, 4.8Hz), 8.96 (1H, t, J=5.6Hz).

Example A-19. 2-Amino-N-(4-(3-fluoro-benzyloxy)-benzyl)-nicotinamide

The title compound (172mg, 0.489mmol, 40%) was obtained from 2-aminonicotinic acid (170mg, 1.23mmol) and 4-(3-fluoro-benzyloxy)-benzylamine described in Preparation Example 6 (312mg, 1.35mmol) according to an analogous method to Example H-1.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.35 (2H, d, J=6.0Hz), 5.10 (2H, s), 6.45-6.60 (1H, m), 6.85-7.46 (10H, m), 7.85-7.92 (1H, m), 8.03-8.07 (1H, m), 8.75-8.92 (1H, m).

Example A-20. 2-Amino-N-(4-(2-fluoro-benzyloxy)-benzyl)-nicotinamide

The title compound (67mg, 0.19mmol, 45%) was obtained from 2-aminonicotinic acid (58mg, 0.42mmol) and 4-(2-fluoro-benzyloxy)-benzylamine described in Preparation Example 154 (117mg, 0.506mmol) according to an analogous method to Example H-1 (with the proviso that only the reaction temperature was changed to 60°C).

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.34 (2H, d, J=6.0Hz), 5.10 (2H, s), 6.56 (1H,

dd, J=4.8, 7.6Hz), 6.90-7.00 (2H, m), 7.04 (2H, brs), 7.15-7.28 (4H, m), 7.35-7.44 (1H, m), 7.50-7.58 (1H, m), 7.90 (1H, dd, J=1.2, 7.6Hz), 8.05 (1H, dd, J=1.2, 4.8Hz), 8.86-8.95 (1H, m).

5 Example A-21. 2-Amino-N-(4-(4-fluoro-benzyloxy)-benzyl)-nicotinamide

The title compound (187mg, 0.532mmol, 96%) was obtained from 2-aminonicotinic acid (77mg, 0.56mmol) and 4-(4-fluoro-benzyloxy)-benzylamine described in Preparation Example 155 (155mg, 0.670mmol) according to an analogous method to Example H-1 (with the proviso that only the reaction  
10 temperature was changed to 60°C).

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.34 (2H, d, J=5.6Hz), 5.04 (2H, s), 6.56 (1H, dd, J=5.2, 8.0Hz), 6.94 (2H, d, J=8.4Hz), 7.03 (2H, brs), 7.12-7.25 (4H, m), 7.46 (2H, dd, J=6.0, 8.4Hz), 7.90 (1H, d, J=8.0Hz), 8.05 (1H, d, J=5.2Hz), 8.88 (1H, t, J=5.6Hz).

15

Example A-22. 2-Amino-N-(4-benzyloxy-benzyl)-thionicotinamide

A mixture of 2-amino-N-(4-benzyloxy-benzyl)-nicotinamide described in Example A-17 (220mg, 0.67mmol), 2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulphide (Lawesson's  
20 reagent) (670mg, 1.7mmol) and toluene (8mL) was stirred at 80°C for 15 minutes, then, refluxed for 45 minutes. After cooling, the precipitate was filtered, and the filtrate was evaporated *in vacuo*. Purification was carried out by NH silica gel column chromatography (ethyl acetate), and the title compound (28mg, 0.080mmol, 12%) was obtained as a pale yellow oil.

25 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.87 (2H, d, J=4.9Hz), 5.08 (2H, s), 5.88 (2H,

brs), 6.62 (1H, dd, J=4.9, 7.5Hz), 6.98 (2H, d, J=8.6Hz) , 7.30 (2H, d, J=8.6Hz) , 7.32-7.44 (6H, m), 7.71 (1H, brs), 8.07 (1H, dd, J=1.7, 4.9Hz).

Example A-23. 2-Amino-N-(3-(2-butyloxy)benzyl)-nicotinamide

5 Trifluoroacetic acid salt of the title compound (10mg, 0.024mmol, 49%) was obtained from 2-amino-N-(3-hydroxybenzyl)-nicotinamide described in Preparation Example A-17 (12mg, 0.050mmol) and 1-bromo-2-butyne (6.6mg, 0.050mmol) according to an analogous method to Example E-43.

MS m/e (ESI) 296.3 (MH<sup>+</sup>)

10

Example A-24. 2-Amino-N-(4-benzylamino-benzyl)-6-chloro-nicotinamide

To a solution of (4-aminomethylphenyl)-benzylamine described in Preparation Example 19 (369mg, 1.74mmol) and 2-amino-6-chloro-nicotinic acid (300mg, 1.74mmol) in N,N-dimethylformamide (10mL) were added  
 15 benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (924mg, 2.09mmol) and triethylamine (0.49mL, 3.48mmol), and the solution was stirred overnight at room temperature. Ethyl acetate and water were added to the reaction solution, which was then partitioned, the organic layer was washed with water, and then, dried over anhydrous magnesium sulfate. The solvent was evaporated, the  
 20 residue was purified by silica gel column chromatography (hexane : ethyl acetate), and the title compound (310mg, 49%) was obtained as a pale yellow solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.32-4.34(3H, m), 4.45(2H, d, J=5.6Hz), 6.07(1H, brs), 6.54-6.63(5H, m), 7.03(1H, dd, J=2.4, J=8.4Hz), 7.14(2H, d, J=8.4Hz), 7.35-7.36(4H, m), 7.48(1H, d, J=8.0Hz).

25

Example A-25. 2-Amino-6-chloro-N-(4-phenylamino-benzyl)-nicotinamide

To a solution of (4-aminomethyl-phenyl)-phenylamine described in Preparation Example 20 (345mg, 1.74mmol) and 2-amino-6-chloro-nicotinic acid (300mg, 1.74mmol) in N,N-dimethylformamide (10mL) were  
5 benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (924mg, 2.09mmol) and triethylamine (0.49mL, 3.48mmol), and the solution was stirred overnight at room temperature. Ethyl acetate and water were added to the reaction solution, which was then partitioned, the organic layer was washed with water, and then, dried over anhydrous magnesium sulfate. The solvent was evaporated, the  
10 residue was purified by silica gel column chromatography (hexane : ethyl acetate), and the title compound (360mg, 59%) was obtained as a pale yellow solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.51(2H, d, J=5.2Hz), 5.74(1H, s), 6.16(1H, brs), 6.57(1H, d, J=8.0Hz), 6.58(2H, s), 6.94-6.97(1H, m), 7.04-7.09(4H, m), 7.21-7.30(4H, m), 7.52(1H, d, J=8.0Hz).

15

Example A-26. 2-Amino-6-chloro-N-(4-phenylaminomethyl-benzyl)-nicotinamide

To a solution of (4-aminomethyl-benzyl)-phenylamine described in Preparation Example 21 (369mg, 1.74mmol) and 2-amino-6-chloro-nicotinic acid (300mg, 1.74mmol) in N,N-dimethylformamide (10mL) were added  
20 benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (924mg, 2.09mmol) and triethylamine (0.49mL, 3.48mmol), and the solution was stirred overnight at room temperature. Ethyl acetate and water were added to the reaction solution, which was then partitioned, the organic layer was washed with water, and then, dried over anhydrous magnesium sulfate. The solvent was evaporated, the  
25 residue was purified by silica gel column chromatography (hexane : ethyl acetate),

and the title compound (479mg, 75%) was obtained as a pale yellow solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.08(1H, brs), 4.31(2H, s), 4.57(2H, d, J=5.6Hz), 6.21(1H, brs), 6.57(1H, d, J=8.0Hz), 6.58(2H, s), 6.61-6.63(2H, m), 6.70-6.74(1H, m), 7.15-7.19(2H, m), 7.30(2H, d, J=8.0Hz), 7.37(2H, d, J=8.0Hz), 7.51(1H, d, J=8.0Hz).

5

**Example A-27. 2-Amino-6-chloro-N-(4-benzyloxy-benzyl)-nicotinamide**

2-Amino-6-chloro-nicotinic acid described in Preparation Example A-1 (220mg, 1.4mmol), triethylamine (0.47mL, 3.37mmol) and benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (764mg, 1.73mmol) were dissolved in N,N-dimethylformamide (3mL), 4-benzyloxy-benzylamine described in Preparation Example 1 (399mg, 1.87mmol) was added thereto, followed by stirring at room temperature for 17 hours 30 minutes. After the reaction was completed, the reaction solution was poured into brine, which was then extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, then concentrated, the obtained residue was purified by silica gel column chromatography (ethyl acetate : hexane = 1 : 1), and the title compound (115mg, 0.31mmol, 22%) was obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.35 (2H, d, J=6.0Hz), 5.08 (2H, s), 6.63 (1H, d, J=8.0Hz), 6.94-6.98 (2H, m), 7.23 (2H, d, J=8.4Hz), 7.29-7.45 (5H, m), 7.52 (2H, brs), 7.97 (1H, d, J=8.0Hz), 8.96 (1H, t, J=6.0Hz).

20

**Example A-28. 2-Amino-N-(4-benzyloxy-benzyl)-6-cyclopropylamino-nicotinamide**

2-Amino-6-chloro-N-(4-benzyloxy-benzyl)-nicotinamide described in Example A-27 (80mg, 0.22mmol) was dissolved in tetrahydrofuran (3mL), cyclopropylamine (0.3mL, 4.3mmol) was added thereto, and the solution was heated in a sealed tube

25



for 16 hours (oil bath temperature: 140°C). The reaction solution was allowed to room temperature, the solution was concentrated, the obtained residue was purified by NH silica gel column chromatography (ethyl acetate : hexane = 2 : 1), and the title compound (12mg, 0.031mmol, 14%) was obtained.

- 5 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.52-0.58 (2H, m), 0.74-0.81 (2H, m), 2.47-2.56 (1H, m), 4.45 (2H, d, J=5.2 Hz), 5.07 (2H, s), 5.02 (1H, brs), 5.96-6.01 (1H, m), 6.04 (1H, d, J=8.4 Hz), 6.39 (2H, brs), 6.95 (2H, d, J=8.8 Hz), 7.23-7.45 (8H, m).

Example A-29. 2-Amino-N-(4-benzyloxy-benzyl)-6-ethoxy-nicotinamide

- 10 Sodium hydride (70mg, 1.7mmol, 60% in oil), catalytic amount of copper(I) iodide, 2-amino-6-chloronicotinic acid described in Preparation Example A-4 (30mg, 0.17mmol) were added sequentially to ethanol (0.5mL), the solution was stirred at 110°C for 3 hours, then, stirred overnight at 80°C. After cooling to room temperature, water, diethyl ether and an aqueous solution of 29% ammonia were added to the
- 15 reaction solution, which was then partitioned, and the aqueous layer was neutralized with citric acid. Dichloromethane was added to the aqueous layer, the organic layer was partitioned, washed with brine, then, the solvent was evaporated *in vacuo*. Trifluoroacetic acid salt of the title compound (3.4mg, 0.0069mmol, 14%) was obtained from a portion (10mg) of the residue (35mg) and 4-benzyloxybenzylamine
- 20 (10mg, 0.047mmol) according to an analogous method to Example Q-6.
- MS m/e (ESI) 378.5 (MH<sup>+</sup>)

Example A-30. (6-amino-5-(4-benzyloxy-benzylcarbamoyl)-pyridin-2-ylamino)-acetic acid

- 25 Glycine (935mg, 12.5mmol) and 1,8-diazabicyclo[5,4,0]undec-7-ene (1.86mL,

12.5mmol) were added to 2-amino-N-(4-benzyloxy-benzyl)-6-chloro-nicotinamide described in Preparation Example A+-18 (454mg, 1.25mmol) under nitrogen atmosphere, and the solution was stirred for 6 hours at 130°C. Dimethylsulfoxide (35mL) was added to the reaction mixture, which was then filtered with  
 5 polytetrafluoroethylene membrane filter (Whatman Inc), the filtrate was purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), and the title compound (287mg, 0.551mmol, 44%) was obtained.

MS m/e (ESI) 406.91 (MH<sup>+</sup>)

10 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.82-4.00(2H, m), 4.29(2H, d, J=6.0Hz), 5.06(2H, s), 5.77-5.88(1H, m), 6.93(2H, d, J=8.8Hz), 7.18(2H, d, J=8.8Hz), 7.28-7.42(5H, m), 7.68-7.80(1H, m).

Example A-31. 2-Amino-6-methoxymethyl-N-(4-(pyridin-2-ylmethoxy)-benzyl)-

15 nicotinamide

To a solution of 2-amino-6-methoxymethyl-nicotinic acid described in Preparation Example A-11 (100mg, 0.55mmol) and 4-(pyridin-2-ylmethoxy)-benzylamine described in Preparation Example 171 (170mg, 0.82mmol) in N,N-dimethylformamide (5mL) were added  
 20 benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (290mg, 0.66mmol) and triethylamine (0.23mL, 1.7mmol), and the solution was stirred overnight at room temperature. Ethyl acetate and water were added to the reaction solution for extraction, the organic layer was washed with brine, then, the solvent was evaporated *in vacuo*. The residue was purified by silica gel column chromatography  
 25 (heptane : ethyl acetate = 1 : 2), and the title compound (150mg, 0.40mmol, 73%)

was obtained as a colorless solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.32(3H, s) , 4.27(2H, s) , 4.33(2H, d, J=5.9Hz) , 5.14(2H, s) , 6.58(1H, d, J=7.9Hz) , 6.96(2H, d, J=8.6Hz) , 7.10(2H, br s) , 7.22(2H, d, J=8.6Hz) , 7.31 (1H, ddd, J=7.5, 4.8, 1.1Hz) , 7.47 (1H, d, J=7.9Hz) , 7.80  
5 (1H, td, J=7.6, 1.8Hz) , 7.94 (1H, d, J=8.1Hz) , 8.54-8.56 (1H, m) , 8.87 (1H, t, J=5.9Hz).

Example A-32. 2-Amino-N-(4-(4-fluoro-benzyloxy)-benzyl)-6-methoxymethyl-nicotinamide

10 To a solution of 2-amino-6-methoxymethyl-nicotinic acid described in Preparation Example A-11 (10mg, 0.055mmol) and 4-(4-fluoro-benzyloxy)-benzylamine described in Preparation Example 155 (19mg, 0.082mmol) in N,N-dimethylformamide (1mL) were added benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (29mg,  
15 0.066mmol) and triethylamine (0.022mL, 0.16mmol), and the solution was stirred overnight at room temperature. Water was added to the reaction solution, the precipitated solid was filtered, and the title compound (13mg, 0.033mmol, 60%) was obtained as a colorless solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.32(3H, s) , 4.27(2H, s) , 4.33(2H, d, J=5.9Hz) , 5.04(2H, s) , 6.58(1H, d, J=7.9Hz) , 6.94(2H, d, J=8.8Hz) , 7.09(2H, br s) ,  
20 7.16-7.22(4H, m) , 7.46 (2H, dd, J=8.7, 5.6Hz) , 7.93 (1H, d, J=8.1Hz) , 8.85-8.88 (1H, m).

Example A-33. 2-Amino-N-(4-benzyloxy-benzyl)-6-methoxymethyl-nicotinamide

25 MS m/e (ESI) 378 (MH<sup>+</sup>)

Example A-34. 2-Amino-6-methoxymethyl-N-(4-(pyridin-2-yloxymethyl)-benzyl)-nicotinamide

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.46(3H, s), 4.40(2H, s), 4.60(2H, d, J=5.6Hz),  
5 5.38(2H, s), 6.25(1H, brs), 6.40(2H, brs), 6.68-6.70(1H, m), 6.77-6.81(1H, m),  
6.87-6.91(1H, m), 7.35-7.37(2H, m), 7.46-7.48(2H, m), 7.57-7.61(2H, m),  
8.16-8.18(1H, m).

Example A-35. 2-Amino-N-(4-(3-fluoro-benzyloxy)-benzyl)-6-methoxymethyl-nicotinamide

MS m/e (ESI) 396 (MH<sup>+</sup>)

Example A-36. 2-Amino-N-(4-benzyloxy-benzyl)-6-(3-methoxypropoxy)-nicotinamide

Trifluoroacetic acid salt of the title compound (0.65mg, 0.0012mmol, 2.4%)  
15 was obtained from 2-amino-6-chloronicotinic acid described in Preparation Example A-4 (8.6mg, 0.050mmol) and 3-methoxypropanol (0.5mL) according to an analogous method to Example A-29.

MS m/e (ESI) 422.5 (MH<sup>+</sup>)

Example A-37. 2-Amino-N-(4-benzyloxy-benzyl)-6-methyl-nicotinamide

Trifluoroacetic acid salt of the title compound (0.38mg, 0.00082mmol, 4.6%)  
was obtained from 2-amino-N-(4-benzyloxy-benzyl)-6-chloro-nicotinamide described  
in Preparation Example A+-18 (6.5mg, 0.018mmol) and methylmagnesium bromide  
(0.93M tetrahydrofuran solution, 0.12mL, 0.11mmol) according to an analogous  
25 method to Example E-40.

MS m/e (ESI) 348.5 (MH<sup>+</sup>)

Example A-38. 2-Amino-N-(4-benzyloxy-benzyl)-6-propoxy-nicotinamide

Trifluoroacetic acid salt of the title compound (1.5mg, 0.0030mmol, 5.9%)  
5 was obtained from 2-amino-6-chloronicotinic acid described in Preparation Example A-4 (8.6mg, 0.050mmol) and propanol (0.5mL) according to an analogous method to Example A-29.

MS m/e (ESI) 406.6 (MH<sup>+</sup>)

10 Example A-39. 6-Amino-N-(4-benzyloxybenzyl)-nicotinamide

Trifluoroacetic acid salt of the title compound (7.1mg, 0.016mmol, 32%) was obtained from 4-benzyloxybenzylamine described in Preparation Example 1 (11mg, 0.050mmol) and 6-aminonicotinic acid (6.9mg, 0.050mmol) according to an analogous method to Example Q-6.

15 MS m/e (ESI) 334.3 (MH<sup>+</sup>)

Example A-40. 6-Amino-N-(3-phenoxybenzyl)-nicotinamide

Trifluoroacetic acid salt of the title compound (16mg, 0.037mmol, 74%) was obtained from 3-phenoxybenzylamine described in Preparation Example 4 (10mg,  
20 0.050mmol) and 6-aminonicotinic acid (6.9mg, 0.050mmol) according to an analogous method to Example Q-6.

MS m/e (ESI) 320.2 (MH<sup>+</sup>)

Example A-41. 6-Chloro-N-(3-phenoxy-benzyl)-nicotinamide

25 The title compound (240mg, 0.71mmol, 61%) was obtained as a white solid

from 3-phenoxy-benzylamine described in Preparation Example 4 (230mg, 1.1mmol) and 6-chloronicotinic acid (180mg, 1.1mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.63 (2H, d, J=5.7Hz), 6.42 (1H, brs), 6.94 (1H, dd, J=1.8, 8.1Hz), 6.98-7.03 (3H, m), 7.08(1H, d, J=7.5Hz), 7.11-7.15 (1H, m), 7.30-7.37 (3H, m), 7.41-7.43 (1H, m), 8.09 (1H, dd, J=2.6, 8.2Hz), 8.74 (1H, d, J=2.2Hz).

Example A-42. N-(4-Benzyloxy-benzyl)-6-methylamino-nicotinamide

10 The title compound (71mg, 0.19mmol, 88%) was obtained as a white solid from N-(4-benzyloxy-benzyl)-6-(ethoxymethyl-amino)-nicotinamide described in Preparation Example A+-8 (90mg, 0.22mmol) according to an analogous method to Example A-163.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 2.96 (3H, s), 4.56 (2H, d, J=5.5Hz), 5.07 (2H, s), 15 6.12 (1H, brs), 6.38 (1H, d, J=8.8Hz), 6.96 (2H, d, J=8.8Hz), 7.22-7.34 (4H, m), 7.36-7.44 (3H, m), 7.89 (1H, dd, J=2.4, 8.6Hz), 8.50 (1H, d, J=2.4Hz).

Example A-43. N-(4-benzyloxybenzyl)-nicotinamide

20 The title compound (8.5mg, 0.027mmol, 33%) was obtained as a white solid from 4-benzyloxybenzylamine described in Preparation Example 1 (26mg, 0.12mmol) and nicotinic acid (10mg, 0.081mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.59 (2H, d, J=5.5Hz), 5.07 (2H, s), 6.41 (1H, brs), 6.97 (2H, d, J=8.6Hz), 7.29 (2H, d, J=8.6Hz), 7.32-7.44 (6H, m), 8.11-8.14 (1H, 25 m), 8.71 (1H, dd, J=1.5, 4.8Hz), 8.96 (1H, d, J=1.8Hz).

Example A-44. 2-Amino-N-(4-benzyloxy-3-hydroxy-benzyl)-nicotinamide

To a solution of 4-benzyloxy-3-methoxymethoxy-benzonitrile obtained in Preparation Example 132 (100mg, 0.371mmol) in tetrahydrofuran (5mL) was added  
5 lithium aluminum hydride (75mg, 1.98mmol) portionwise under nitrogen atmosphere on an ice bath, the solution was stirred for 24 hours. In addition, lithium aluminum hydride (75mg, 1.98mmol) was added portionwise on an ice bath, then, the solution was stirred at 50-60°C for 3 hours. Ethyl acetate (10mL) and methanol (5mL) were added little by little to the reaction solution on an ice bath, then, NH silica gel (50mL)  
10 was added, and a pale yellow oily residue (73mg) was obtained by NH silica gel column chromatography (hexane : ethyl acetate = 7 : 3). This residue was purified again by NH silica gel column chromatography (hexane : ethyl acetate = 7 : 3), and 4-benzyloxy-3-methoxymethoxy-benzylamine (30mg, 0.11mmol, 30%) was obtained as a pale yellow oil.

15 Next, a solution of 2-amino-nicotinic acid (16mg, 0.116mmol), 4-benzyloxy-3-methoxymethoxy-benzylamine (15mg, 0.0549mmol), benzotriazol-1-yl-tris(dimethylamino)phosphonium hexafluorophosphate (55mg, 0.124mmol) and triethylamine (0.08mL, 0.574mmol) in dimethylsulfoxide (4mL) was stirred under nitrogen atmosphere at room temperature for 24 hours. Water  
20 (100mL) and brine (50mL) were added to the reaction solution, which was then extracted with ethyl acetate (100mL) twice, and washed with water twice. The organic layer was dried over anhydrous magnesium sulfate, which was then filtered, the filtrate was evaporated *in vacuo*, and 2-amino-N-(4-benzyloxy-3-methoxymethoxy-benzyl)-nicotinamide was obtained as a  
25 pale yellow oil. This was purified by NH silica gel column chromatography (hexane :

ethyl acetate = 1 : 1, then hexane : ethyl acetate = 3 : 7), and  
 2-amino-N-(4-benzyloxy-3-methoxymethoxy-benzyl)-nicotinamide (9.4mg,  
 0.0239mmol, 43.5%) was obtained as a pale yellow oil. A solution of the resulting  
 2-amino-N-(4-benzyloxy-3-methoxymethoxy-benzyl)-nicotinamide (7.8mg,  
 5 0.0198mmol) and 2M hydrochloric acid (2mL) in methanol (3mL) was stirred for 21  
 hour at room temperature. Sodium bicarbonate (600mg, 7.14mmol) was added to  
 the reaction mixture to basify it, which was then filtered, evaporation *in vacuo* was  
 carried out, then, the obtained residue was purified by thin layer NH silica gel  
 chromatography (methanol : ethyl acetate = 5 : 95), and the title compound (2.0mg,  
 10 0.0057mmol, 29%) was obtained as a white solid.  
 MS m/e (ESI) 350(MH<sup>+</sup>)

ExampleA-45.2-Amino-N-(6-benzyloxypyridin-3-ylmethyl)-6-methoxymethyl-nicotinamide

15 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.45(3H, s), 4.39(2H, s), 4.53(2H, d, J=5.6Hz),  
 5.38(2H, s), 6.23(1H, brs), 6.39(2H, brs), 6.69-6.71(1H, m), 6.80-6.82(1H, m),  
 7.30-7.33(1H, m), 7.36-7.40(2H, m), 7.57-7.59(2H, m), 7.60-7.63(2H, m),  
 8.15-8.15(1H, m).

20 Example A-46. 2,6-Diamino-N-(1-(3-fluoro-benzyl)-1H-pyrrol-3-ylmethyl)-nicotinamide

The title compound (6.2mg, 0.018mmol, 5.5%) was obtained from  
 2,6-diamino-nicotinic acid ethyl ester described in Preparation Example A-14 (60mg,  
 0.33mmol) and C-(1-(3-fluoro-benzyl)-1H-pyrrol-3-yl)-methylaniline described in  
 Preparation Example 59 (159mg, 0.78mmol) according to an analogous method to  
 25 Example A-54.



<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.40 (2H, d, J=4.8 Hz), 4.53 (2H, brs), 5.01 (2H, s), 5.76 (1H, d, J=8.4 Hz), 5.84-5.89 (1H, m), 6.17 (1H, t, J=2.0 Hz), 6.48 (2H, brs), 6.62-6.67 (2H, m), 6.78-6.82 (1H, m), 6.91-6.93 (1H, m), 6.98 (1H, dt, J=2.4, 8.4 Hz), 7.27-7.33 (1H, m), 7.36 (1H, d, J=8.4 Hz).

5

Example A-47. 2-Amino-N-(1-(3-fluoro-benzyl)-1*H*-pyrrol-3-ylmethyl)nicotinamide

The title compound (106mg, 0.327mmol, 66.7%) was obtained as a white solid from C-(1-(3-fluoro-benzyl)-1*H*-pyrrol-3-yl)methylamine described in Preparation Example 59 (100mg, 0.49mmol) and 2-aminonicotinic acid (68mg, 0.49mmol) according to an analogous method to Example Q-6.

10

<sup>1</sup>H-NMR Spectrum (DMSO-*d*<sub>6</sub>) δ(ppm) : 4.22 (2H, d, J=5.6Hz), 5.03 (2H, s), 5.97-6.01 (1H, m), 6.50-6.55 (1H, m), 6.73 (2H, s), 6.95-7.10 (5H, m), 7.32-7.38 (1H, m), 7.83-7.88 (1H, m), 8.00-8.05 (1H, m), 8.63 (1H, t, J=5.6Hz).

15

Example A-48.

2-Amino-N-(1-(3-fluoro-benzyl)-1*H*-pyrrol-3-ylmethyl)-6-methylamino-nicotinamide

2-Amino-6-chloro-N-(1-(3-fluoro-benzyl)-1*H*-pyrrol-3-ylmethyl)-nicotinamide described in Preparation Example A+-5 (50mg, 0.14mmol) was dissolved in a mixture solution of dimethylsulfoxide (1mL) and N,N-diisopropylethylamine (0.5mL), methylamine (2.0M tetrahydrofuran solution) (1mL, 2mmol) was added thereto, and the solution was heated in a sealed tube for 15 hours (oil bath temperature: 135°C). The reaction mixture was allowed to room temperature, poured into brine, and the solution was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate then concentrated, the obtained residue was purified by NH silica gel column chromatography (ethyl acetate), and the title compound

20

25

(7.3mg, 0.021mmol, 15%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 2.88 (3H, d, J=5.2 Hz), 4.40 (2H, d, J=5.2 Hz), 4.58-4.66 (1H, m), 5.01 (2H, s), 5.66 (1H, J=8.8 Hz), 5.81-5.87 (1H, m), 6.17 (1H, t, J=2.4 Hz), 6.45 (2H, brs), 6.63 (1H, t, J=2.4 Hz), 6.66 (1H, brs), 6.78-6.83 (1H, m), 6.92 (1H, br d, J=7.2 Hz), 6.97 (1H, dt, J=2.4, 8.4 Hz), 7.27-7.33 (1H, m), 7.36 (1H, d, J=8.8 Hz).

Example A-49. N-(1-(3-Fluoro-benzyl)-1H-pyrrol-3-ylmethyl)-6-methyl-nicotinamide

The title compound (61mg, 0.18mmol, 65.1%) was obtained as a colorless oil from C-(1-(3-fluoro-benzyl)-1H-pyrrol-3-yl)methylamine described in Preparation Example 59 (60mg, 0.29mmol) and 6-methylnicotinic acid (40mg, 0.29mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.48(3H, s), 4.27 (2H, d, 5.6Hz), 5.03 (2H, s), 5.99-6.02 (1H, m), 6.74-6.78 (2H, m), 6.96-7.10 (3H, m), 7.28-7.38 (2H, m), 8.06 (1H, dd, J=2.4, 8.0Hz), 8.80 (1H, t, J=5.6Hz), 8.87 (1H, d, J=2.4Hz).

Example A-50. 2-Amino-N-(2-phenoxy-thiazol-5-ylmethyl)-nicotinamide

The title compound (13.5mg, 41μmol, 87%) was obtained as a white solid from C-(2-phenoxy-thiazol-5-yl)-methylamine described in Preparation Example 117 (9.8mg, 48μmol) and 2-amino-nicotinic acid (7.9mg, 58μmol) according to an analogous method to Example A-26.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.44 (2H, d, J=5.7Hz), 6.55 (1H, dd, J=4.8, 7.7Hz), 7.05 (2H, s), 7.18 (1H, s), 7.27-7.31 (3H, m), 7.45 (2H, t, J=8.2Hz), 7.84 (1H, d, J=7.5Hz), 8.06 (1H, d, J=4.6Hz), 9.05 (1H, t, J=6.0Hz).

ExampleA-51.2-((furan-2-ylmethyl)-amino)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

The title compound (2.29mg, 0.0044mmol, 4.4%) was obtained as a trifluoroacetic acid salt from 2-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Preparation Example A+-7 (35mg, 0.10mmol) and furfurylamine (16μl, 0.18mmol) according to an analogous method to Example A-133.

MS m/e (ESI) 406.15(MH<sup>+</sup>)Example A-52. 4-((3-((5-Phenoxy-thiophen-2-ylmethyl)-carbamoyl)-pyridin-2-yl)amino)-methyl) benzoic acid

The title compound (2.75mg, 0.0048mmol, 4.8%) was obtained as a trifluoroacetic acid salt from 2-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Preparation Example A+-7 (36mg, 0.10mmol) and 4-(aminomethyl)benzoic acid (16mg, 0.11mmol) according to an analogous method to Example A-133.

MS m/e (ESI) 460.17(MH<sup>+</sup>)

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.50 (2H, d, J=6.0Hz), 4.71 (2H, d, J=4.8Hz), 6.51(1H, d, J=3.6Hz), 6.63 (1H, dd, J=4.8, 8.0Hz), 6.79 (1H, d, J=3.6Hz), 7.09 (2H, d, J=8.0Hz), 7.09-7.19 (1H, m), 7.32-7.47 (4H, m), 7.87 (2H, d, J=7.6Hz), 7.98 (1H, dd, J=0.8, 8.0Hz), 8.13 (1H, dd, J=0.8, 4.8Hz), 8.87 (1H, brs), 9.16-9.24 (1H, m).

ExampleA-53.2,6-Diamino-N-(5-(4-fluoro-phenoxy)-thiophen-2-ylmethyl)-nicotinamide

The title compound (180mg, 0.50mmol, 38.6%) was obtained as a white solid from C-(5-(4-fluoro-phenoxy)-thiophen-2-yl)-methylamine described in Preparation

Example 28 (290mg, 1.3mmol) and 2,6-diaminonicotinic acid described in Preparation Example A-15 (200mg, 1.3mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.40 (2H, d, J=5.6Hz), 5.64-5.68 (1H, m),  
5 6.10 (2H, s), 6.45-6.49 (1H, m), 6.70-6.74 (1H, m), 6.95 (2H, s), 7.10-7.18 (2H, m),  
7.18-7.26 (2H, m), 7.60 (1H, d, J=8.0Hz), 8.40 (1H, t, J=5.6Hz).

Example A-54. 2,6-diamino-N-(5-phenoxy-thiophene-2-ylmethyl)-nicotinamide

To a solution of 2,6-diamino-nicotinic acid ethyl ester described in Preparation  
10 Example A-14 (18mg, 0.1mmol) in ethanol (10mL) was added 1N sodium hydroxide  
aqueous solution (5mL), and the solution was stirred for 1 hour 10 minutes under  
reflux. After cooling the reaction solution, the solution was neutralized with 1N  
hydrochloric acid and concentrated. The resulting crude product was suspended in  
N,N-dimethylformamide (3mL), triethylamine (0.02mL, 0.15mmol),  
15 benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (65mg,  
0.15mmol) and C-(5-phenoxy-thiophen-2-yl)-methylamine described in Preparation  
Example 26 (30mg, 0.15mmol) were added thereto, followed by stirring at room  
temperature for 19 hours 40 minutes. After the reaction was completed, reaction  
solution was poured into brine, and the solution was extracted with ethyl acetate.  
20 The organic layer was dried over anhydrous magnesium sulfate then concentrated,  
the obtained residue was purified by NH silica gel column chromatography (ethyl  
acetate), and the title compound (8.7mg, 0.025mol, 25%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.58-4.61 (2H, m), 4.62 (2H, brs), 5.75 (1H, d,  
J=8.4Hz), 6.21-6.27 (1H, m), 6.36-6.38 (1H, m), 6.45 (2H, brs), 6.69-6.72 (1H, m),  
25 7.06-7.12 (3H, m), 7.28-7.34 (2H, m), 7.39 (1H, d, J=8.4Hz).

Example A-55. 2,6-Diamino-N-(5-benzyl-thiophen-2-ylmethyl)-nicotinamide

To a solution of 2,6-diaminonicotinic acid described in Preparation Example A-15 (173mg, 1.13mmol) in dimethylsulfoxide (15mL) were added  
 5 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (433mg, 2.26mmol),  
 1-hydroxybenzotriazole (346mg, 2.26mmol) and  
 C-(5-benzyl-thiophen-2-yl)-methylamine described in Preparation Example 42 (230mg, 1.13mmol), and the solution was stirred at room temperature for 16 hours 30  
 minutes. The reaction solution was poured into brine, the solution was extracted  
 10 with ethyl acetate, the organic layer was dried over anhydrous magnesium sulfate  
 and then concentrated. The obtained residue was subjected to NH silica gel column  
 chromatography, eluted with solvent (ethyl acetate) and the title compound (114mg,  
 0.34mmol, 30%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.00 (2H, s), 4.53 (2H, d, J=5.6 Hz), 4.70 (2H, br  
 15 s), 5.69 (1H, d, J=8.4 Hz), 6.39 (2H, br s), 6.55 (1H, d, J=3.2 Hz), 6.67-6.95 (2H, m),  
 7.09-7.25 (5H, m), 7.38 (1H, d, J=8.4 Hz).

Example A-56. 2,6-Diamino-N-(5-benzyloxy-thiophen-2-ylmethyl)-nicotinamide

To a solution of 2,6-diaminonicotinic acid described in Preparation Example  
 20 A-15 (109mg, 0.71mmol) in dimethylsulfoxide (10mL) were added  
 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (272mg, 1.42mmol),  
 1-hydroxybenzotriazole (217mg, 1.42mmol) and  
 C-(5-benzyloxy-thiophen-2-yl)methylamine described in Example E-76 (156mg,  
 0.71mmol), and the solution was stirred at room temperature for 14 hours. The  
 25 reaction solution was poured into brine, extracted with ethyl acetate, the fractionated

organic layer was dried over anhydrous magnesium sulfate and then concentrated. The obtained residue was subjected to NH silica gel column chromatography, eluted with solvent (ethyl acetate) and the title compound (157mg, 0.44mmol, 62%) was obtained.

- 5  $^1\text{H-NMR}$  Spectrum ( $\text{CDCl}_3$ )  $\delta$ (ppm) : 4.53 (2H, d,  $J=5.2$  Hz), 4.69 (2H, br s), 5.03 (2H, s), 5.76 (1H, d,  $J=8.8$  Hz), 6.08 (1H, d,  $J=4.0$  Hz), 6.46 (3H, br), 6.59 (1H, d,  $J=4.0$  Hz), 7.31-7.44 (6H, m).

Example A-57.

10 2,6-Diamino-N-(5-(3-fluoro-phenoxy)-thiophen-2-ylmethyl)-nicotinamide

$^1\text{H-NMR}$  Spectrum ( $\text{CDCl}_3$ )  $\delta$ (ppm) : 4.42 (2H, d,  $J=6.0\text{Hz}$ ), 5.66 (1H, d,  $J=8.4\text{Hz}$ ), 6.11 (2H, brs), 6.55-6.59 (1H, m), 6.75 (1H, d,  $J=4.0\text{Hz}$ ), 6.90-7.02 (5H, m), 7.38-7.45 (1H, m), 7.60 (1H, d,  $J=8.4\text{Hz}$ ), 8.42 (1H, t,  $J=6.0\text{Hz}$ ).

15 Example A-58.

2,6-Diamino-N-(5-benzofuran-2-ylmethyl-thiophen-2-ylmethyl)-nicotinamide

$^1\text{H-NMR}$  Spectrum ( $\text{CDCl}_3$ )  $\delta$ (ppm) : 4.29 (2H, s), 4.43 (2H, d,  $J=6.0\text{Hz}$ ), 5.64 (1H, d,  $J=8.4\text{Hz}$ ), 6.09 (2H, brs), 6.66 (1H, d,  $J=0.8\text{Hz}$ ), 6.92 (2H, s), 6.94 (2H, brs), 7.17-7.26 (2H, m), 7.47-7.64 (3H, m), 8.37 (1H, t,  $J=6.0\text{Hz}$ ).

20

Example A-59.

2,6-Diamino-N-(5-benzo[b]thiophen-2-ylmethyl-thiophen-2-ylmethyl)-nicotinamide

- $^1\text{H-NMR}$  Spectrum ( $\text{CDCl}_3$ )  $\delta$ (ppm) : 4.39 (2H, s), 4.44 (2H, d,  $J=5.6\text{Hz}$ ), 5.64 (1H, d,  $J=8.4\text{Hz}$ ), 6.09 (2H, brs), 6.78-6.84 (2H, m), 6.95 (2H, brs), 7.23-7.36 (3H, m), 7.59 (1H, d,  $J=8.4\text{Hz}$ ), 7.74 (1H, dd,  $J=0.8$ ,  $J=6.0\text{Hz}$ ), 7.85 (1H, dd,  $J=0.8$ ,  $J=6.0\text{Hz}$ ),
- 25

8.38(1H, t, J=5.6Hz).

Example A-60. N-(5-(3-Fluorophenoxy)thiophen-2-ylmethyl)-2,6-dimethylnicotinamide

The title compound (56mg, 0.157mmol, 47.6%) was obtained as a light brown solid from 2,6-dimethylnicotinic acid (50mg, 0.33mmol) and C-(5-(3-fluorophenoxy)thiophen-2-yl)methylamine described in Preparation Example 23 (66mg, 0.297mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.44 (3H, s), 2.46 (3H, s), 4.51 (2H, d, J=5.6Hz), 6.59-6.62 (1H, m), 6.80-6.84 (1H, m), 6.90-7.03 (3H, m), 7.12 (1H, d, J=8.0Hz), 7.40-7.47 (1H, m), 7.60 (1H, d, J=8.0Hz), 9.00 (1H, t, J=5.6Hz).

Example A-61. 2-Acetylamino-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

To a mixture of 2-amino-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-67 (50mg, 0.15mmol), acetonitrile(5mL) and tetrahydrofuran (2mL) was added nitronium tetrafluoroborate (0.50M sulfolane solution, 0.46mL, 0.23mmol) on an ice bath, and the solution was stirred overnight at room temperature. Water, ethyl acetate, tetrahydrofuran and an aqueous solution of saturated sodium bicarbonate were added to the reaction solution for extraction, washed with brine, then, the solvent was evaporated *in vacuo*. The residue was purified by silica gel column chromatography (methanol : ethyl acetate = 1 : 30), and the title compound (1.3mg, 0.0035mmol, 2.3%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 2.31 (3H, s), 4.67 (2H, d, J=5.5Hz), 6.40 (1H, d, J=3.9Hz), 6.60 (1H, brs), 6.77 (1H, d, J=3.7Hz), 7.05 (1H, dd, J=4.9, 7.8Hz), 7.08-7.14 (3H, m), 7.32-7.36 (2H, m), 7.84 (1H, brs), 8.51 (1H, brs).

Example A-62. 2-Amino-N-(5-(3-cyano-phenoxy)-thiophen-2-ylmethyl)-nicotinamide

To a solution of C-(5-(3-bromophenoxy)-thiophen-2-yl)-methylaniline described in Preparation Example 17 (366mg, 1.29mmol) and 2-aminopyridine-3-carboxylic acid (178mg, 1.29mmol) in tetrahydrofuran (5mL) were added benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (685mg, 1.55mmol) and triethylamine (0.36mL, 2.58mmol), and the solution was stirred overnight at room temperature. Ethyl acetate and water were added to the reaction solution, which was then partitioned, the organic layer was washed with water, and then, dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate), and a mixture of 2-amino-N-(5-(3-bromophenoxy)-thiophen-2-ylmethyl)-nicotinamide and debrominated compound (344mg, 66%) was obtained as a yellow solid.

Next, to a solution of a mixture of 2-amino-N-(5-(3-bromophenoxy)-thiophen-2-ylmethyl)-nicotinamide and debrominated compound (100mg, 0.247mmol) in N,N-dimethylformamide (3.0mL) were added zinc cyanide (58mg, 0.495mmol) and tetrakis(triphenylphosphine)palladium(0) (285mg, 0.247mmol) under nitrogen atmosphere, and the solution was stirred at 140°C for 3 hours and a half. The reaction solution was allowed to room temperature, ethyl acetate and water were added to the reaction solution, which was then partitioned, the organic layer was washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated, then, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate), and the title compound (9mg, 10%) was obtained as a pale yellow oil.



<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.69(2H, d, J=6.0Hz), 6.36(2H, brs), 6.41(1H, brs), 6.47-6.48(1H, m), 6.59-6.62(1H, m), 6.79-6.80(1H, m), 7.31-7.34(2H, m), 7.37-7.44(2H, m), 7.61-7.63(1H, m), 8.17-8.19(1H, m).

5 Example A-64. 2-Amino-N-(5-*m*-tolxyloxy-thiophen-2-ylmethyl)-nicotinamide

The title compound (126mg, 0.37mmol, 80.8%) was obtained as a brown oil from 2-aminonicotinic acid (69mg, 0.51mmol) and C-(5-*m*-tolxyloxy-thiophen-2-yl)-methylamine described in Preparation Example 30 (100mg, 0.46mmol) according to an analogous method to Example Q-6.

10 <sup>1</sup>H-NMR Spectrum (DMSO-*d*<sub>6</sub>) δ(ppm) : 2.26 (3H, s), 4.46 (2H, d, J=5.6Hz), 6.46 (1H, d, J=3.6Hz), 6.55 (1H, dd, J=4.8, 8.0Hz), 6.74 (1H, d, J=3.6Hz), 6.84-6.96 (3H, m), 7.04 (2H, s), 7.22 (1H, dd, J=8.0, 8.0Hz), 7.86 (1H, dd, J=1.6, 8.0Hz), 8.05 (1H, dd, J=1.6, 4.8Hz), 9.02 (1H, t, J=5.6Hz).

15 Example A-65. 2-Amino-N-(5-*p*-tolxyloxy-thiophen-2-ylmethyl)nicotinamide

The title compound (72mg, 0.212mmol, 57.4%) was obtained as a light brown solid from 2-aminonicotinic acid (55mg, 0.41mmol) and C-(5-*p*-tolxyloxythiophen-2-yl)methylamine described in Preparation Example 32 (80mg, 0.37mmol) according to an analogous method to Example Q-6.

20 <sup>1</sup>H-NMR Spectrum (DMSO-*d*<sub>6</sub>) δ(ppm) : 2.24 (3H, s), 4.44 (2H, d, J=5.6Hz), 6.40-6.44 (1H, m), 6.53-6.58 (1H, m), 6.72 (1H, d, J=3.6Hz), 6.95-7.00 (2H, m), 7.04 (2H, s), 7.12-7.18 (2H, m), 7.85 (1H, d, J=7.6Hz), 8.03-8.08 (1H, m), 9.01 (1H, t, J=5.6Hz).

Example A-66. 2-amino-4-(5-(3-chloro-benzyl)thiophene-2-ylmethyl)-nicotinamide

25 The title compound (63mg, 0.176mmol, 51.9%) was obtained as a white solid

from 2-aminonicotinic acid (51mg, 0.37mmol) and C-(5-(3-chloro-benzyl)thiophen-2-yl)-methylamine described in Preparation Example 45 (80mg, 0.34mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.06 (2H, s), 4.47 (2H, d, J=5.6Hz), 6.54 (1H, dd, J=4.8, 7.6Hz), 6.72 (1H, d, J=3.6Hz), 6.80 (1H, d, J=3.6Hz), 7.05 (2H, brs), 7.18-7.34 (4H, m), 7.84 (1H, dd, J=1.6, 7.6Hz), 8.04 (1H, dd, J=1.6, 4.8Hz), 8.98 (1H, t, J=5.6Hz).

Example A-67. 2-Amino-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

The title compound (148mg, 0.455mmol, 73%) was obtained from 2-aminonicotinic acid (87mg, 0.63mmol) and C-(5-phenoxy-thiophen-2-yl)methylamine described in Preparation Example 24 (143mg, 0.697mmol) according to an analogous method to Example H-1.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.46 (2H, d, J=5.6Hz), 6.49 (1H, d, J=4.0Hz), 6.55 (1H, dd, J=4.8, 7.6Hz), 6.76 (1H, d, J=4.0Hz), 7.00-7.17 (5H, m), 7.32-7.40 (2H, m), 7.87 (1H, dd, J=1.6, 7.6Hz), 8.06 (1H, dd, J=1.6, 4.8Hz), 9.00-9.09 (1H, m).

Example A-68. 2-Amino-N-(5-(3-fluoro-phenoxy)-thiophen-2-ylmethyl)-nicotinamide

The title compound (112mg, 0.326mmol, 75%) was obtained from 2-aminonicotinic acid (60mg, 0.43mmol) and C-(5-(3-fluoro-phenoxy)-thiophen-2-yl)methylamine described in Preparation Example 23 (106mg, 0.475mmol) according to an analogous method to Example H-1.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.48 (2H, d, J=5.2Hz), 6.52-6.59 (2H, m), 6.79 (1H, d, J=2.8Hz), 6.87-7.00 (3H, m), 7.06 (2H, brs), 7.34-7.45 (1H, m), 7.87 (1H,

dd, J=2.0, 7.6Hz), 8.06 (1H, dd, J=2.0, 4.8Hz), 9.05 (1H, t, J=5.2Hz).

Example A-69. 2-Amino-N-(5-(4-fluoro-phenoxy)-thiophen-2-ylmethyl)-nicotinamide

The title compound (174mg, 0.507mmol, 65%) was obtained from  
5 2-aminonicotinic acid (107mg, 0.777mmol) and  
C-(5-(4-fluoro-phenoxy)-thiophen-2-yl)methylamine described in Preparation  
Example 28 (191mg, 0.856mmol) according to an analogous method to Example  
H-1.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.45 (2H, d, J=6.0Hz), 6.46 (1H, d, J=4.0Hz),  
10 6.55 (1H, dd, J=4.8, 7.6Hz), 6.74 (1H, d, J=4.0Hz), 6.85-7.25 (6H, m), 7.86 (1H, d,  
J=7.6Hz), 8.05 (1H, d, J=4.8Hz), 8.96-9.08 (1H, m).

Example A-70. 2-Amino-N-(5-benzyl-thiophen-2-ylmethyl)-nicotinamide

The title compound (67mg, 0.21mmol, 92%) was obtained from  
15 2-aminonicotinic acid (31mg, 0.224mmol) and  
C-(5-benzyl-thiophen-2-yl)-methylamine described in Preparation Example 42 (50mg,  
0.245mmol) according to an analogous method to Example H-1.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.04 (2H, s), 4.46 (2H, d, J=5.2Hz), 6.54 (1H,  
dd, J=4.8, 8.0Hz), 6.69 (1H, d, J=3.6Hz), 6.78 (1H, d, J=3.6Hz), 7.05 (2H, brs),  
20 7.15-7.30 (5H, m), 7.85 (1H, dd, J=2.0, 8.0Hz), 8.04 (1H, d, J=2.0, 4.8Hz), 8.98 (1H, t,  
J=5.2Hz).

Example A-71. 2-Amino-N-(5-(3-fluoro-benzyl)-thiophen-2-ylmethyl)-nicotinamide

The title compound (13mg, 0.038mmol, 19%) was obtained from  
25 2-aminonicotinic acid (28mg, 0.205mmol) and

C-(5-(3-fluoro-benzyl)-thiophen-2-yl)-methylamine obtained by the method described in Example A-146 (50mg, 0.226mmol) according to an analogous method to Example H-1.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.07 (2H, s), 4.40-4.53 (2H, m), 6.50-6.58 (1H, m), 6.71 (1H, d, J=3.2Hz), 6.79 (1H, d, J=3.2Hz), 6.94-7.09 (5H, m), 7.22-7.37 (1H, m), 7.84 (1H, dd, J=1.6, 8.0Hz), 8.03-8.06 (1H, m), 8.92-9.03 (1H, m).

Example A-72. 2-Amino-N-(4-phenoxy-thiophen-2-ylmethyl)-nicotinamide

The title compound (108mg, 0.331mmol, 74%) was obtained as a white solid from C-(4-phenoxy-thiophen-2-yl)-methylamine described in Preparation Example 111 (92mg, 0.45mmol) and 2-amino-nicotinic acid (68mg, 0.49mmol) according to an analogous method to Example A-26.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.70 (2H, d, J=5.7Hz), 6.35 (3H, brs), 6.48 (1H, d, J=1.7Hz), 6.60 (1H, dd, J=4.9, 7.7Hz), 6.82 (1H, d, J=1.6Hz), 7.05 (2H, dd, J=1.1, 8.6Hz), 7.11 (1H, tt, J=1.1, 7.7Hz), 7.34 (2H, t, J=8.6Hz), 7.61 (1H, dd, J=1.7, 7.7Hz), 8.18 (1H, dd, J=1.8, 4.8Hz).

Example A-73. 2-Amino-N-(5-(3-chloro-phenoxy)-thiophen-2-ylmethyl)-nicotinamide

C-(5-(3-Chloro-phenoxy)-thiophen-2-yl)-methylamine (1.02g, 4.25mmol, 98%) was obtained as an oil from 5-(3-chloro-phenoxy)-thiophene-2-carbonitrile described in Preparation Example 121 (1.02g, 4.32mmol) according to an analogous method to Example E-24. Then, title compound (12.1mg) was obtained from the resulting C-(5-(3-chloro-phenoxy)-thiophen-2-yl)-methylamine (30mg, 0.13mmol) and 2-amino-nicotinic acid (17mg, 0.13mmol). Trifluoroacetic acid salt of the title compound (12.1mg) was obtained by reverse phase high performance liquid

chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used).

MS m/e (ESI): 360.3(MH<sup>+</sup>)

5 Example A-74. 2-Amino-N-(5-(2-fluoro-benzyl)-thiophen-2-ylmethyl)nicotinamide

The title compound was obtained from C-(5-(2-fluoro-benzyl)thiophen-2-yl)-methylamine described in Preparation Example 126 (30mg, 0.14mmol) and 2-amino-nicotinic acid (21mg, 0.15mmol) according to an analogous method to Example A-26. Trifluoroacetic acid salt of the title compound  
10 (16.2mg) was obtained by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used).

MS m/e (ESI) 342.34(MH<sup>+</sup>)

Example A-75. 2-Amino-N-(5-(4-chloro-phenoxy)-thiophen-2-ylmethyl)-nicotinamide

15 A solution of 2-amino-nicotinic acid (21mg, 0.15mmol), C-(5-(4-chloro-phenoxy)-thiophen-2-yl)-methylamine described in Preparation Example 157 (36mg, 0.15mmol), benzotriazol-1-yl-tris(dimethylamino)phosphonium hexafluorophosphate (86mg, 0.195mmol) and triethylamine (0.065mL, 0.45mmol) in dimethylsulfoxide (1mL) was stirred under nitrogen atmosphere for 17 hours at room  
20 temperature. This reaction solution was purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), and trifluoroacetic acid salt of the title compound (17.7mg, 0.037mmol, 24.9%) was obtained as a pale yellow solid.

MS m/e (ESI) 360(MH<sup>+</sup>)

25

Example A-76. 2-Amino(5-(2-chloro-phenoxy)-thiophen-2-ylmethyl)-nicotinamide

Trifluoroacetic acid salt of the title compound (31.1mg, 0.07mmol, 43.7%) was obtained as a light brown oil from 2-amino-nicotinic acid (21mg, 0.15mmol) and C-(5-(2-chloro-phenoxy)-thiophen-2-yl)-methylamine described in Preparation Example 159 (36mg, 0.15mmol) according to an analogous method to Example A-75.  
MS m/e (ESI) 360(MH<sup>+</sup>)

Example A-77. 2-Amino-N-(5-(2,2-dicyclopropylvinyl)thiophen-2-ylmethyl)nicotinamide

The title compound (25mg, 0.0742mol, 53.8%) was obtained as a white solid from 2-aminonicotinic acid (19mg, 0.138mmol) and C-(5-(2,2-dicyclopropylvinyl)thiophen-2-yl)methylamine described in Preparation Example 152 (30mg, 0.138mmol) according to an analogous method to Example A-149.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.42-0.46(2H, m), 0.61-0.66(2H, m), 0.83-0.88(4H, m), 1.16-1.23(1H, m), 1.96-2.03(1H, m), 4.73(2H, d, J=5.6Hz), 6.27(1H, s), 6.34(3H, s), 6.58(1H, dd, J=4.8, 7.6Hz), 6.84(1H, d, J=3.2Hz), 6.92(1H, d, J=3.2Hz), 7.57(1H, dd, J=1.6, 7.6Hz), 8.16(1H, dd, J=1.6, 4.8Hz).

Example A-78. 2-Amino-5-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

Trifluoroacetic acid salt of the title compound (1.7mg, 0.0036mmol, 5.6%) was obtained as a by-product from Example A-171.

MS m/e (ESI) 360.1 (MH<sup>+</sup>)

Example A-79. 2-Amino-5-methyl-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

The title compound was obtained from 2-amino-5-iodine-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Preparation Example A+-16 (10mg, 22 $\mu$ mol) according to an analogous method to Example A-170. Trifluoroacetic acid salt of the title compound was obtained by  
5 reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used).

MS m/e (ESI) 340.12(MH<sup>+</sup>)

Example A-80.

10 2-Amino-6-(1-pentynyl)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

Trifluoroacetic acid salt of the title compound (0.70mg, 0.00014mmol, 3.3%) was obtained from 2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-101 (15mg, 0.042mmol) and 1-pentyne (3.4mg, 0.050mmol) according to an analogous method to Example A-91.

15 MS m/e (ESI) 392.2 (MH<sup>+</sup>)

Example A-81. 2-Amino-N-(5-phenoxy-thiophen-2-ylmethyl)-6-(3-[1,2,3]-triazol-2-yl-propylamino)-nicotinamide

The title compound (14.96mg, 0.027mmol, 9.2%) was obtained from  
20 2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-101 (105mg, 0.292mmol) and 3-[1,2,3]triazol-2-yl-propylamine (279mg, 2.21mmol) according to an analogous method to Example A-126.

MS m/e(ESI) 450.38(MH<sup>+</sup>)

25 Example A-82.

2-Amino-6-(furfurylamino)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

2-Amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-101 (180mg, 0.5mmol) was dissolved in a mixture solution of dimethylsulfoxide (2mL) and diisopropylethylamine (1mL), furfurylamine (0.663mL, 5 7.5mmol) was added, and the solution was heated in a sealed tube for 13 hours 30 minutes (oil bath temperature: 135°C). The reaction mixture was allowed to room temperature, poured into brine, and the solution was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, then concentrated, the obtained residue was purified by NH silica gel column chromatography (ethyl 10 acetate : hexane = 2 : 1), and the title compound (144mg, 0.34mmol, 68%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.50 (2H, d, J=5.6 Hz), 4.61 (2H, dd, J=0.8, 5.6 Hz), 4.87-4.94 (1H, m), 5.74 (1H, d, J=8.8 Hz), 6.04 (1H, t, J=5.2 Hz), 6.22 (1H, dd, J=0.8, 3.2 Hz), 6.31 (1H, dd, J=2.0, 3.2Hz), 6.38 (1H, d, J=3.6 Hz), 6.45 (2H, brs), 15 6.69-6.72 (1H, m), 7.06-7.12 (3H, m), 7.29-7.38 (4H, m).

Example A-83. 2-Amino-N-(5-phenoxy-thiophen-2-ylmethyl)-6-(2-pyridin-2-yl-ethylamino)-nicotinamide

2-Amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described 20 in Example A-101 (20mg, 37μmol) and 2-pyridin-2-yl-ethyl amine (66μl, 0.56mmol) were dissolved in a mixture solvent of dimethylsulfoxide (1mL) and N,N-diisopropylethylamine (0.5mL), and the solution was stirred at 130°C for 17 hours. The reaction solution was cooled to room temperature, water and ethyl acetate were added, the organic layer was partitioned, washed with water and brine, 25 and dried over anhydrous magnesium sulfate. The solvent was evaporated *in*



*vacuo*, the residue was purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), and the title compound (17.7mg) was obtained as a trifluoroacetic acid salt.

5 MS m/e (ESI) 446.05(MH<sup>+</sup>)

Example A-84. 2-Amino-N-(5-phenoxy-thiophen-2-ylmethyl)-6-(tetrahydrofuran-2-ylmethoxy)-nicotinamide

Trifluoroacetic acid salt of the title compound (16mg, 0.030mmol, 31%) was  
 10 obtained from 2-amino-6-chloronicotinic acid described in Preparation Example A-4 (17mg, 0.096mmol), tetrahydrofuran-2-ylmethanol (0.5mL) and C-(5-phenoxy-thiophen-2-yl)-methylamine (20mg, 0.097mmol) according to an analogous method to Example A-29.

MS m/e (ESI) 426.2 (MH<sup>+</sup>)

15

Example A-85. 2-Amino-6-N-(5-phenoxy-thiophen-2-ylmethyl)-6-(2-(R)-(-)-tetrahydrofurfurylamino)-nicotinamide

2-Amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-101 (30mg, 0.083mmol) was dissolved in a mixture solution of  
 20 dimethylsulfoxide (1mL) and diisopropylethylamine (0.5mL), (R)-(-)-tetrahydrofurfuryl amine (0.086mL, 0.83mmol) was added thereto, followed by heating in a sealed tube for 22 hours 30 minutes (oil bath temperature: 130°C). The reaction mixture was allowed to room temperature, poured into brine, and the solution was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, then  
 25 concentrated, the obtained residue was purified by silica gel column chromatography

(ethyl acetate : hexane = 1 : 1), and the title compound (22mg, 0.052mmol, 62%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 1.56-1.67 (1H, m), 1.86-2.04 (3H, m), 3.23-3.31 (1H, m), 3.54-3.62 (1H, m), 3.73-3.80 (1H, m), 3.84-3.91 (1H, m), 4.02-4.09 (1H, m),  
 5 4.61 (2H, d, J= 5.6 Hz), 4.91-5.02 (1H, m), 5.71 (1H, d, J=8.8 Hz), 5.98-6.04 (1H, m),  
 6.38 (1H, d, J=3.6 Hz), 6.46 (2H, brs), 6.71 (1H, d, J=3.6 Hz), 7.06-7.12 (3H, m),  
 7.29-7.37 (3H, m).

Example A-86. 2-Amino-N-(5-phenoxy-thiophen-2-ylmethyl)-6-thiazol-2-yl-

10 nicotinamide

To a solution of 2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-101 (101mg, 0.281mmol) in xylene (7mL) were added 2-tributylstanyl thiazole (137mg, 0.365mmol) and tetrakis(triphenylphosphine)palladium(0) (81mg,  
 15 0.070mmol) under nitrogen atmosphere, and the solution was stirred for 12 hours at 120°C. The reaction mixture was concentrated, the obtained residue was purified by silica gel chromatography (toluene-ethyl acetate), then, the obtained residue was washed by hexane-ethyl acetate (20:1), and the title compound (22mg, 0.054mmol, 19%) was obtained.

20 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.50 (2H, d, J=6.0Hz), 6.50 (1H, d, J=3.6Hz), 6.78 (1H, d, J=3.6Hz), 7.00-7.18 (3H, m), 7.22-7.50 (5H, m), 7.85 (1H, d, J=2.8Hz), 7.97 (1H, d, J=2.8Hz), 8.05 (1H, d, J=8.4Hz), 9.12-9.22 (1H, m).

Example A-87. 2-Amino-6-(3-methyl-2-butenyl)-N-(5-phenoxy-thiophen-2-ylmethyl)-

25 nicotinamide

Trifluoroacetic acid salt of the title compound (0.71mg, 0.0014mmol, 1.7%) was obtained from 2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-101 (30mg, 0.083mmol) and tributyl(3-methyl-2-butenyl)tin (0.084mL, 0.25mmol) according to an analogous method to Example A-29, followed by purifying by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used).  
MS m/e (ESI) 394.2 (MH<sup>+</sup>)

10 Example A-88. 2-Amino-6-(3-dimethylamino-1-propynyl)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

Trifluoroacetic acid salt of the title compound (1.00mg, 0.00019mmol, 4.6%) was obtained from 2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-101 (15mg, 0.042mmol) and 1-dimethylamino-2-propyn (4.2mg, 0.050mmol) according to an analogous method to Example A-91.  
15 MS m/e (ESI) 407.2 (MH<sup>+</sup>)

Example A-89. 2-Amino-6-(3-fluoro-benzylamino)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

The title compound (20.6mg, 0.0365mmol, 43%) was obtained from 2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-101 (30mg, 0.085mmol) and 3-fluorobenzylamine (146μl, 1.28mmol) according to an analogous method to Example A-94.  
MS m/e (ESI) 449.50(MH<sup>+</sup>)

25 Example A-90. 2-Amino-6-(3-methoxy-1-(Z)-propenyl)-N-(5-phenoxy-thiophen-2-

ylmethyl)-nicotinamide

To a mixture of 2-amino-6-(3-methoxy-1-propynyl)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-91 (11mg, 0.028mmol) and tetrahydrofuran (1mL) were added quinoline (5.4mg, 0.042mmol) and Lindlar catalyst (5.0mg), and the mixture was stirred under hydrogen atmosphere at room temperature for 15 minutes. The interior of the reaction system was exchanged with nitrogen, then, filtration was carried out through Celite pad, and the solvent was evaporated *in vacuo*. The residue was filtered by NH silica gel, then, purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), and trifluoroacetic acid salt of the title compound (4.4mg, 0.0086mmol, 31%) was obtained.

MS m/e (ESI) 396.5 (MH<sup>+</sup>)

15 Example A-91. 2-Amino-6-(3-methoxy-1-propynyl)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

A mixture of 2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-101 (15mg, 0.042mmol), methylpropargyl ether (3.5mg, 0.050mmol), diisopropylethylamine (0.023mL, 0.13mmol), pyridine (0.011mL, 0.13mmol), catalytic amount of copper(I) iodide, tetrakis(triphenylphosphine)palladium(0) (9.6mg, 0.0083mmol) and N-methylpyrrolidinone (1mL) was stirred for 4 hours at 120°C. After cooling, water and dichloromethane were added to the reaction solution for extraction, and the organic layer was filtered through a membrane filter. The solvent was evaporated *in vacuo*, then, the residue was purified by reverse phase high

performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), trifluoroacetic acid salt of the title compound (1.5mg, 0.00030mmol, 7.2%) was obtained.

MS m/e (ESI) 394.2 (MH<sup>+</sup>)

5

Example A-92.

2-Amino-6-(2-(4-amino-phenylamine)-ethylamino)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

2-Amino-6-(2-(4-nitro-phenylamino)-ethylamino)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Preparation Example A+-13 (17mg, 28μmol), iron powder (7.7mg, 138μmol) and ammonium chloride (4.41mg, 83μmol) were suspended in a mixture solvent of ethanol (1mL) and water (250μl), and the solution was stirred at 90°C for 8 hours. The reaction suspension was cooled to room temperature, then, filtered through Celite pad, water was added to the filtrate, which was then extracted with ethyl acetate, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by NH silica gel column chromatography (ethyl acetate : methanol = 10 : 1), and the title compound (10mg, 21μmol, 77%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 2.80 (3H, brs), 3.28 (2H, t, J=5.7Hz), 3.53 (2H, q, J=5.7Hz), 4.60 (2H, d, J=4.9Hz), 4.91 (1H, t, J=5.7Hz), 5.68 (1H, d, J=8.6Hz), 6.07 (1H, t, J=5.3Hz), 6.38 (1H, d, J=3.7Hz), 6.46 (2H, s), 6.52 (2H, d, J=8.8Hz), 6.60 (2H, d, J=8.8Hz), 6.71 (1H, d, J=3.7Hz), 7.08-7.12 (3H, m), 7.29-7.34 (3H, m).

Example A-93.

25 2-Amino-N-(5-phenoxy-thiophen-2-ylmethyl)-6-(2-(4-sulfamoyl-phenylamino)-

ethylamino)-nicotinamideMS m/e (ESI) 539.47 (MH<sup>+</sup>)Example A-94.5 2-Amino-6-(4-chloro-benzylamino)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

To a solution of 2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-101 (35mg, 0.10mmol) in dimethylsulfoxide (1mL) were added 4-chlorobenzylamine (234μl, 1.92mmol) and N,N-diisopropylethylamine (1.0mL, 5.74mmol), and the solution was stirred at 140°C for 2.5 days. Ethanolamine (116μl, 1.92mmol) and N,N-diisopropylethylamine (1.0mL, 5.74mmol) were added to the reaction mixture, which was then further stirred at 140°C for 2.5 days. Water was added to the reaction mixture, which was then extracted with ethyl acetate, the organic layer was sequentially washed with water and brine, dried over anhydrous sodium sulfate, and then, concentrated *in vacuo*. The obtained residue was purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), and the title compound (13.8mg, 0.024mmol, 24%) was obtained as a trifluoroacetic acid salt.

MS m/e (ESI) 465.07(MH<sup>+</sup>)

20

Example A-95.2-Amino-6-(4-fluoro-benzylamino)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

The title compound (10.6mg, 0.0188mmol, 16%) was obtained as a trifluoroacetic acid salt from 2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in

25

Example A-101 (41mg, 0.12mmol) and 4-fluorobenzylamine (200μl, 1.75mmol) according to an analogous method to Example A-94.

MS m/e (ESI) 449.56(MH<sup>+</sup>)

5 Example A-96. 2-Amino-6-(4-methoxy-benzylamino)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

The title compound (19.4mg, 0.034mmol, 37%) was obtained as a trifluoroacetic acid salt from 2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in

10 Example A-101 (32mg, 0.091mmol) and 4-methoxybenzylamine (238μl, 1.82mmol) according to an analogous method to Example A-126.

MS m/e (ESI) 461.21 (MH<sup>+</sup>)

Example A-97. 2-Amino-N-(5-phenoxy-thiophen-2-ylmethyl)-6-(4-trifluoromethyl-benzylamino)-nicotinamide

15

The title compound (15.0mg, 0.024mmol, 16%) was obtained from 2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-101 (54mg, 0.15mmol) and 4-(trifluoromethyl)benzylamine (330μl, 2.45mmol) according to an analogous method to Example A-126.

20 MS m/e(ESI) 499.10(MH<sup>+</sup>)

Example A-98. 6-Acetyl-2-amino-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

To a mixture of 2-amino-6-(1-ethoxy vinyl)-N-(5-phenoxy-thiophene-2-ylmethyl)-nicotinamide described in Preparation

25 Example A+-14 (2.0mg, 0.0051mmol) and acetone (2mL) were added water (1mL)

and concentrated sulfuric acid (0.2mL), and the solution was stirred at room temperature for 3 hours. The reaction solution was neutralized with an aqueous solution of saturated sodium bicarbonate, and ethyl acetate was added for extraction. The organic layer was washed with brine, the solvent was then evaporated *in vacuo*,  
 5 the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 2 : 1), and the title compound (1.0mg, 0.0027mmol, 53%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 2.62 (3H, s), 4.66 (2H, d, J=5.7Hz), 6.39-6.40 (4H, m), 6.75 (1H, d, J=3.9Hz), 7.08-7.13 (3H, m), 7.29-7.35 (3H, m), 7.71 (1H, d,  
 10 J=7.9Hz).

Example A-99.

(6-Amino-5-((5-phenoxy-thiophen-2-ylmethyl)-carbamoyl)-pyridin-2-ylamino)-acetic acid

15 Glycine (610mg, 8.13mmol) and 1,8-diazabicyclo[5,4,0]undec-7-ene (405μl, 2.71mmol) were added to 2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-101 (195mg, 0.542mmol) under nitrogen atmosphere, and the solution was stirred at 190°C for 4 hours. Dimethylsulfoxide (5mL) was added to the  
 20 reaction mixture, which was then filtered with a polytetrafluoroethylene membrane filter (Whatman Inc), the filtrate was purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), title compound (55.36mg, 0.108mmol, 20%) was obtained.

25 MS m/e(ESI) 399.30(MH<sup>+</sup>)



Example A-100. 2-amino-6-(1-(Z)-hydroxyimino-ethyl)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

To a mixture of 5 6-acetyl-2-amino-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-98 (9.0mg, 0.024mmol), ethanol (1mL) and water (0.5mL) were added hydroxylamine hydrochloride (2.6mg, 0.037mmol) and sodium acetate (3.0mg, 0.037mmol), and the solution was stirred under reflux for 6 hours. After cooling the reaction solution, water and ethyl acetate were added for extraction. The solvent 10 was evaporated *in vacuo*, the residue was purified by NH silica gel column chromatography (methanol : ethyl acetate = 1 : 50), and the title compound (8.3mg, 0.022mmol, 90%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 2.28 (3H, s), 4.65 (2H, d, J=5.7Hz), 6.36 (1H, brs), 6.39 (1H, d, J=3.9Hz), 6.49 (2H, brs), 6.74 (1H, d, J=3.8Hz) , 7.06-7.13 (4H, m), 15 7.31-7.34 (2H, m), 7.58 (1H, d, J=8.1Hz).

Example A-101. 2-Amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

2-Amino-6-chloro-nicotinic acid described in Preparation Example A-1 (400mg, 2.31mmol) was dissolved in N,N-dimethylformamide (10mL), triethylamine 20 (0.78mL, 5.6mmol), benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (1.23g, 2.8mmol) and C-(5-phenoxy-thiophen-2-yl)-methylamine described in Preparation Example 26 (572mg, 2.8mmol) were added, and the solution was stirred for 13 hours 30 minutes at room temperature. After the reaction was completed, the reaction solution was 25 poured into brine, which was then extracted with ethyl acetate. The organic layer

was dried over anhydrous magnesium sulfate, then concentrated, the obtained residue was purified by silica gel column chromatography (ethyl acetate : hexane = 1 : 1), and the title compound (380mg, 1.05mmol, 46%) was obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.47 (2H, d, J=6.0Hz), 6.50 (1H, d, J=4.0Hz),  
 5 6.64 (1H, d, J=8.0Hz), 6.78 (1H, d, J=4.0Hz), 7.07-7.17 (3H, m), 7.36-7.41 (2H, m),  
 7.53 (2H, brs), 7.93 (1H, d, J=8.0Hz), 9.11 (1H, t, J=6.0Hz).

Example A-102. 2-Amino-6-cyclopropyl-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

10 To a mixture of  
 2-amino-N-(5-phenoxy-thiophen-2-ylmethyl)-6-vinyl-nicotinamide described in  
 Example A-168 (6.0mg, 0.017mmol) and toluene (0.5mL) were added diiodine  
 methane (0.0055mL, 0.068mmol) and diethyl zinc (1.1M toluene solution, 0.046mL,  
 0.051mmol) on an ice bath, and the solution was stirred at room temperature for 30  
 15 minutes. Water, ethyl acetate and an aqueous solution of 29% ammonia were  
 added to the reaction solution for extraction, the organic layer was then washed with  
 brine. The organic layer was evaporated *in vacuo*, then, residue was purified by  
 reverse phase high performance liquid chromatography (acetonitrile-water mobile  
 phase (containing 0.1% trifluoroacetic acid) was used), and trifluoroacetic acid salt of  
 20 the title compound (0.40mg, 0.00083mmol, 4.9%) was obtained.

MS m/e (ESI) 366.1 (MH<sup>+</sup>)

Example A-103. 2-Amino-6-cyclopropylamino-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

25 2-Amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described

in Example A-101 (30mg, 0.083mmol) was dissolved in a mixture solution of dimethylsulfoxide (1mL) and N,N-diisopropylethylamine (0.5mL), cyclopropylamine (0.058mL, 0.84mmol) was added thereto, followed by heating in a sealed tube for 15 hours 30 minutes (oil bath temperature: 130°C). The reaction mixture was allowed to room temperature, poured into brine, which was then extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, then concentrated, the obtained residue was purified by NH silica gel column chromatography (ethyl acetate : hexane = 2 : 1), and the title compound (15mg, 0.039mmol, 47.5%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.52-0.57 (2H, m), 0.74-0.80 (2H, m), 2.47-2.54 (1H, m), 4.62 (2H, d, J=5.6 Hz), 5.09 (1H, brs), 6.06 (1H, d, J=8.4 Hz), 6.08-6.14 (1H, m), 6.34-6.42 (3H, m), 6.72 (1H, d, J=3.6 Hz), 7.06-7.13 (3H, m), 7.29-7.35 (2H, m), 7.45 (1H, d, J=8.4 Hz).

Example A-104. 2-Amino-6-(cyclopropylmethyl-amino)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

MS m/e (ESI) 395.22 (MH<sup>+</sup>)

Example A-105. 2-Amino-6-(2-ethoxy-ethylamino)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

2-Amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-101 (100mg, 0.28mmol) was dissolved in a mixture solution of dimethylsulfoxide (2mL) and N,N-diisopropylethylamine (1mL), 2-ethoxyethylamine (0.051mL, 0.49mmol) was added thereto, followed by heating in a sealed tube for 32 hours 40 minutes (oil bath temperature: 130°C). The reaction mixture was allowed

to room temperature, poured into brine, and the solution was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, then concentrated, the obtained residue was purified by NH silica gel column chromatography (ethyl acetate : hexane = 2 : 1), and the title compound (37mg, 0.09mmol, 32%) was obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 1.11 (3H, t, J=7.2 Hz), 3.35-3.47 (6H, m), 4.41 (2H, d, J=6.0 Hz), 5.72 (1H, d, J=8.8 Hz), 6.48 (1H, d, J=3.6 Hz), 6.67-6.77 (2H, m), 7.05 (2H, brs), 7.05-7.16 (3H, m), 7.35-7.41 (2H, m), 7.59 (1H, d, J=8.8 Hz), 8.39 (1H, t, J=6.0Hz).

#### Example A-106.

##### 2-Amino-6-ethylamino-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

2-Amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-101 (200mg, 0.56mmol) was dissolved in a mixture solution of dimethylsulfoxide (1mL) and N,N-diisopropylethylamine (0.5mL), ethylamine (2M tetrahydrofuran solution) (2mL, 4mmol) was added thereto, followed by heating in a sealed tube for 17 hours (oil bath temperature: 135°C). The reaction mixture was allowed to room temperature, poured into brine, and the solution was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, then concentrated, the obtained residue was purified by NH silica gel column chromatography (ethyl acetate : hexane = 2 : 1), and the title compound (117mg, 0.32mmol, 57%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 1.21 (3H, t, J=7.2 Hz), 3.24-3.32 (2H, m), 4.56-4.63 (3H, m), 5.67 (1H, d, J=8.8 Hz), 6.06-6.11 (1H, m), 6.37 (1H, d, J=4.0 Hz), 6.42 (2H, brs), 6.71 (1H, d, J=4.0 Hz), 7.06-7.12 (3H, m), 7.29-7.34 (2H, m), 7.37 (1H,

J=8.8 Hz).

Example A-107. (±)-2-(6-Amino-5-((5-phenoxy-thiophen-2-ylmethyl)-carbamoyl)-pyridin-2-ylamino)-(R)-3-hydroxy-butyric acid

5 Trifluoroacetic acid salt of the title compound (12mg, 0.022mmol, 26%) was obtained from 2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-101 (30mg, 0.083mmol) and (L)-threonine (99mg, 0.83mmol) according to an analogous method to Example A-99.

MS m/e (ESI) 443.1 (MH<sup>+</sup>)

10

Example A-108. (±)-2-(6-Amino-5-((5-phenoxy-thiophen-2-ylmethyl)-carbamoyl)-pyridin-2-ylamino)-3-phenyl-propionic acid

Trifluoroacetic acid salt of the title compound (11mg, 0.019mmol, 23%) was obtained from 2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide  
15 described in Example A-101 (30mg, 0.083mmol) and (±)-phenylalanine (140mg, 0.83mmol) according to an analogous method to Example A-99, trifluoroacetic acid salt of the title compound (11mg, 0.019mmol, 23%) was obtained.

MS m/e (ESI) 489.1 (MH<sup>+</sup>)

20 Example A-109. (±)-2-(6-Amino-5-((5-phenoxy-thiophen-2-ylmethyl)-carbamoyl)-pyridin-2-ylamino)-4-methyl-pentanoic acid

Trifluoroacetic acid salt of the title compound (6.8mg, 0.012mmol, 14%) was obtained from 2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-101 (30mg, 0.083mmol) and (D)-leucine (110mg, 0.83mmol)  
25 according to an analogous method to Example A-99.

MS m/e (ESI) 455.2 (MH<sup>+</sup>)

Example A-110. (±)-2-(6-Amino-5-((5-phenoxy-thiophen-2-ylmethyl)-carbamoyl)-pyridin-2-ylamino)-3-methoxy-propionic acid

5 Trifluoroacetic acid salt of the title compound (5.5mg, 0.0099mmol, 12%) was obtained from 2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-101 (30mg, 0.083mmol) and (±)-O-methylserine (99mg, 0.83mmol) according to an analogous method to Example A-99.

MS m/e (ESI) 443.1 (MH<sup>+</sup>)

10

Example A-111. (±)-2-(6-Amino-5-((5-phenoxy-thiophene-2-ylmethyl)-carbamoyl)-pyridin-2-ylamino)-pentanedioic acid

Trifluoroacetic acid salt of the title compound (1.7mg, 0.0029mmol, 3.5%) was obtained from 2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide  
15 described in Example A-101 (30mg, 0.083mmol) and (±)-glutamic acid (122mg, 0.83mmol) according to an analogous method to Example A-99.

MS m/e (ESI) 471.4 (MH<sup>+</sup>)

Example A-112. 2-Amino-6-(2-hydroxyethoxy)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

20

Sodium hydride (3.1mg, 0.078mmol, 60% in oil), catalytic amount of copper(I) iodide, 2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-102 (4.0mg, 0.011mmol) were added sequentially to ethyleneglycol (0.7mL), and the solution was stirred at 65°C for 2 hours. After further stirring at  
25 90°C, the solution was allowed to cool to room temperature. Water,

dichloromethane and an aqueous solution of saturated ammonium chloride were added to the reaction solution for extraction, which was then washed with brine, then, the solvent was evaporated *in vacuo*. The residue was purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing  
 5 0.1% trifluoroacetic acid) was used), and trifluoroacetic acid salt of the title compound (0.26mg, 0.00052mmol, 4.7%) was obtained.

MS m/e (ESI) 386.2 (MH<sup>+</sup>)

Example A-113.

10 2-Amino-6-(2-hydroxy-ethylamino)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

2-Amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-101 (170mg, 0.47mmol) was dissolved in a mixture solution of dimethylsulfoxide (2mL) and diisopropylethylamine (1mL), ethanolamine (0.428mL, 7.1mmol) was added thereto, followed by heating in a sealed tube for 15 hours 20 minutes (oil bath temperature: 135°C). The reaction mixture was allowed to room temperature, poured into brine, and the solution was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, then concentrated, the obtained residue was purified by NH silica gel column chromatography (ethyl acetate : hexane = 2 : 1), and the title compound (138mg, 0.36mmol, 76%) was  
 20 obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.24-3.31 (2H, m), 3.44-3.51 (2H, m), 4.39 (2H, d, J=5.6 Hz), 4.68 (1H, t, J=5.2 Hz), 5.70 (1H, d, J= 8.4 Hz), 6.46 (1H, d, J=4.0 Hz), 6.66 (1H, brs), 6.70 (1H, d, J=4.0Hz), 7.02 (2H, brs), 7.04-7.14 (3H, m), 7.34-7.39 (2H, m), 7.57 (1H, d, J=8.4 Hz), 8.37 (1H, t, J=5.6 Hz).

25

Example A-114. 2-Amino-6-hydroxymethyl-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

To a mixture of 2-amino-N-(5-phenoxy-thiophen-2-ylmethyl)-6-tributylstanyl-nicotinamide described in Preparation Example A-15 (98mg, 0.16mmol) in tetrahydrofuran (1.5mL) was added *n*-butyl lithium (2.4M hexane solution, 0.32mL, 0.80mmol) dropwise at -78°C, and the solution was stirred for 1 hour 40 minutes at the same temperature. N,N-dimethylformamide (0.037mL, 0.48mmol) was added at the same temperature, the solution was stirred for 35 minutes, then, a solution of sodium borocyanide (50mg, 0.80mmol) in tetrahydrofuran (1mL) was added dropwise at the same temperature, and the solution was stirred at -3°C for 1 hour. The reaction solution was cooled to -78°C, acetic acid (0.091mL, 1.6mmol) was added, and the solution was warmed gradually to 0°C. Water, ethyl acetate and tetrahydrofuran were added to the reaction solution for extraction, which was then washed with brine, then, the solvent was evaporated *in vacuo*. The residue was purified by silica gel column chromatography (methanol : ethyl acetate = 1 : 50), and the title compound (19mg, 0.053mmol, 33%) was obtained as a pale yellow oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.58 (2H, s), 4.64 (2H, d, J=5.5Hz), 6.38 (1H, d, J=3.7Hz), 6.43 (1H, brs), 6.48-6.50 (3H, m), 6.74 (1H, d, J=3.7Hz), 7.08-7.13 (3H, m), 7.30-7.34 (2H, m), 7.59 (1H, d, J=7.9Hz).

Example A-115. 2-Amino-6-isopropoxy-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

Trifluoroacetic acid salt of the title compound (4.3mg, 0.0086mmol, 9.0%) was obtained from 2-amino-6-chloronicotinic acid described in Preparation Example



A-4 (17mg, 0.096mmol), isopropanol (0.5mL) and C-(5-phenoxy-thiophen-2-yl)-methylamine (20mg, 0.097mmol) according to an analogous method to Example A-29.

MS m/e (ESI) 384.2 (MH<sup>+</sup>)

5

Example A-116. 2-Amino-6-methoxy-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

2,5-Diamino-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Preparation Example A-10 (27mg, 59μmol), sodium nitrite (4.1mg, 59μmol) and sulfuric acid (several drops) were dissolved in methanol (5mL), and the solution was stirred for 30 minutes under reflux. An aqueous solution of saturated sodium bicarbonate was added to the reaction solution at 0°C, which was then extracted with ethyl acetate, and the organic layer was washed with brine. The solvent was evaporated *in vacuo*, the residue was purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), then, further by NH silica gel column chromatography (hexane : ethyl acetate = 5 : 1), and the title compound (0.7mg) was obtained as a white solid.

10

15

MS m/e (ESI) 356.32(MH<sup>+</sup>).

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.87 (3H, s), 4.63 (2H, d, J=5.7Hz), 6.01 (1H, d, J=8.6Hz), 6.11 (1H, brs), 6.39 (1H, d, J=3.8Hz), 6.51 (1H, brs), 6.73 (1H, d, J=3.3Hz), 7.08-7.12 (3H, m), 7.22-7.26 (1H, m), 7.32 (2H, t, J=8.6Hz), 7.50 (1H, t, J=8.8Hz).

20

Example A-117. 2-Amino-N-(5-benzofuran-5-ylmethyl-thiophen-2-ylmethyl)-6-methoxymethyl-nicotinamide

MS m/e (ESI) 407.85 (MH<sup>+</sup>)

25

Example A-118. 2-Amino-N-(5-benzo[1,3]dioxol-5-ylmethyl-thiophen-2-ylmethyl)-6-methoxymethyl-nicotinamide

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.33 (3H, s), 3.96 (2H, s), 4.28 (2H, s), 4.47 (2H, d, J=6.0Hz), 5.96 (2H, d, J=1.2Hz), 6.59 (1H, d, J=8.0Hz), 6.68-6.74 (2H, m), 6.77-6.84 (3H, m), 7.12 (2H, brs), 7.90 (1H, d, J=8.0Hz), 8.97 (1H, t, J=6.0Hz).

Example A-119. 2-Amino-6-methoxymethyl-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.46(3H, s) , 4.41(2H, s) , 4.64-4.66(2H, m) , 6.32(1H, br s), 6.39(1H, d, J=3.8Hz) , 6.47(2H, br s) , 6.71(1H, d, J=7.9Hz) , 6.74(1H, d, J=3.8Hz) , 7.08-7.13(3H, m) , 7.31-7.35(2H, m) , 7.62 (1H, d, J=7.9Hz).

Example A-120. 2-Amino-N-(4-benzylamino-benzyl)-6-methoxymethyl-nicotinamide

MS m/e (ESI) 377 (MH<sup>+</sup>)

Example A-121. 2-Amino-N-(5-benzyl-thiophen-2-ylmethyl)-6-methoxymethyl-nicotinamide

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.44 (3H, s), 4.09 (2H, s), 4.38 (2H, s), 4.66 (2H, d, J=5.2 Hz), 6.25-6.34 (1H, m), 6.38 (2H, br s), 6.65 (1H, d, J=3.6 Hz), 6.67 (1H, d, J=8.0 Hz), 6.82 (1H, d, J=3.6 Hz), 7.20-7.27 (3H, m), 7.27-7.34 (2H, m), 7.57 (1H, d, J=8.0 Hz).

Example A-122. 2-Amino-N-(5-(3-chloro-phenoxy)-thiophen-2-ylmethyl)-6-methoxymethyl-nicotinamide

MS m/e (ESI) 404 (MH<sup>+</sup>)

Example A-123. 2-Amino-N-(5-(3-fluoro-phenoxy)-thiophen-2-ylmethyl)-  
6-methoxymethyl-nicotinamide

5 MS m/e (ESI) 388 (MH<sup>+</sup>)

Example A-124. 2-Amino-6-(3-methoxy-propyl)-N-(5-phenoxy-thiophen-2-ylmethyl)-  
nicotinamide

To a mixture of  
10 2-amino-6-(3-methoxy-1-(Z)-propenyl)-N-(5-phenoxy-thiophen-2-ylmethyl)-  
nicotinamide described in Example A-90 (3.0mg, 0.0059mmol) and tetrahydrofuran  
(1mL) were added triethylamine (3.6mg, 0.036mmol) and 10% palladium-carbon  
(50% water wet, 5mg), and the solution was stirred under hydrogen atmosphere at  
room temperature for 15 minutes. The interior of the reaction system was  
15 exchanged with nitrogen, then, filtration was carried out through Celite pad, the  
solvent was evaporated *in vacuo*. The residue was purified by reverse phase high  
performance liquid chromatography (acetonitrile-water mobile phase (containing  
0.1% trifluoroacetic acid) was used), and trifluoroacetic acid salt of the title compound  
(0.48mg, 0.00094mmol, 16%) was obtained.  
20 MS m/e (ESI) 398.3 (MH<sup>+</sup>)

Example A-125. 2-Amino-6-methylamino-N-(5-phenoxy-thiophen-2-  
ylmethyl)-nicotinamide

2-Amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described  
25 in Example A-101 (200mg, 0.56mmol) was dissolved in a mixture solution of

dimethylsulfoxide (1mL) and diisopropylethylamine (0.5mL), methylamine (2M tetrahydrofuran solution) (2mL, 4mmol) was added thereto, followed by heating in a sealed tube for 14 hours (oil bath temperature: 135°C). The reaction mixture was allowed to room temperature, poured into brine, and the solution was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, then concentrated, the obtained residue was purified by NH silica gel column chromatography (ethyl acetate : hexane = 2 : 1), and the title compound (144mg, 0.41mmol, 73%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 2.89 (3H, d, J=5.2 Hz), 4.61 (2H, d, J=5.6 Hz), 4.64-4.71 (1H, m), 5.69 (1H, d, J=8.4 Hz), 6.03-6.09 (1H, m), 6.38 (1H, d, J=4.0 Hz), 6.44 (2H, brs), 6.71 (1H, d, J=4.0 Hz), 7.06-7.12 (3H, m), 7.29-7.35 (2H, m), 7.39 (1H, d, J=8.4 Hz).

Example A-126. 2-Amino-6-benzylamino-N-(5-phenoxy-thiophen-2-ylmethyl)

-nicotinamide

To a solution of 2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-101 (110mg, 0.31mmol) in dimethylsulfoxide (1mL) were added benzylamine (1.8mL, 16.5mmol) and N,N-diisopropylethylamine (0.5mL, 2.94mmol), and the solution was stirred for 17 hours at 135°C. Water was added to the reaction mixture, which was then extracted with ethyl acetate, the organic layer was sequentially washed with water and brine, dried over anhydrous sodium sulfate, and then, concentrated *in vacuo*. The obtained residue was purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), and the title compound (23.4mg, 0.0429mmol,

14%) was obtained as a trifluoroacetic acid salt.

MS m/e(ESI) 431.27(MH<sup>+</sup>)

Example A-127. 2-Amino-N-(5-phenoxy-thiophen-2-ylmethyl)-6-propoxy-nicotinamide

5 Trifluoroacetic acid salt of the title compound (5.3mg, 0.011mmol, 11%) was obtained from 2-amino-6-chloronicotinic acid described in Preparation Example A-4 (17mg, 0.096mmol), propanol (0.5mL) and C-(5-phenoxy-thiophen-2-yl)-methylamine (20mg, 0.097mmol) according to an analogous method to Example A-29.

MS m/e (ESI) 384.1 (MH<sup>+</sup>)

10

Example A-128. 2-Amino-N-(5-phenoxy-thiophen-2-ylmethyl)-6-propylamino-nicotinamide

2-Amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-101 (200mg, 0.56mmol) was dissolved in a mixture solution of  
15 dimethylsulfoxide (2mL) and N,N-diisopropylethylamine (1mL), propylamine (0.685mL, 8.3mmol) was added thereto, followed by heating in a sealed tube for 13 hours (oil bath temperature: 135°C). The reaction mixture was allowed to room temperature, poured into brine, which was then extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, then concentrated, the  
20 obtained residue was purified by NH silica gel column chromatography (ethyl acetate : hexane = 2 : 1), and the title compound (89mg, 0.23mmol, 42%) was obtained.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.96 (3H, t, J=7.6Hz), 1.54-1.64 (2H, m), 3.16-3.22 (2H, m), 4.59 (2H, d, J=5.6 Hz), 4.69 (1H, t, J=5.2 Hz), 5.67 (1H, d, J=8.8  
25 Hz), 6.16 (1H, t, J=5.6 Hz), 6.36 (1H, d, J=3.6 Hz), 6.42 (2H, brs), 6.69 (1H, d, J=3.6

Hz), 7.06-7.12 (3H, m), 7.29-7.35 (2H, m), 7.37 (1H, J=8.8 Hz).

Example A-129. 2-Amino-N-(5-phenoxy-thiophen-2-ylmethyl)-6-((pyrazin-2-ylmethyl)-amino)-nicotinamide

5 MS m/e (ESI) 433.15(MH<sup>+</sup>)

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.33(2H, d, J=5.6Hz), 4.68(2H, s), 5.99(1H, d, J=8.4Hz), 6.48(1H, d, J=3.6Hz), 6.73(1H, d, J=3.6Hz), 7.06-7.15(3H, m), 7.34-7.39(2H, m), 7.82-7.94 (1H, m), 8.55 (1H, d, J=2.4Hz), 8.60 (1H, dd, J=2.4,1.2Hz), 8.67 (1H, d, J=1.2Hz), 8.69-8.79(1H, m).

10

Example A-130. 2-Amino-N-(5-phenoxy-thiophen-2-ylmethyl)-6-((pyridin-2-ylmethyl)-amino)-nicotinamide

MS m/e (ESI) 432.17 (MH<sup>+</sup>)

15 Example A-131.

3-(3-(5-Phenoxy-thiophen-2-ylmethyl)-carbamoyl)-pyridin-2-ylamino)-propionic acid

To a solution of 2-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Preparation Example A+-7 (51mg, 0.15mmol) in dimethylsulfoxide (2mL) were added *tert*-butyl 3-aminopropanoate hydrochloride (32mg, 0.178mmol) and triethylamine (27μl, 0.192mmol), and the solution was stirred for 2.5 hours at 120°C. Potassium carbonate (49mg, 0.36mmol) was added to the reaction mixture, which was then stirred for 20 hours 120°C. Water was added to the reaction mixture, which was then extracted with ethyl acetate and concentrated. To a solution of the resulting residue in dichloromethane (1mL) was added trifluoroacetic acid (500μl, 6.49mmol), and the solution was stirred for 2.5 hours at room temperature. The

20

25

reaction mixture was concentrated, and the obtained residue was purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), and the title compound (3.44mg, 0.0067mmol, 4.5%) was obtained as a trifluoroacetic acid salt.

5 MS m/e(ESI) 398.52(MH<sup>+</sup>)

Example A-132. 2-Chloro-N-(5-(3-fluorophenoxy)thiophen-2-ylmethyl)-6-methylnicotinamide

The title compound (330mg, 0.877mmol, 65.0%) was obtained as a white  
 10 solid from 2-chloro-6-methylnicotinic acid (230mg, 1.35mmol) and C-(5-(3-fluorophenoxy)thiophen-2-yl)methylamine described in Preparation Example 23 (300mg, 1.35mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.47 (3H, s), 4.52 (2H, d, J=5.6Hz), 6.60-6.63 (1H, m), 6.84 (1H, d, J=4.0Hz), 6.92-7.04 (3H, m), 7.32-7.35 (1H, m), 7.40-7.47 (1H,  
 15 m), 7.77 (1H, d, J=7.2Hz), 9.15 (1H, t, J=5.6Hz).

Example A-133. 2-(Cyclopropylmethyl-amino)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

To a solution of 2-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide  
 20 described in Preparation Example A+-7 (64mg, 0.186mmol) in dimethylsulfoxide (1mL) was added (aminomethyl)cyclopropane (48μl, 0.56mmol), and the solution was stirred for 14 hours at 120°C. The reaction mixture was purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), and the title compound (15.6mg, 0.0316mmol,  
 25 17%) was obtained as a trifluoroacetic acid salt.

MS m/e(ESI) 380.43(MH<sup>+</sup>)

Example A-134. 2-(2-Methoxy-ethylamino)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

- 5           To a solution of 2-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Preparation Example A+-7 (16mg, 0.046mmol) in 1-methyl-2-pyrrolidone (2mL) were added 2-methoxyethylamine (6μl, 0.07mmol) and sodium hydride (4mg, 0.092mmol, 60% in oil), and the solution was stirred for 8 hours at 110°C. An aqueous solution of saturated ammonium chloride was added to the reaction mixture, 10 which was then extracted with ethyl acetate, and concentrated. The obtained residue was purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), and the title compound (0.67mg, 0.0013mmol, 2.8%) was obtained as a trifluoroacetic acid salt.
- 15   MS m/e(ESI) 384.16(MH<sup>+</sup>)

Example A-135. 2-Methyl-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

- The title compound (40mg, 0.123mmol, 42.4 %) was obtained as a light brown solid from 2-methylnicotinic acid (40mg,0.29mmol) and 20 C-(5-phenoxy-thiophen-2-yl)-methylamine described in Preparation Example 26 (60mg,0.29mmol) according to an analogous technique to Example Q-6.
- <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.47 (3H, s), 4.49 (2H, d, J=5.6Hz), 6.50-6.63 (1H, m), 6.77-6.80 (1H, m), 7.06-7.16 (3H, m), 7.23-7.28 (1H, m), 7.35-7.40 (2H, m), 7.66-7.70 (1H, m), 8.48 (1H, dd, J=1.6, 4.8Hz), 9.05 (1H, t, J=5.6Hz).



Example A-136. 2-Methylamino-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

To a solution of 2-(ethoxymethyl-amino)-N-(5-phenoxy-thiophene-2-ylmethyl)-nicotinamide described in Preparation Example A-6 (148mg, 0.385mmol) in dimethylsulfoxide (3mL) was added sodium borohydride (44mg, 1.15mmol), and the solution was stirred for 30 minutes at 100°C, followed by stirring for 20 minutes at 110°C. Furthermore, sodium borohydride (35mg, 0.925mmol) was added thereto, followed by stirring for 20 minutes at 110°C. Water was added to the reaction mixture, which was then extracted with ethyl acetate, the organic layer was sequentially washed with water and brine, dried over anhydrous sodium sulfate, and then, concentrated *in vacuo*. The obtained residue was purified by silica gel chromatography (hexane-ethyl acetate), and the title compound (86mg, 0.26mmol, 67%) was obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.87 (3H, d, J=4.8Hz), 4.46 (2H, d, J=5.6Hz), 6.48 (1H, d, J=3.6Hz), 6.53 (1H, dd, J=4.8, 7.6Hz), 6.76 (1H, d, J=3.6Hz), 7.00-7.18(3H, m), 7.30-7.41 (2H, m), 7.88 (1H, dd, J=2.0, 7.6Hz), 8.10-8.25 (2H, m), 9.07 (1H, t, J=5.6Hz).

Example A-137. 6-Amino-N-(5-(3-fluorophenoxy)thiophen-2-ylmethyl)nicotinamide

The title compound (20mg, 0.058mmol, 21.6%) was obtained as a white solid from 6-amino-nicotinic acid (37mg, 0.27mmol) and C-(5-(3-fluorophenoxy)thiophen-2-yl)methylamine described in Preparation Example 23 (60mg, 0.27mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.56 (2H, d, 6.0Hz), 6.39 (1H, d, J=8.8Hz), 6.48 (2H, brs), 6.56 (1H, d, J=3.6Hz), 6.76 (1H, d, J=3.6Hz), 6.88-7.00 (3H, m), 7.39 (1H, ddd, J=8.0, 8.0, 8.0), 7.79 (1H, dd, J=2.0, 8.8Hz), 8.43 (1H, d, J=2.0Hz), 8.77

(1H, t, J=6.0Hz).

Example A-138. 6-Amino-N-(5-phenoxy thiophen-2-ylmethyl)-nicotinamide

To a solution of C-(5-phenoxythiophen-2-yl)methylamine described in  
5 Preparation Example 24 (170mg, 0.83mmol) and 6-aminonicotinic acid (130mg,  
0.91mmol) in N,N-dimethylformamide (10mL) were added  
benzotriazol-1-yl-tris(dimethylamino)phosphonium hexafluorophosphate (400mg,  
0.91mmol) and triethylamine (0.3mL, 2.2mmol), and the solution was stirred for 35  
minutes at 60°C. Water and ethyl acetate were added to the reaction solution,  
10 which was then partitioned, the organic layer was washed with water twice, NH silica  
gel was added to the organic layer, the solvent was evaporated *in vacuo* for  
adsorption, purification was carried out by NH silica gel column chromatography  
(ethyl acetate, then ethyl acetate : methanol = 50 : 1), and the title compound (130mg,  
0.40mmol, 48.2%) was obtained as a solid.  
15 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.45 (2H, d, 5.2Hz), 6.39 (1H, d, J=8.8Hz),  
6.43-6.54 (3H, m), 6.70-6.77 (1H, m), 7.20-7.16 (3H, m), 7.31-7.41 (2H, m), 7.78 (1H,  
d, J=8.8Hz), 8.43 (1H, s), 8.76 (1H, d, J=5.2Hz).

Example A-139. 6-Amino-N-(5-(4-fluoro-phenoxy)-thiophen-2-ylmethyl)-nicotinamide

20 To a solution of C-(5-(4-fluorophenoxy)thiophen-2-yl)methylamine described  
in Preparation Example 28 (500mg, 2.24mmol) and 6-aminonicotinic acid (340mg,  
2.46mmol) in N,N-dimethylformamide (10mL) were added  
benzotriazol-1-yl-tris(dimethylamino)phosphonium hexafluorophosphate (1.1g,  
2.46mmol) and triethylamine (0.6mL, 4.48mmol), and the solution was stirred for 30  
25 minutes at 60°C. Water and ethyl acetate were added to the reaction solution,

which was then partitioned, and the organic layer was washed with water three times. The organic layer was passed through a glass filter lined with NH silica gel and silica gel (1:1), and eluted with a mixture solvent of ethyl acetate and methanol (20 : 1). The solvent was evaporated *in vacuo*, ethyl acetate and hexane were added to the  
5 residue, the generated solid was collected by filtration, and the title compound (560mg, 1.63mmol, 72.8%) was obtained as a slightly yellow solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.44 (2H, d, J=5.2Hz), 6.39 (1H, d, J=8.8Hz), 6.44-6.50 (3H, m), 6.72 (1H, d, J=2.4Hz), 7.09-7.16 (2H, m), 7.16-7.24 (2H, m), 7.78 (1H, d, J=8.8Hz), 8.42 (1H, s), 8.74 (1H, t, J=5.2Hz).

10

Example A-140. 6-Amino-N-(5-(4-chloro-phenoxy)-thiophen-2-ylmethyl)-nicotinamide

The title compound (32mg, 0.089mmol, 30.7%) was obtained as a white solid from C-(5-(4-chlorophenoxy)thiophen-2-yl)methylamine obtained according to an analogous method to Example E-66 (70mg, 0.29mmol) and 6-amino-nicotinic acid  
15 (40mg, 0.29mmol).

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.45 (2H, d, 5.2Hz), 6.39 (1H, d, J=8.4Hz), 6.48 (2H, s), 6.51 (1H, d, J=3.6Hz), 6.74 (1H, d, J=3.6Hz), 7.04-7.14 (2H, m), 7.36-7.46 (2H, m), 7.78 (1H, d, J=8.4Hz), 8.43 (1H, s), 8.76 (1H, t, J=5.2Hz).

20 Example A-141. 6-Amino-N-(5-*m*-tolylloxy-thiophen-2-ylmethyl)nicotinamide

The title compound (243mg, 0.717mmol, 52.7%) was obtained as a light brown solid from C-(5-*m*-tolylloxythiophen-2-yl)methylamine described in Preparation Example 30 (300mg, 1.36mmol) and 6-aminonicotinic acid (210mg, 1.52mmol) according to an analogous method to Example Q-6.

25 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.26 (3H, s), 4.44 (2H, d, J=5.6Hz), 6.39 (1H,

d, J=8.8Hz), 6.42-6.52 (3H, m), 6.70-6.74 (1H, m), 6.84-6.95 (3H, m), 7.22 (1H, dd, J=8.0, 8.0Hz), 7.78 (1H, dd, J=2.4, 8.8Hz), 8.43 (1H, d, J=2.4Hz), 8.74 (1H, t, J=5.6Hz).

5 Example A-142. 6-Amino-N-(5-*p*-tolxyloxy-thiophene-2-ylmethyl)-nicotinamide

The title compound (265mg, 0.78mmol, 52.1%) was obtained as a light brown solid from 6-aminonicotinic acid (210mg, 1.50mmol) and C-(5-*p*-tolxyloxy-thiophen-2-yl)-methylanine described in Preparation Example 32 (300mg, 1.37mmol) according to an analogous method to Example Q-6.

10 <sup>1</sup>H-NMR Spectrum (DMSO-*d*<sub>6</sub>) δ(ppm) : 2.24 (3H, s), 4.43 (2H, d, J=5.6Hz), 6.38 (1H, d, J=8.8Hz), 6.41 (1H, d, J=3.6Hz), 6.47 (2H, s), 6.70 (1H, d, J=3.6Hz), 6.94-7.00 (2H, m), 7.13-7.19 (2H, m), 7.78 (1H, dd, J=2.4, 8.8Hz), 8.42 (1H, d, J=2.4Hz), 8.73 (1H, t, J=5.6Hz).

15 Example A-143. 6-Amino-N-(5-(4-fluoro-benzyl)-thiophen-2-ylmethyl)-nicotinamide

The title compound (46mg, 0.13mmol, 81.2%) was obtained as a white solid from C-(5-(4-fluoro-benzyl)-thiophen-2-yl)-methylanine described in Preparation Example 38 (35mg, 0.16mmol) and 6-aminonicotinic acid (24mg, 0.17mmol) according to an analogous method to Example Q-6.

20 <sup>1</sup>H-NMR Spectrum (DMSO-*d*<sub>6</sub>) δ(ppm) : 4.03 (2H, s), 4.45 (2H, d, J=5.6Hz), 6.38 (1H, d, J=8.8Hz), 6.45 (2H, s), 6.67 (1H, d, J=3.6Hz), 6.76 (1H, d, J=3.6Hz), 7.04-7.14 (2H, m), 7.20-7.30 (2H, m), 7.77 (1H, dd, J=2.4, 8.8Hz), 8.41 (1H, d, J=2.4Hz), 8.69 (1H, t, J=5.6Hz).

25 Example A-144. 6-Amino-N-(5-benzyl-thiophen-2-ylmethyl)-nicotinamide

The title compound (27mg, 0.082mmol, 31.0 %) was obtained as a white solid from 6-aminonicotinic acid (37mg, 0.27mmol) and C-(5-benzyl-thiophen-2-yl)-methylamine described in Preparation Example 42 (54mg, 0.27mmol) according to an analogous technique to Example Q-6.

5 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>)δ(ppm) : 4.03 (2H, s), 4.44 (2H, d, J=5.2Hz), 6.37 (1H, d, J=8.8Hz), 6.45 (2H, s), 6.67 (1H, d, J=3.2Hz), 6.75 (1H, d, J=3.2Hz), 7.15-7.30 (5H, m), 7.76 (1H, dd, J=2.0, 8.8Hz), 8.41 (1H, d, J=2.0Hz), 8.68 (1H, t, J=5.2Hz).

Example A-145. 6-Amino-N-(5-(3-chloro-benzyl)-thiophen-2-ylmethyl)-nicotinamide

10 The title compound (42mg, 0.12mmol, 56.0%) was obtained as a white solid from C-(5-(3-chloro-benzyl-2-yl)methylamine described in Preparation Example 45 (50mg, 0.21mmol) and 6-aminonicotinic acid (32mg, 0.23mmol) according to an analogous method to Example Q-6.

15 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.05 (2H, s), 4.45 (2H, d, J=5.6Hz), 6.38 (1H, d, J=8.8Hz), 6.45 (2H, s), 6.71 (1H, d, J=3.2Hz), 6.76 (1H, d, J=3.2Hz), 7.16-7.34 (4H, m), 7.77 (1H, d, J=8.8Hz), 8.41 (1H, s), 8.70 (1H, t, J=5.6Hz).

Example A-146. 6-Amino-N-(5-(3-fluoro-benzyl)-thiophen-2-ylmethyl)-nicotinamide

20 To a solution of 5-(3-fluoro-benzyl)-thiophene-2-carbaldehyde described in Preparation Example 53 (485mg, 2.2mmol) in 7N ammonia/methanol (30mL) was added Raney nickel (1g), and the solution was stirred under hydrogen atmosphere at room temperature for 16 hours. The reaction solution was filtered through Celite pad to remove the catalyst, then, the organic layer was concentrated *in vacuo*, the residue was purified by NH silica gel column chromatography (hexane, then hexane :  
25 ethyl acetate = 4 : 1), and C-(5-(3-fluoro-benzyl-2-yl)-methylamine (290mg, 1.3mmol,

59.6%) was obtained as a brown oil. Using the resulting C-(5-(3-fluoro-benzyl-2-yl)-methylamine (50mg, 0.226mmol) and 6-aminonicotinic acid (34mg, 0.248mmol) according to an analogous technique to Example Q-6, the title compound (44mg, 0.129mmol, 57.1 %) was obtained as a white solid.

5 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.06 (2H, s), 4.45 (2H, d, J=5.6Hz), 6.38 (1H, d, J=8.8Hz), 6.45 (2H, s), 6.70 (1H, d, J=2.0Hz), 6.77 (1H, d, J=2.0Hz), 6.96-7.10 (3H, m), 7.28-7.36 (1H, m), 7.77 (1H, dd, J=2.0, 8.8Hz), 8.41 (1H, d, J=2.0Hz), 8.70 (1H, t, J=5.6Hz).

10 Example A-147. 6-Amino-N-(5-(3-chloro-phenoxy)-thiophen-2-ylmethyl)-nicotinamide

The title compound (8.21mg) was obtained from C-(5-(3-chloro-phenoxy)-thiophen-2-yl)-methylamine described in Example A-73 (30mg, 0.13mmol) and 6-amino-nicotinic acid (17mg, 0.13mmol) according to an analogous method to Example E-24, the title compound (8.21mg) was obtained.

15 Trifluoroacetic acid salt of the tile compound was obtained by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used).

MS m/e (ESI) 360.3(MH<sup>+</sup>)

20 Example A-148. 6-Amino-N-(5-(2-fluoro-phenoxy)-thiophen-2-ylmethyl)-nicotinamide

Trifluoroacetic acid salt of the title compound (26.6mg) was obtained as a pale yellow oil from 6-amino-nicotinic acid (21mg, 0.15mmol) and C-(5-(2-fluoro-phenoxy)-thiophen-2-yl)-methylamine described in Preparation Example 161 (33.5mg, 0.15mmol) according to an analogous method to Example

25 A-75. Then, the title compound (7.8mg, 0.023mmol, 15.1%) was obtained as a pale

yellow solid by purifying again by thin layer NH silica gel chromatography (ethyl acetate).

MS m/e (ESI) 344(MH<sup>+</sup>)

5 Example A-149. 6-Amino-N-(5-(2-cyclopropylvinyl)thiophen-2-ylmethyl) nicotinamide

6-Aminonicotinic acid (35mg, 0.251mmol), C-(5-(2-cyclopropylvinyl)thiophen-2-yl)methylamine described in Preparation Example 151 (45mg, 0.251mmol), benzotriazol-1-yl-tris(dimethylamino)phosphonium hexafluorophosphate (133mg, 0.301mmol) and triethylamine (0.042mL, 0.301mmol) were dissolved in N,N-dimethylformamide (3mL), and the solution was stirred at room temperature for 3 hours. Water (10mL) was added to the reaction solution, which was then extracted with ethyl acetate (30mL). The organic layer was washed, dried over anhydrous magnesium sulfate, then, the solvent was evaporated *in vacuo*, the obtained residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 3 : 1), and the title compound (50mg, 0.167mmol, 66.6%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.44-0.52(2H, m), 0.76-0.84(2H, m), 1.45-1.55(1H, m), 4.70(2H, d, J=5.6Hz), 4.76(2H, s), 5.53(1H, dd, J=8.8, 15.6Hz), 6.21(1H, d, J=5.6Hz), 6.49(1H, d, J=8.8Hz), 6.52(1H, d, J=15.6Hz), 6.66(1H, d, J=3.6Hz), 6.83(1H, d, J=3.6Hz), 7.88(1H, dd, J=2.4, 8.8Hz), 8.48(1H, d, J=2.4Hz).

Example A-150. 6-Amino-N-(5-(4-fluoro-phenoxy)-thiophen-2-ylmethyl)-nicotinamide

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.44 (2H, d, J=5.2Hz), 6.39 (1H, d, J=8.8Hz), 6.43-6.52 (3H, m), 6.72 (1H, d, J=2.4Hz), 7.09-7.16 (2H, m), 7.16-7.24 (2H, m), 7.78 (1H, dd, J=2.4, 8.8Hz), 8.42 (1H, s), 8.74 (1H, t, J=5.2Hz).

Example A-151. 6-Amino-N-(5-(3-cyano-phenoxy)-thiophen-2-ylmethyl)-nicotinamide

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.68(2H, d, J=5.6Hz), 4.78(2H, s), 6.27(1H, brs),  
6.38-6.40(1H, m), 6.49-6.52(1H, m), 6.74-6.75(1H, m), 7.08-7.10(2H, m),  
5 7.30-7.33(2H, m), 7.88-7.91(1H, m), 8.48-8.49(1H, m).

Example A-152. N-(5-Benzyl-thiophen-2-ylmethyl)-6-(carbamoylmethyl-amino)-  
nicotinamide

5N Hydrochloric acid (1.5mL) and ethanol (10mL) were added to  
10 (5-((5-benzyl-thiophen-2-ylmethyl)-carbamoyl)-pyridin-2-yl)-carbamoylmethyl-  
carbamic acid *tert*-butyl ester described in Preparation Example A-4 (23mg,  
0.047mmol), and the solution was stirred for 10 minutes at 80°C. An aqueous  
solution of sodium bicarbonate and ethyl acetate were added to the reaction solution,  
which was then partitioned, the organic layer was washed with water twice and dried  
15 over anhydrous magnesium sulfate. The solvent was evaporated, and the title  
compound (12mg, 0.031mmol, 67.1%) was obtained as a brown oily substance.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.00-4.07 (4H, m), 4.46 (2H, d, J=5.6Hz),  
6.55 (1H, d, J=8.8Hz), 6.66-6.69 (1H, m), 6.76 (1H, d, J=3.6Hz), 7.15-7.30 (5H, m),  
7.46 (1H, t, J=6.0Hz), 7.80 (1H, dd, J=2.4, 8.8Hz), 8.44 (1H, d, J=2.4Hz), 8.74 (1H, t,  
20 J=5.6Hz).

Example A-153. 6-(Ethoxymethyl-amino)-N-(5-phenoxy-thiophen-2-ylmethyl)-  
nicotinamide

To a solution of 6-amino-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide  
25 described in Example A-138 (600mg, 1.8mmol) in ethanol (40mL) were added



5,5-dimethylimidazo phospho-2,4-dione (260mg,2.0mmol) and 37% formaldehyde aqueous solution (3mL), and the solution was stirred under reflux for 30 minutes. Water and ethyl acetate were added to the reaction solution, which was then partitioned, and the organic layer was washed with water twice. NH silica gel was added to the organic layer, the solvent was evaporated *in vacuo* for adsorption, purification was carried out by NH silica gel column chromatography (ethyl acetate), and the title compound (457mg, 1.19mmol, 66.1%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) :1.05 (3H, t, J=7.2Hz), 3.42 (2H, q, J=7.2Hz), 4.47 (2H, d, J=5.6Hz), 4.73 (2H, d, J=6.8Hz), 6.44-6.50 (1H, m), 6.55 (1H, d, J=8.8Hz), 6.74 (1H, d, J=3.6Hz), 7.03-7.15 (3H, m), 7.30-7.40 (2H, m), 7.83-7.92 (2H, m), 8.53 (1H, d, J=2.0Hz), 8.85 (1H, t, J=5.6Hz).

#### Example A-154.

6-(Ethoxymethyl-amino)-N-(5-(4-fluoro-phenoxy)-thiophen-2-ylmethyl)-nicotinamide

The title compound (120mg,0.30mmol,51.7%) was obtained as a white solid from (6-amino-N-(5-(4-fluoro-phenoxy)-thiophen-2-ylmethyl)-nicotinamide described in Example A-139 (200mg,0.58mmol) according to an analogous method to Example A-153.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) :1.03-1.10 (3H, m), 3.40-3.46 (2H, m), 4.46 (2H, d, J=5.6Hz), 4.73(2H, d, J=6.8Hz), 6.44-6.49 (1H, m), 6.55 (1H, d, J=8.8Hz), 6.73 (1H, d, J=2.8Hz), 7.08-7.16 (2H, m), 7.16-7.24 (2H, m), 7.83-7.90 (2H, m), 8.52 (1H, s), 8.84 (1H, t, J=5.6Hz).

Example A-155. 6-(Ethoxymethyl-amino)-N-(5-*m*-tolylloxy-thiophen-2-ylmethyl)-

nicotinamide

The title compound (84mg, 2.05mmol, 31.5%) was obtained as a white solid from 6-amino-N-(5-*m*-tolxyloxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-141 (220mg, 0.65mmol) according to an analogous method to Example A-153.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 1.03-1.10 (3H, m), 2.25 (3H, s), 3.40-3.46 (2H, m), 4.46 (2H, d, J=5.6Hz), 4.73 (2H, d, J=6.8Hz), 6.44-6.47 (1H, m), 6.54 (1H, d, J=8.8Hz), 6.73 (1H, d, J=4.0Hz), 6.83-6.96 (3H, m), 7.22 (1H, dd, J=8.0, 8.0Hz), 7.84-7.91 (2H, m), 8.52 (1H, d, J=2.4Hz), 8.84 (1H, t, J=5.6Hz).

Example A-156. N-(5-(3-Fluoro-phenoxy)-thiophen-2-ylmethyl)-6-(2-methoxy-ethylamino)-nicotinamide

To a solution of (5-((5-(3-fluoro-phenoxy)-thiophen-2-ylmethyl)-carbamoyl)-pyridin-2-yl)-carbamic acid *tert*-butyl ester described in Preparation Example A+-2 (100mg, 0.227mmol) and methoxyethyl bromide (38mg, 0.272mmol) in dimethylsulfoxide (5mL) was added sodium hydride (11mg, 0.272mmol, 60% in oil), and the solution was stirred at room temperature for 1 hour. 5N Hydrochloric acid was added to the reaction solution, which was then stirred for 5 minutes at 80°C. Water and ethyl acetate were added to the reaction solution, which was then partitioned, the organic layer was washed with water twice, then, silica gel was added, the solvent was evaporated *in vacuo* for adsorption, purification was carried out by silica gel column chromatography (hexane : ethyl acetate = 1 : 1, then ethyl acetate), and the title compound (39mg, 0.097mmol, 42.8%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.24 (3H, s), 3.41-3.46 (4H, m), 4.47 (2H, d,

J=5.6Hz), 6.48 (1H, d, J=9.2Hz), 6.54-6.58 (1H, m), 6.76 (1H, d, J=2.8Hz), 6.88-7.00 (3H, m), 7.12 (1H, s), 7.36-7.44 (1H, m), 7.77 (1H, d, J=9.2Hz), 8.48 (1H, s), 8.77 (1H, t, J=5.6Hz).

5 Example A-157. 6-Methoxymethyl-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

The title compound (56mg, 0.158mmol, 43.9 %) was obtained as a light brown solid from 6-methoxymethylnicotinic acid (60mg, 0.36mmol) and C-(5-phenoxy-thiophen-2-yl)methylamine described in Preparation Example 26 (64mg, 0.36mmol) according to an analogous technique to Example Q-6.

10 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.38 (3H, s), 4.53-4.57 (4H, m), 6.52 (1H, d, J=3.6Hz), 6.81 (1H, d, J=3.6Hz), 7.07-7.17 (3H, m), 7.35-7.42 (2H, m), 7.50 (1H, d, J=8.0Hz), 8.22 (1H, dd, J=2.0, 8.0Hz), 8.95 (1H, d, J=2.0Hz), 9.31 (1H, t, J=5.6Hz).

Example A-158. N-(5-(3-Chloro-benzyl)-thiophen-2-ylmethyl)-6-

15 (methoxymethyl-amino)-nicotinamide

To a solution of 6-amino-N-(5-(3-chloro-benzyl)-thiophen-2-ylmethyl)-nicotinamide described in Example A-145 (110mg, 0.3mmol) in methanol (10mL) were added 5,5-dimethylimidazo phospho-2,4-dione (43mg, 0.33mmol) and an aqueous solution  
20 of 37% formaldehyde (0.5mL), and the solution was stirred under reflux for 2 hours. NH silica gel was added to the reaction solution, the solvent was evaporated *in vacuo* for adsorption, purification was carried out by NH silica gel column chromatography (hexane : ethyl acetate = 1 : 1, then ethyl acetate), and the title compound (65mg, 0.16mmol, 54%) was obtained as a slightly yellow solid.

25 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.16 (3H, s), 4.06 (2H, s), 4.48 (2H, d,

J=6.0Hz), 4.68 (2H, d, J=6.8Hz), 6.54 (1H, d, J=8.4Hz), 6.69-6.73 (1H, m), 6.76-6.80 (1H, m), 7.17-7.34 (4H, m), 7.84-7.92 (2H, m), 8.51 (1H, s), 8.81 (1H, t, J=6.0Hz).

Example

A-159.

5 N-(5-benzylthiophen-2-ylmethyl)-6-(methoxymethyl-amino)-nicotinamide

To a solution of 6-amino-N-(5-benzylthiophen-2-ylmethyl)-nicotinamide described in Example A-144 (210mg, 0.65mmol) and 5,5-dimethylimidazo phospho-2,4-dione (92mg,0.71mmol) in methanol (15mL) was added an aqueous solution of 37% formaldehyde (2.5mL) under reflux in three time, and the solution  
10 was stirred for 1 hour. NH silica gel was added to the reaction solution, the solvent was evaporated *in vacuo* for adsorption, purification was carried out by NH silica gel column chromatography (hexane : ethyl acetate = 1 : 1, then ethyl acetate), and the title compound (210mg,0.57mmol,87.6%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.14-3.18 (3H, m), 4.03 (2H, s), 4.47 (2H, d, J=4.8Hz), 4.68 (2H, d, J=7.2Hz), 6.54 (1H, d, J=8.8Hz), 6.68 (1H, s), 6.76 (1H, s),  
15 7.14-7.30 (5H, m), 7.84-7.92 (2H, m), 8.51 (1H, s), 8.79 (1H, t, J=4.8Hz).

Example A-160. N-(5-(3-Fluorophenoxy)thiophen-2-ylmethyl)-6-methylnicotinamide

The title compound (53mg, 0.154mmol, 43.0%) was obtained as a white solid  
20 from 6-methylnicotinic acid (49mg,0.36mmol) and C-(5-(3-fluorophenoxy)thiophen-2-yl)methylamine described in Preparation Example 23 (100mg,0.36mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.51 (3H, s), 4.55 (2H, d, J=6.4Hz), 6.58-6.61 (1H, m), 6.82-6.85 (1H, m), 6.92-7.03 (3H, m), 7.34-7.46 (2H, m), 8.10 (1H, dd, J=2.0,  
25 8.0Hz), 8.90 (1H, d, J=2.0Hz), 9.25 (1H, t, J=5.6Hz).

Example A-161. 6-Methyl-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

The title compound (31mg, 0.095mmol, 32.9 %) was obtained as a white solid from 6-methylnicotinic acid (40mg, 0.29mmol) and C-(5-phenoxy-thiophen-2-yl)-methylamine described in Preparation Example 26 (49mg, 0.24mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.49 (3H, s), 4.51 (2H, d, J=5.6Hz), 6.49 (1H, d, J=4.0Hz), 6.78 (1H, d, J=4.0Hz), 7.05-7.15 (3H, m), 7.32-7.40 (3H, m), 8.07 (1H, dd, J=2.0, 8.0Hz), 8.87 (1H, d, J=2.0Hz), 9.21 (1H, t, J=5.6Hz).

Example A-162. N-(5-(3-Chloro-benzyl)-thiophen-2-ylmethyl)-6-methyl-nicotinamide

The title compound (32mg, 0.089mmol, 26.4%) was obtained as a white solid from C-(5-(3-chloro-benzyl-2-yl)-methylamine described in Preparation Example 45 (80mg, 0.34mmol) and 6-methylnicotinic acid (46mg, 0.34mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.50 (3H, s), 4.08 (2H, s), 4.54 (2H, d, 5.6Hz), 6.75 (1H, d, J=3.6Hz), 6.84 (1H, d, J=3.6Hz), 7.20-7.36 (5H, m), 8.08 (1H, dd, J=2.0, 8.0Hz), 8.88 (1H, d, J=2.0Hz), 9.18 (1H, t, J=5.6Hz).

Example A-163. 6-Methylamino-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

To a solution of 6-(ethoxymethyl-amino)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-153 (427mg, 0.92mmol) in dimethylsulfoxide (5mL) was added sodium borohydride (100mg, 2.7mmol), and the solution was stirred for 15 minutes at 100°C. Water and ethyl acetate were added to the reaction solution, which was then

partitioned, and the organic layer was washed with water three times. Silica gel was added to the organic layer, the solvent was evaporated *in vacuo* for adsorption, and purified by silica gel column chromatography (ethyl acetate). The solvent was evaporated *in vacuo*, hexane, diethyl ether and ethyl acetate were added to the residue for solidification, and the title compound (150mg, 0.44mmol, 48.1%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.78 (3H, d, J=4.8Hz), 4.46 (2H, d, J=5.6Hz), 6.40 (1H, d, J=8.8Hz), 6.47 (1H, d, J=3.6Hz), 6.73 (1H, d, J=3.6Hz), 6.98-7.15 (4H, m), 7.32-7.40 (2H, m), 7.79 (1H, dd, J=2.0, 8.8Hz), 8.50 (1H, d, J=2.0Hz), 8.75 (1H, t, J=5.6Hz).

Example A-164. N-(5-(4-Fluoro-phenoxy)-thiophen-2-ylmethyl)-6-methylamino-nicotinamide

The title compound (120mg, 0.30mmol, 51.7%) was obtained as a white solid from 6-(ethoxymethyl-amino)-N-(5-(4-fluoro-phenoxy)-thiophen-2-ylmethyl)-nicotinamide described in Example A-154 according to an analogous method to Example A-163.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.76-2.81 (3H, m), 4.45 (2H, d, J=5.6Hz), 6.40 (1H, d, J=8.4Hz), 6.44-6.48 (1H, m), 6.70-6.75 (1H, m), 6.98-7.06 (1H, m), 7.08-7.24 (4H, m), 7.79 (1H, d, J=8.4Hz), 8.50 (1H, s), 8.75 (1H, t, J=5.6Hz).

Example A-165. N-(5-(3-Fluoro-phenoxy)-thiophen-2-ylmethyl)-6-methylamino-nicotinamide

Ethanol (20mL) and 5N hydrochloric acid (0.8mL) were added to (5-((5-(3-fluoro-phenoxy)-thiophen-2-ylmethyl)carbamoyl)pyridin-2-yl)methyl-

carbamic acid *tert*-butyl ester described in Preparation Example A+-3 (87mg, 0.19mmol), and the solution was stirred for 25 minutes at 80°C. An aqueous solution of sodium bicarbonate and ethyl acetate were added to the reaction solution, which was then partitioned, the organic layer was washed with water twice, washed  
 5 with brine, and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated, and the title compound (58mg, 0.162mmol, 85.5%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.78 (3H, d, J=4.8Hz), 4.47 (2H, d, J=5.6Hz), 6.40 (1H, d, J=8.0Hz), 6.55 (1H, d, J=3.6Hz), 6.76 (1H, d, J=3.6Hz), 6.86-7.06 (4H,  
 10 m), 7.36-7.44 (1H, m), 7.79 (1H, dd, J=2.0, 8.0Hz), 8.50 (1H, d, J=2.0Hz), 8.77 (1H, t, J=5.6Hz).

Example A-166. N-(5-Benzyl-thiophen-2-ylmethyl)-6-methylamino-nicotinamide

To a solution of  
 15 N-(5-benzyl-thiophen-2-ylmethyl)-6-(methoxymethyl-amino)-nicotinamide described in Example A-159 (183mg, 0.52mmol) in dimethylsulfoxide (3mL) was added sodium borohydride (120mg, 3.12mmol), and the solution was stirred at 140°C for 5 hours. Water and ethyl acetate were added to the reaction solution, which was then partitioned, and the organic layer was washed with water twice. Silica gel was  
 20 added to the organic layer, the solvent was evaporated *in vacuo* for adsorption, purification was carried out by silica gel column chromatography (ethyl acetate, then ethyl acetate : methanol = 10 : 1), and the title compound (68mg, 0.20mmol, 38.8%) was obtained as a green solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.77 (3H, d, J=4.8Hz), 4.03 (2H, s), 4.46 (2H,  
 25 d, J=5.6Hz), 6.39 (1H, d, J=8.8Hz), 6.68 (1H, d, J=3.6Hz), 6.76 (1H, d, J=3.6Hz),

6.96-7.04 (1H, m), 7.15-7.32 (5H, m), 7.77 (1H, dd, J=2.0, 8.8Hz), 8.49 (1H, d, J=2.0Hz), 8.70 (1H, t, J=5.6Hz).

**Example A-167. 6-Methylamino-N-(5-*m*-tolxyloxy-thiophen-2-ylmethyl)-nicotinamide**

5 To a solution of  
6-(ethoxymethyl-amino)-N-(5-*m*-tolylloxy-thiophen-2-ylmethyl)-nicotinamide described  
in Example A-155 (70mg, 0.17mmol) in dimethylsulfoxide (5mL) was added sodium  
borohydride (20mg, 0.51mmol), and the solution was stirred for 20 minutes at 120°C.  
Water and ethyl acetate were added to the reaction solution, which was then  
10 partitioned, the organic layer was washed with brine once. The organic layer was  
passed through a glass filter lined with NH silica gel, and eluted thoroughly with ethyl  
acetate. The solvent was evaporated *in vacuo*, and the title compound (53mg,  
0.15mmol, 88.2%) was obtained as a light brown solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.25 (3H, s), 2.77 (3H, d, J=4.4Hz), 4.45 (2H, d, J=5.6Hz), 6.40 (1H, d, J=8.8Hz), 6.45 (1H, d, J=3.6Hz), 6.72 (1H, d, J=3.6Hz), 6.83-6.96 (3H, m), 7.02 (1H, q, J=4.4Hz), 7.22 (1H, dd, J=8.0, 8.0Hz), 7.78 (1H, dd, J=2.4, 8.8Hz), 8.50 (1H, d, J=2.4Hz), 8.75 (1H, t, J=5.6Hz).

**Example A-168. 2-Amino-N-(5-phenoxy-thiophen-2-ylmethyl)-6-vinyl-nicotinamide**

20 To a mixture of  
2-amino-6-chloro-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in  
Example A-101 (30mg, 0.083mmol), tetrakis(triphenylphosphine)palladium(0) (19mg,  
0.017mmol) and xylene (1.5mL) was added vinyl(tri-butyl)tin (0.073mL, 0.25mmol),  
and the solution was stirred at 130°C for 3 hours. After cooling, water and ethyl  
25 acetate were added to the reaction solution for extraction, which was then washed



with brine. The solvent was evaporated *in vacuo*, then, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 2 : 1), and the title compound (19mg, 0.054mmol, 65%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.65 (2H, d, J=5.5Hz), 5.49 (1H, dd, J=1.4, 10.7Hz), 6.22 (1H, dd, J=1.4, 17.3Hz), 6.27 (1H, brs), 6.36 (2H, brs), 6.39 (1H, d, J=3.8Hz), 6.60-6.67 (2H, m), 6.74 (1H, d, J=3.7Hz), 7.08-7.13 (3H, m), 7.30-7.35 (2H, m), 7.56 (1H, d, J=8.1Hz).

Example A-169. 2-Amino-N-(5-bromo-4-phenoxy-thiophen-2-ylmethyl)-nicotinamide

The title compound was obtained from C-(5-bromo-4-phenoxy-thiophen-2-yl)-methylamine described in Preparation Example 129 (500mg, 1.76mmol) and 2-amino-nicotinic acid (267mg, 1.94mmol) according to an analogous method to Example A-26. Trifluoroacetic acid salt of the title compound was obtained by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used).

MS m/e (ESI) 405.94(MH<sup>+</sup>)

Example A-170. 2-Amino-N-(5-methyl-4-phenoxy-thiophen-2-ylmethyl)-nicotinamide

To a mixture of 2-amino-N-(5-bromo-4-phenoxy-thiophen-2-ylmethyl)-nicotinamide described in Example A-169 (30mg, 0.074mmol), dichloro(1,1'-bis(diphenylphosphino)ferrocene) nickel(II) (10mg, 0.015mmol) and tetrahydrofuran (1mL) was added methylmagnesium bromide (638μl, 0.592mmol) at room temperature, the solution was stirred for 2 hours at room temperature, and was further stirred for 2 hours at 50°C.

After cooling, water and ethyl acetate were added for extraction, and the solution was washed with brine, then, the solvent was evaporated *in vacuo*. The residue was purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), and the title  
5 compound was obtained as a trifluoroacetic acid salt.

MS m/e (ESI) 340.09(MH<sup>+</sup>)

Example A-171. 2-Amino-N-(4-chloro-5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

To a mixture of 2-amino-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide  
10 described in Example A-67 (21mg, 0.064mmol) and N,N-dimethylformamide (1mL) was added N-chlorosuccinimide (13mg, 0.096mmol), and the solution was stirred overnight at room temperature. The reaction solution was purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), and trifluoroacetic acid salt of the title compound  
15 (7.4mg, 0.016mmol, 24%) was obtained.

MS m/e (ESI) 360.1 (MH<sup>+</sup>)

Example A-172. 2-Amino-5-chloro-N-(4-chloro-5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

20 Trifluoroacetic acid salt of the title compound (1.5mg, 0.0030mmol, 4.6%) was obtained as a by-product of Example A-171.

MS m/e (ESI) 394.0 (MH<sup>+</sup>)

Example A-173. 2,6-Diamino-N-(4-(3-chloro-benzyloxy)-benzyl)-nicotinamide

25 MS m/e (ESI) 383.246 (MH<sup>+</sup>)

Example A-174. 2,6-Diamino-N-(5-benzyloxy-pyridin-2-ylmethyl)-nicotinamide

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.39(2H, d, J=5.9Hz), 5.16(2H, s), 5.68(1H, d, J=8.6Hz), 6.09(2H, s), 6.94(2H, s), 7.20(1H, d, J=8.8Hz), 7.31-7.46(6H, m), 7.69(1H, d, J=8.6Hz), 8.25(1H, d, J=2.9Hz), 8.35(1H, t, J=5.9Hz).

Example A-175. 2,6-Diamino-N-(5-phenoxy-methyl-pyridin-2-ylmethyl)-nicotinamide

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.64(2H, s), 4.68(2H, d, J=4.9Hz), 5.07(2H, s), 5.80(1H, d, J=8.4Hz), 6.50(2H, s), 6.96-7.01(3H, m), 7.26-7.35(4H, m), 7.56(1H, d, J=8.4Hz), 7.76(1H, dd, J=2.2Hz, 8.0Hz), 8.61(1H, d, J=1.7Hz).

Example A-176. 2-Amino-6-(2-cyano-ethyl)-N-(4-(6-fluoro-pyridin-2-yloxy-methyl)-benzyl)-nicotinamide

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 2.78(2H, t, J=7.2Hz), 2.94(2H, t, J=7.2Hz), 4.60(2H, d, J=5.6Hz), 5.33(2H, s), 6.24(1H, brs), 6.39(2H, brs), 6.47-6.48(1H, m), 6.49-6.50(1H, m), 6.64-6.67(1H, m), 7.35(2H, d, J=8.4Hz), 7.46(2H, d, J=8.4Hz), 7.53(1H, d, J=8.0Hz), 7.66(1H, q, J=8.0Hz).

Example A-177. 2-Amino-6-(2-fluoro-ethyl)-N-(5-phenoxy-thiophen-2-ylmethyl)-nicotinamide

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.93 (2H, td, J=6.0, 25.6 Hz), 4.47 (2H, d, J=5.6 Hz), 4.75 (2H, td, J=6.0, 47.2 Hz), 6.50 (1H, d, J=4.0 Hz), 6.53 (1H, d, J=8.0 Hz), 6.77 (1H, d, J=4.0 Hz), 7.07-7.18 (5H, m), 7.38 (2H, t, J=8.0 Hz), 7.85 (1H, d, J=8.0 Hz), 8.99 (1H, t, J=5.6 Hz).

Example A-178. 2-Amino-6-ethoxymethyl-N-(4-(pyridin-2-yloxymethyl)-benzyl)-nicotinamide

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 1.27(3H, t, J=7.0Hz), 3.60(2H, q, J=7.2Hz), 4.44(2H, s), 4.60(2H, d, 5.6Hz), 5.38(2H, s), 6.24(1H, brs), 6.38(2H, brs), 6.73(1H, d, J=8.0Hz), 6.80(1H, dd, J=0.8, 7.6Hz), 6.88-6.91(1H, m), 7.36(2H, d, J=8.0Hz), 7.46(2H, d, J=8.0Hz), 7.56-7.61(2H, m), 8.17-8.18(1H, m).

Example A-179.

2-Amino-N-(5-isopropoxymethyl-thiophen-2-ylmethyl)-6-methoxymethyl-nicotinamide

10 MS m/e (ESI) 350 (MH<sup>+</sup>)

Example A-180. 2-Amino-N-(4-(3-methoxy-benzyloxy)-benzyl)-6-methoxymethyl-nicotinamide

MS m/e (ESI) 408.27 (MH<sup>+</sup>)

15 <sup>1</sup>H-NMR Spectrum (CD<sub>3</sub>OD) δ(ppm) : 3.50(3H, s), 3.77(3H, s), 4.48(2H, d, J=4Hz), 4.57(2H, s), 5.04(2H, s), 6.84-6.88(2H, m), 6.94-6.98(4H, m), 7.23-7.29(3H, m), 8.27-8.29(1H, m).

Example A-181. 2-Amino-N-(4-(3-chloro-benzyloxy)-benzyl)-6-

20 methoxymethyl-nicotinamide

MS m/e (ESI) 412.26(MH<sup>+</sup>)

Example A-182. 2-Amino-N-(4-butoxymethyl-benzyl)-6-methoxymethyl-nicotinamide

MS m/e (ESI) 358 (MH<sup>+</sup>)

25

Example A-183. 2-Amino-6-methoxymethyl-N-(4-propoxymethyl-benzyl)-nicotinamide

MS m/e (ESI) 344 (MH<sup>+</sup>)

Example A-184. 2-Amino-N-(3-cyclopropylmethoxy-benzyl)-6-methoxymethyl-

5 nicotinamide

MS m/e (ESI) 342 (MH<sup>+</sup>)

Example A-185.

2-Amino-N-(4-(6-fluoro-pyridin-2-yloxymethyl)-benzyl)-6-methoxymethyl-

10 nicotinamide

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.46(3H, s), 4.40(2H, s), 4.61(2H, d, J=5.6Hz), 5.33(2H, s), 6.26(1H, brs), 6.40(2H, s), 6.49(1H, dd, J=2.4, 7.6Hz), 6.66(1H, dd, J=1.1, 7.6Hz), 6.70(1H, d, J=8.0Hz), 7.36(2H, d, J=8.0Hz), 7.46(2H, d, J=8.0Hz), 7.60(1H, d, J=8.0Hz), 7.66(1H, dd, J=8.0, 8.0Hz).

15

Example A-186. 2-Amino—6-methoxymethyl-N-(4-(4-methyl-pyridin-2-yloxymethyl)-benzyl)-nicotinamide

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 2.30(3H, s), 3.45(3H, s), 4.39(2H, s), 4.60(2H, d, J=5.6Hz), 5.36(2H, s), 6.25(1H, brs), 6.40(2H, brs), 6.62(1H, s), 6.65(1H, d, J=8.0Hz), 6.72(1H, d, J=5.2Hz), 7.35(2H, d, J=8.4Hz), 7.45(2H, d, J=8.4Hz), 7.59(1H, d, J=8.0Hz), 8.02(1H, d, J=5.2Hz).

20

Example A-187. 2-Amino-N-(4-(6-fluoro-pyridin-2-ylmethoxy)-benzyl)-6-methoxymethyl-nicotinamide

25 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.45(3H, s), 4.39(2H, s), 4.53(2H, d, J=5.6Hz),

5.13(2H, s), 6.22(1H, brs), 6.39(2H, brs), 6.69(1H, d, J=7.6Hz), 6.87(1H, dd, J=2.8, 8.4Hz), 6.96(2H, d, J=8.4Hz), 7.28(2H, d, J=8.4Hz), 7.40-7.42(1H, m), 7.58(1H, d, J=8.0Hz), 7.81(1H, dd, J=8.0, 8.0Hz).

5 Example A-188. 2-Amino-N-(4-butoxy-benzyl)-6-methoxymethyl-nicotinamide

MS m/e (ESI) 344 (MH<sup>+</sup>)

Example A-189. 2-Amino-N-(4-(2-ethoxy-ethyl)-benzyl)-6-methoxymethyl-nicotinamide

10 MS m/e (ESI) 344 (MH<sup>+</sup>)

Example A-190. 2-Amino-6-methoxymethyl-N-(4-(5-methyl-pyridin-2-ylloxymethyl)-benzyl)-nicotinamide

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 2.25(3H, s), 3.45(3H, s), 4.40(2H, s), 4.60(2H, d, J=5.6Hz), 5.34(2H, s), 6.24(1H, brs), 6.40(2H, brs), 6.69(1H, d, J=8.0Hz), 6.71(1H, d, J=8.0Hz), 7.35(2H, d, J=8.0Hz), 7.41(1H, dd, J=2.4, 8.4Hz), 7.46(2H, d, J=8.0Hz), 7.59(1H, d, J=8.4Hz), 7.96(1H, s).

Example A-191. 2-Amino-N-(6-benzyl-pyridin-3-ylmethyl)-6-methoxymethyl-

20 nicotinamide

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.45(3H, s), 4.15(2H, s), 4.39(2H, s), 4.57(2H, d, J=5.6Hz), 6.32(1H, brs), 6.38(2H, brs), 6.69(1H, d, J=8.0Hz), 7.12(1H, d, J=8.4Hz), 7.20-7.32(5H, m), 7.58(7.61(2H, m), 8.52(1H, d, J=2.0Hz).

25 Example A-192. 2-Amino-6-methoxymethyl-N-(5-phenoxy-methyl-pyridin-2-ylmethyl)-

nicotinamide

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.46(3H, s), 4.41(2H, s), 4.72(2H, d, J=4.8Hz), 5.09(2H, s), 6.45(2H, s), 6.74(1H, d, J=7.9Hz), 6.96-7.02(3H, m), 7.29-7.36(3H, m), 7.53(1H, s), 7.79(2H, d, J=7.9Hz), 8.63(1H, d, J=1.7Hz).

5 Example A-193. 2-Amino-6-methoxymethyl-N-(5-phenoxy-pyridin-2-ylmethyl)-nicotinamide

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.34 (3H, s), 4.30 (2H, s), 4.52 (2H, d, J=6.0 Hz), 6.62 (1H, d, J=8.0 Hz), 7.02-7.07 (2H, m), 7.10-7.20 (3H, m), 7.33-7.46 (4H, m), 8.02 (1H, d, J=8.0 Hz), 8.31 (1H, d, J=2.8 Hz), 9.02 (1H, t, J=6.0 Hz).

10

Example A-194. 2-Amino-6-methoxymethyl-N-(4-(2-propoxy-ethyl)-benzyl)-nicotinamide

MS m/e (ESI) 358 (MH<sup>+</sup>)

15 Example AA-1. 3-Amino-pyrazine-2-carboxylic acid  
(5-phenoxy-thiophen-2-ylmethyl)-amide

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.47 (2H, d, J=6.4 Hz), 6.47 (1H, d, J=4.0 Hz), 6.74 (1H, d, J=4.0 Hz), 7.05-7.15 (3H, m), 7.24-7.40 (2H, m), 7.50 (2H, brs), 7.80 (1H, d, J=2.4 Hz), 8.18 (1H, d, J=2.4 Hz), 9.28 (1H, t, J=6.4 Hz).

20

Example AA-2. 3,5-Diamino-pyrazine-2-carboxylic acid  
4-(pyridin-2-yloxymethyl)-benzylamide

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.38(2H, d, J=6.4Hz), 5.3(2H, s), 6.67(2H, s), 6.84(1H, d, J=8.2Hz), 6.98(1H, t, J=7.2Hz), 7.13(1H, s), 7.28(2H, d, J=8.1Hz), 7.38(2H, d, J=8.1Hz), 7.71(1H, dt, J=2.0Hz, 7.8Hz), 8.16(1H, dd, J=2.0Hz, 4.8Hz),

25

8.54(1H, t, J=6.4Hz).

Example B-1. 4-Amino-pyrimidine-5-carboxylic acid  
(5-phenoxy-thiophen-2-ylmethyl)-amide

5 The title compound (7mg, 0.021mmol, 4.4 %) was obtained as a white solid from 4-amino-pyrimidine-5-carboxylic acid (68mg, 0.49mmol) and C-(5-phenoxy-thiophen-2-yl)-methylamine described in Preparation Example 26 (100mg, 0.49mmol) according to an analogous technique to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.47 (2H, d, J=5.2Hz), 6.47-6.50 (1H, m),  
 10 6.76-6.78 (1H, m), 7.04-7.15 (3H, m), 7.34-7.40 (2H, m), 7.77 (2H, brs), 8.40 (1H, d, J=1.6Hz), 8.59 (1H, d, J=1.6Hz), 9.19 (1H, t, J=5.2Hz).

Example B-2. 4-Amino-2-propylamino-pyrimidine-5-carboxylic acid  
4-(pyridin-2-yloxymethyl)-benzylamide

15 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 0.86 (3H, t, J=7.2 Hz), 1.44-1.55 (2H, m), 3.14-3.24 (2H, m), 4.38 (2H, d, J=6.4 Hz), 5.32 (2H, s), 6.83-6.87 (1H, m), 6.96-7.01 (1H, m), 7.28 (2H, d, J=8.0 Hz), 7.38 (2H, d, J=8.0 Hz), 7.69-7.75 (1H, m), 8.15-8.19 (1H, m), 8.42 (1H, s), 8.60 (1H, brs).

20 Example C-1. (2-Amino-6,7-dihydro-5H-cyclopenta[b]pyridine-3-carboxylic acid  
(5-phenoxy-thiophen-2-ylmethyl)-amide

2-Amino-6,7-dihydro-5H-cyclopenta[b]pyridine-3-carboxylic acid described in Preparation Example C-5 (100mg, 0.56mmol), triethylamine (0.188mL, 1.35mmol) and benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate  
 25 (298mg, 0.67mmol) were dissolved in N,N-dimethylformamide (3mL),



C-(5-phenoxy-thiophen-2-yl)-methylaniline described in Preparation Example 26 (138mg, 0.67mmol) was added, followed by stirring for 15 hours 20 minutes at room temperature. After the reaction was completed, the reaction solution was poured into brine, and the solution was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, then concentrated, the obtained residue was purified by silica gel column chromatography (ethyl acetate : hexane = 1 : 1), and the title compound (77mg, 0.21mmol, 38%) was obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 1.95-2.03 (2H, m), 2.68-2.76 (4H, m), 4.46 (2H, d, J=5.6Hz), 6.45 (1H, d, J=4.0Hz), 6.76 (1H, d, J=4.0Hz), 6.97 (2H, brs), 7.07-7.17 (3H, m), 7.36-7.41 (2H, m), 7.76 (1H, s), 8.88-8.93 (1H, m).

#### Example D-1. [1,5]Naphthylidene-2-carboxylic acid 3-phenoxybenzylamide

The title compound (14mg, 76%) was obtained as a colorless oil from [1,5]naphthylidene-2-carboxylic acid described in Preparation Example D-1 (9mg, 0.0517mmol) and 3-phenoxybenzylamine described in Preparation Example 4 (6mg, 0.0517mmol) according to an analogous method to Example L-4.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.73(2H, d, J=6.0Hz), 6.92-6.94(1H, m), 7.01-7.03(2H, m), 7.07-7.17(3H, m), 7.26-7.36(3H, m), 7.70(1H, dd, J=4.4, 8.8Hz), 8.39(1H, dd, J=1.6, 8.8Hz), 8.51(1H, brs), 8.57(2H, s), 9.06(1H, dd, J=1.6, 4.4Hz).

#### Example D-2. [1,5]Naphthylidene-2-carboxylic acid 4-benzyloxybenzylamide

The title compound (11mg, 52%) was obtained as a white solid from [1,5]naphthylidene-2-carboxylic acid described in Preparation Example D-1 (10mg, 0.0574mmol) and 4-benzyloxybenzylamine described in Preparation Example 1 (12mg, 0.0574mmol) according to an analogous method to Example L-4.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.68(2H, d, J=6.4Hz), 5.08(2H, s), 6.97-6.99(2H, m), 7.32-7.45(7H, m), 7.69(1H, dd, J=4.4, 8.8Hz), 8.36-8.39(1H, m), 8.43(1H, brs), 8.54-8.60(2H, m), 9.04-9.06(1H, m).

5 Example D-3. [1,5]Naphthylidene-2-carboxylic acid 4-phenoxyethyl-benzylamide

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.76(2H, d, J=6Hz), 5.07(2H, s), 6.94-6.99(3H, m), 7.27-7.31(3H, m), 7.43-7.45(4H, m), 7.68-7.71(1H, m), 8.37-8.40(1H, m), 8.50(1H, brs), 8.55-8.60(1H, m), 9.05-9.06(1H, m).

10 Example D-4. [1,5]Naphthylidene-2-carboxylic acid  
(1-(3-fluoro-benzyl)-1H-pyrrol-3-ylmethyl)-amide

To a solution of C-(1-(3-fluoro-benzyl)-1H-pyrrol-3-yl)-methylamine described in Preparation Example 59 (59mg, 0.287mmol) and [1,5]naphthylidene-2-carboxylic acid described in Preparation Example D-1 (50mg, 0.287mmol) in  
 15 N,N-dimethylformamide (3mL) were added benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (152mg, 0.344mmol) and triethylamine (80μL, 0.574mmol), and the solution was stirred overnight at room temperature. Ethyl acetate and water were added to the reaction solution, which was then partitioned, the organic layer was washed with water, and  
 20 then, dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate), and the title compound (49mg, 47%) was obtained as a pale yellow solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>)δ(ppm) : 4.59(2H, d, J=5.6Hz), 5.03(2H, s), 6.26-6.27(1H, m), 6.66-6.67(1H, m), 6.74-6.75(1H, m), 6.81-6.84(1H, m), 6.92-6.94(1H, m),  
 25 6.96-7.01(1H, m), 7.28-7.33(1H, m), 7.67-7.70(1H, m), 8.31(1H, brs), 8.37-8.40(1H,

m), 8.53-8.59(2H, m), 9.03-9.05(1H, m).

Example D-5. [1,5]Naphthylidene-2-carboxylic acid  
(5-(3-fluorophenoxy)thiophen-2-ylmethyl) amide

5 The title compound (66mg, 0.17mmol, 72.5%) was obtained as a brown oil from [1,5]-naphthylidene-2-carboxylic acid (42mg, 0.24mmol) and C-(5-(3-fluorophenoxy)thiophen-2-yl)methylamine described in Preparation Example 23 (54mg,0.24mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.61 (2H, d, J=6.4Hz), 6.57 (1H, d, J=3.6Hz),  
 10 6.83 (1H, d, J=3.6Hz), 6.87-7.00 (3H, m), 7.36-7.42 (1H, m), 7.87 (1H, dd, J=4.0, 8.8Hz), 8.37 (1H, d, J=8.8Hz), 8.49 (1H, d, J=8.8Hz), 8.59 (1H, d, J=8.8Hz), 9.09 (1H, d, J=4.0Hz), 9.58 (1H, t, J=6.4Hz).

Example D-6. [1,5]Naphthylidene-2-carboxylic acid  
 15 (5-phenoxy-thiophen-2-ylmethyl)-amide

The title compound (15mg, 0.042mmol, 17.3 %) was obtained as a light brown solid from [1,5]-naphthylidene-2-carboxylic acid (42mg,0.24mmol) and C-(5-phenoxy-thiophen-2-yl)methylamine described in Preparation Example 26 (49mg,0.24mmol) according to an analogous technique to Example Q-6.

20 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.59 (2H, d, J=5.6Hz), 6.49 (1H, d, J=3.6Hz), 6.81 (1H, d, J=3.6Hz), 7.03-7.14 (3H, m), 7.30-7.40 (2H, m), 7.88 (1H, dd, J=4.0, 8.8Hz), 8.36 (1H, d, J=8.8Hz), 8.48 (1H, d, J=8.8Hz), 8.58 (1H, d, J=8.8Hz), 9.09 (1H, d, J=4.0Hz), 9.56 (1H, t, J=5.6Hz).

25 Example D-7. [1,5]Naphthylidene-2-carboxylic acid

(5-(4-fluoro-phenoxy)-thiophen-2-ylmethyl)-amide

To a solution of [1,5]naphthylidene-2-carboxylic acid described in Preparation Example D-1 (20mg, 0.115mmol) and described in Preparation Example 28C-(5-(4-fluorophenoxy)thiophen-2-yl)methylamine) (51mg, 0.23mmol) in dimethylsulfoxide (9mL) were added benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (102mg, 0.23mmol) and triethylamine (56μL, 0.43mmol), and the solution was stirred at 60°C for 30 minutes. Ethyl acetate and water were added to the reaction solution, which was then partitioned, the organic layer was sequentially washed with an aqueous solution of saturated sodium bicarbonate, water and brine, and then, dried over anhydrous sodium sulfate. The solvent was evaporated, the residue was sequentially purified by silica gel column chromatography (hexane : ethyl acetate), silica gel column chromatography (dichloromethane : ethyl acetate), reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), and the title compound (8.6mg, 0.017mmol, 15%) was obtained.

MS m/e (ESI) 379.76(MH<sup>+</sup>)

<sup>1</sup>H-NMR Spectrum (CD<sub>3</sub>OD) δ(ppm) : 4.71(2H, d, J=6.4Hz), 6.38(1H, d, J=4.0Hz), 6.83 (1H, d, J=4.0Hz), 7.02-7.09 (4H, m), 7.85 (1H, dd, J=4.0, 8.8Hz), 8.46 (1H, d, J=8.8Hz), 8.56-8.61(2H, m), 9.04(1H, dd, J=1.6, 4.0Hz), 9.55-9.64 (1H, m).

Example D-8. [1,5]Naphthylidene-2-carboxylic acid(5-(3-chloro-phenoxy)-thiophen-2-ylmethyl)-amide

MS m/e (ESI) 396.28 (MH<sup>+</sup>)

Example D-9. [1,5]Naphthylidene-2-carboxylic acid  
(5-benzofuran-2-ylmethyl-thiophen-2-ylmethyl)-amide

MS m/e (ESI) 400.51 (MH<sup>+</sup>)

5 Example D-10. [1,5]Naphthylidene-2-carboxylic acid  
4-(6-fluoro-pyridin-2-yloxymethyl)-benzylamide

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.75(2H, d, J=6.0Hz), 5.34(2H, s), 6.47-6.49(1H, m), 6.64-6.66(1H, m), 7.43-7.48(4H, m), 7.62-7.70(2H, m), 8.36-8.39(1H, m), 8.50(1H, brs), 8.55-8.60(2H, m), 9.04-9.06(1H, m).

10

Example E-1. Quinoline-6-carboxylic acid (5-benzyl-furan-2-yl)-amide

The title compound (200mg, 5.8mmol, 54.6%) was obtained as a slightly yellow solid from 5-benzylfuran-2-carbaldehyde described in Preparation Example 39 and 6-quinolinecarboxylic acid according to an analogous method to Example Q-1.

15 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.92 (2H, s), 4.45 (2H, d, J=5.6Hz), 6.02 (1H, d, J=2.8Hz), 6.20 (1H, d, J=2.8Hz), 7.16-7.30 (5H, m), 7.59 (1H, dd, J=4.4, 8.0Hz), 8.05 (1H, d, J=8.8Hz), 8.16 (1H, dd, J=2.0, 8.8Hz), 8.44 (1H, dd, J=1.6, 8.0Hz), 8.50 (1H, d, J=2.0Hz), 8.97 (1H, dd, J=1.6, 4.4Hz), 9.16 (1H, t, J=5.6Hz).

20 Example E-2. Quinoline-6-carboxylic acid (5-phenoxy-furan-2-ylmethyl)-amide

The title compound (68mg, 0.197mmol, 17.2%) was obtained as a white solid from quinoline-6-carboxylic acid (200mg, 1.15mmol) and C-(5-phenoxy-furan-2-yl)-methylamine described in Preparation Example 49 (200mg, 1.12mmol) according to an analogous method to Example Q-6.

25 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.45 (2H, d, J=5.6Hz), 5.72 (1H, d, J=3.2Hz),

6.34 (1H, d, J=3.2Hz), 7.02-7.08 (2H, m), 7.11-7.17 (1H, m), 7.14-7.40 (2H, m), 7.60 (1H, dd, J=2.0, 8.0Hz), 8.06 (1H, d, J=8.8Hz), 8.16 (1H, d, J=8.8Hz), 8.45 (1H, d, J=8.0Hz), 8.51 (1H, s), 8.97 (1H, d, J=4.0Hz), 9.18 (1H, t, J=5.6Hz).

5 Example E-3. Quinoline-6-carboxylic acid  
(5-(3-fluoro-phenoxy)-furan-2-ylmethyl)-amide

To a solution of (5-(3-fluoro-phenoxy)-furan-2-yl)-methanol described in Preparation Example 51 (1.5g, 7.2mmol), phthalimide (1.1g, 7.2mmol) and triphenyl phosphine (1.9g, 7.2mmol) in tetrahydrofuran (10mL) was added diethyl  
 10 azodicarboxylate (3.5g, 7.9mmol) dropwise at 0°C, and the solution was stirred at room temperature for 30 minutes. Silica gel was added to the reaction solution, the solvent was evaporated *in vacuo* for adsorption, purification was carried out by silica gel column chromatography (hexane : ethyl acetate = 2 : 1), and a white solid (700mg) was obtained. Ethanol (10mL) and hydrazine monohydrate (0.3mL) were  
 15 added to this solid (700mg), and stirred for 15 minutes at 90°C. The solution was allowed to room temperature, the solid was removed by filtration, and a pale yellow oil (360mg) containing C-(5-(3-fluoro-phenoxy)-furan-2-yl)-methylamine was obtained. The title compound (17mg, 0.046mmol, 2.7%) was obtained as a brown solid from this oil (360mg) and quinoline-6-carboxylic acid (300mg, 1.7mmol) according to an  
 20 analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.46 (2H, d, J=5.2Hz), 5.83 (1H, d, J=3.2Hz), 6.36 (1H, d, J=3.2Hz), 6.87-7.03 (3H, m), 7.38-7.45 (1H, m), 7.60 (1H, d, J=4.0, 8.4Hz), 8.06 (1H, d, J=8.8Hz), 8.16 (1H, dd, J=2.0, 8.8Hz), 8.45 (1H, dd, J=1.6, 8.4Hz), 8.51(1H, d, J=2.0Hz), 8.97(1H, dd, J=1.6, 4.0Hz), 9.19(1H, t, J=5.2Hz).

Example E-4. Quinoline-6-carboxylic acid (5-phenyl-furan-2-ylmethyl)-amide

To a solution of quinoline-6-carboxylic acid (5-bromo-furan-2-ylmethyl)-amide described in Preparation Example E+2 (200mg, 0.60mmol) in 1,4-dioxane (5mL) were added phenylboronic acid (150mg, 1.2mmol),  
 5 tetrakis(triphenylphosphine)palladium(0) (55mg, 0.047mmol) and an aqueous solution of potassium carbonate (2mol), and the solution was stirred at 110°C for 2 hours. Water and ethyl acetate were added to the reaction solution, which was then partitioned, NH silica gel was added to the organic layer, the solvent was evaporated *in vacuo* for adsorption, purification was carried out by NH silica gel column  
 10 chromatography (hexane : ethyl acetate = 2 : 1), and the title compound (65mg, 0.198mmol, 33.0%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.59 (2H, d, J=6.0Hz), 6.44 (1H, d, J=3.2Hz), 6.88 (1H, d, J=3.2Hz), 7.23-7.28 (1H, m), 7.37-7.42 (2H, m), 7.60 (1H, dd, J=4.0, 8.4Hz), 7.64-7.70 (2H, m), 8.07 (1H, d, J=8.8Hz), 8.20 (1H, dd, J=2.0, 8.8Hz),  
 15 8.45-8.50 (1H, m), 8.55 (1H, d, J=2.0Hz), 8.97 (1H, dd, J=1.6, 4.0Hz), 9.26 (1H, t, J=6.0Hz).

Example E-5. Quinoline-6-carboxylic acid  
(5-(2,4-difluoro-phenoxy)-furan-2-ylmethyl)-amide

20 The title compound (71mg, 0.187mmol, 18%) was obtained from quinoline-6-carboxylic acid (180mg, 1.04mmol) and C-(5-(2,4-difluoro-phenoxy)-furan-2-yl)-methylaniline described in Preparation Example 77 (258mg, 1.15mmol) according to an analogous method to Example H-1.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.44 (2H, d, J=5.6Hz), 5.59 (1H, d, J=3.6Hz),  
 25 6.31 (1H, d, J=3.6Hz), 7.06-7.13 (1H, m), 7.27-7.34 (1H, m), 7.44-7.51 (1H, m),

7.59(1H, dd, J=4.0, 8.8Hz), 8.06 (1H, d, J=9.2 Hz), 8.16 (1H, dd, J=1.2, 8.8Hz), 8.45 (1H, d, J=9.2Hz), 8.51 (1H, s), 8.97 (1H, dd, J=1.2, 4.0Hz), 9.19 (1H, t, J=5.6Hz).

Example E-6. Quinoline-6-carboxylic acid

5 (5-(2,5-difluoro-phenoxy)-furan-2-ylmethyl)-amide

The title compound (194mg, 0.51mmol, 32%) was obtained from quinoline-6-carboxylic acid (275mg, 1.59mmol) and C-(5-(2,5-difluoro-phenoxy)-furan-2-yl)-methylamine described in Preparation Example 79 (357mg, 1.59mmol) according to an analogous method to Example H-1.

10 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.46 (2H, d, J=5.2Hz), 5.78 (1H, d, J=3.2Hz), 6.35 (1H, d, J=3.2Hz), 7.04-7.18 (2H, m), 7.42-7.50 (1H, m), 7.60 (1H, dd, J=4.0, 8.0Hz), 8.06 (1H, d, J=8.8Hz), 8.17 (1H, dd, J=1.2, 8.0Hz), 8.45 (1H, d, J=8.8Hz), 8.52(1H, s), 8.97 (1H, dd, J=1.2, 4.0Hz), 9.19 (1H, t, J=5.2Hz).

15 Example E-7. Quinoline-6-carboxylic acid

(5-(3-fluoro-benzyl)-furan-2-ylmethyl)-amide

Title compound (301mg, 0.835mmol, 80%) was obtained from quinoline-6-carboxylic acid (188mg, 1.16mmol) and C-(5-(3-fluoro-benzyl)-furan-2-yl)-methylamine described in Preparation Example 84 (279mg, 1.36mmol) according to an analogous method to Example H-1.

20 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.96 (2H, s), 4.46 (2H, d, J=5.6Hz), 6.07 (1H, d, J=3.2Hz), 6.21 (1H, d, J=3.2Hz), 7.00-7.09 (3H, m), 7.28-7.35 (1H, m), 7.59 (1H, dd, J=4.4, 8.4Hz), 8.05 (1H, d, J=8.8Hz), 8.16 (1H, d, J=2.0, 8.4Hz), 8.44 (1H, dd, J=1.2, 8.8Hz), 8.50 (1H, d, J=1.2Hz), 8.97 (1H, dd, J=2.0, 4.4Hz), 9.17 (1H, t, J=5.6Hz).

25



Example E-8. Quinoline-6-carboxylic acid 4-benzyloxybenzylamide

To a solution of 6-quinoline carboxylic acid (100mg, 0.577mmol) in tetrahydrofuran (50mL) was added N,N'-dicyclohexylcarbodiimide (1.90g, 11.7mmol), and the solution was stirred at room temperature for 1 hour. Then, to this solution was added a solution of 4-benzyloxybenzylamine described in Preparation Example 1 (2.49g, 11.7mmol) in tetrahydrofuran (5mL), and the solution was stirred overnight at room temperature. The solvent was evaporated, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate), and the title compound (4.31g, quantitatively) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.65 (2H, d, J=5.6Hz), 5.08(2H, s), 6.48(1H, brs), 6.97-7.00(2H, m), 7.31-7.35(3H, m), 7.37-7.45(4H, m), 7.47(1H, dd, J=4.4, 8.4Hz), 8.05(1H, dd, J=2.0, 8.8Hz), 8.15(1H, d, J=8.8Hz), 8.22-8.25(1H, m), 8.32(1H, d, J=2.0Hz), 8.98-9.00(1H, m).

Example E-9. Quinoline-6-carboxylic acid 3-benzyloxybenzylamide

The title compound (102mg, 48%) was obtained as a colorless oil from 6-quinolinecarboxylic acid (100mg, 0.58mmol) and 3-benzyloxybenzylamine described in Preparation Example 2 (126mg, 0.58mmol) according to an analogous method to Example E-8.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.69 (2H, d, J=5.6Hz), 5.07(2H, s), 6.57(1H, brs), 6.92-6.95(1H, m), 6.98-7.02(2H, m), 7.27-7.32(2H, m), 7.34-7.38(2H, m), 7.41-7.43(2H, m), 7.47(1H, dd, J=4.4, 8.4Hz), 8.05(1H, dd, J=2.0, 8.8Hz), 8.16(1H, d, J=8.8Hz), 8.22-8.25(1H, m), 8.32(1H, d, J=2.0Hz), 8.98-9.00(1H, m).

Example E-10. Quinoline-6-carboxylic acid 4-phenoxybenzylamide

The title compound (63mg, 31%) was obtained as a colorless solid from 6-quinolinecarboxylic acid (100mg, 0.58mmol) and 4-phenoxybenzylamine described in Preparation Example 3 (115mg, 0.58mmol) according to an analogous method to

5 Example E-8.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.69 (2H, d, J=5.6Hz), 6.59(1H, brs), 7.00-7.03(4H, m), 7.10-7.14(1H, m), 7.32-7.38(4H, m), 7.46-7.49(1H, m), 8.05-8.08(1H, m), 8.15-8.17(1H, m), 8.23-8.25(1H, m), 8.34(1H, s), 8.98-9.00(1H, m).

10 Example E-11. Quinoline-6-carboxylic acid 3-phenoxybenzylamide

The title compound (140mg, 69%) was obtained as a colorless oil from 6-quinolinecarboxylic acid (100mg, 0.58mmol) and 4-phenoxybenzylamine described in Preparation Example 3 (115mg, 0.58mmol) according to an analogous method to Example E-8.

15 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.69 (2H, d, J=6.0Hz), 6.63(1H, brs), 6.92-6.95(1H, m), 7.01-7.05(3H, m), 7.09-7.14(2H, m), 7.30-7.36(3H, m), 7.47(1H, dd, J=4.0, 8.0Hz), 8.05(1H, dd, J=2.0, 8.8Hz), 8.15(1H, d, J=8.8Hz), 8.21-8.24(1H, m), 8.32(1H, d, J=2.0Hz), 8.98-8.99(1H, m).

20 Example E-12. Quinoline-6-carboxylic acid 4-(pyridin-2-ylmethoxy)-benzylamide

To a solution of quinoline-6-carboxylic acid 4-hydroxybenzylamide described in Preparation Example E+-1 (20mg, 0.0719mmol) and 2-chloromethyl-pyridine hydrochloride (12mg, 0.0719mmol) in N,N-dimethyl formamide (1.0mL) was added potassium carbonate (298mg, 2.16mmol), and the solution was stirred at room

25 temperature for 12 hours. Ethyl acetate and water were added to the reaction

solution, which was then partitioned, and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), and trifluoroacetic acid salt of the title compound was obtained.

MS m/e (ESI) 369.2 (MH<sup>+</sup>)

<sup>1</sup>H-NMR Spectrum (CD<sub>3</sub>OD) δ(ppm) : 4.60(2H, s), 5.37(2H, s), 7.35(2H, d, J=8.8Hz), 7.40(2H, d, J=8.8Hz), 7.74-7.77(1H, m), 7.82-7.85(1H, m), 7.92-7.94(1H, m), 8.18-8.20(1H, m), 8.29-8.35(2H, m), 8.59-8.60(1H, m), 8.70-8.71(1H, m), 8.79-8.81(1H, m), 9.08-9.09(1H, m).

#### Example E-13. Quinoline-6-carboxylic acid (biphenyl-3-ylmethyl)-amide

The title compound (20mg, 21%) was obtained as a colorless oil from 6-quinolinecarboxylic acid (50mg, 0.289mmol) and C-biphenyl-3-yl-methylamine described in Preparation Example 5 (48mg, 0.263mmol) according to an analogous method to Example E-8.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.79(2H, d, J=5.6Hz), 6.62(1H, brs), 7.34-7.49(6H, m), 7.55-7.62(4H, m), 8.08(1H, dd, J=2.0, 8.8Hz), 8.16(1H, d, J=8.8Hz), 8.23-8.25(1H, m), 8.35(1H, d, J=2.0Hz), 8.98-9.00(1H, m).

#### Example E-14. Quinoline-6-carboxylic acid 4-(3-methyl-benzyloxy)-benzylamide

The title compound (2.5mg, 18%) was obtained as a white solid from quinoline-6-carboxylic acid 4-hydroxybenzylamide obtained in Preparation Example E+-1 (10mg, 0.0359mmol) and 3-methylbenzyl chloride (5mg, 0.0359mmol) according to an analogous method to Example E-12.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 2.38(3H, s), 4.66(2H, d, J=5.2Hz), 5.04(2H, s), 6.50(1H, brs), 6.97-7.00(2H, m), 7.14-7.16(1H, m), 7.22-7.35(5H, m), 7.46-7.50(1H, m), 8.06(1H, dd, J=2.0, 8.8Hz), 8.17(1H, d, J=8.8Hz), 8.23-8.26(1H, m), 8.33(1H, d, J=2.0Hz), 8.99-9.00(1H, m).

5

Example E-15. Quinoline-6-carboxylic acid 3-(4-fluoro-phenoxy)-benzylamide

The title compound (28mg, 25%) was obtained as a colorless oil from 6-quinolinecarboxylic acid (52mg, 0.30mmol) and 3-(4-fluorophenoxy)-benzylamine described in Preparation Example 8 (65mg, 0.30mmol) according to an analogous method to Example L-4.

10

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.70(2H, d, J=5.6Hz), 6.64(1H, brs), 6.88-6.91(1H, m), 6.99-7.06(5H, m), 7.12-7.14(1H, m), 7.31-7.35(1H, m), 7.47-7.50(1H, m), 8.04-8.07(1H, m), 8.15-8.17(1H, m), 8.23-8.25(1H, m), 8.33(1H, m), 9.00-9.01(1H, m).

15

Example E-16. Quinoline-6-carboxylic acid 3-(4-methoxy-phenoxy)-benzylamide

The title compound (28mg, 25%) was obtained as a colorless oil from 6-quinolinecarboxylic acid (50mg, 0.29mmol) and 3-(4-methoxyphenoxy) benzylamine described in Preparation Example 9 (66mg, 0.30mmol) according to an analogous method to Example L-4.

20

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.79(3H, s), 4.66(2H, d, J=5.6Hz), 6.55(1H, brs), 6.84-6.89(3H, m), 6.95-6.99(3H, m), 7.05-7.07(1H, m), 7.46(2H, dd, J=4.4, 8.4Hz), 8.03(1H, dd, J=2.4, 8.8Hz), 8.14(1H, d, J=8.8Hz), 8.21-8.23(1H, m), 8.29(1H, d, J=2.0Hz), 8.98(1H, dd, J=2.0, 4.4Hz).

25

Example E-17. Quinoline-6-carboxylic acid 3-(3-trifluoromethyl-phenoxy)-benzylamide

The title compound (39mg, 32%) was obtained as a colorless oil from 6-quinolinecarboxylic acid (50mg, 0.29mmol) and 3-(3-trifluoromethyl-phenoxy)-benzylamine described in Preparation Example 10 (77mg, 0.29mmol) according to an analogous method to Example L-4.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.70(2H, d, J=5.6Hz), 6.68(1H, brs), 6.93-6.95(1H, m), 7.05-7.06(1H, m), 7.14-7.25(3H, m), 7.31-7.47(4H, m), 8.04(1H, dd, J=1.6, 8.8Hz), 8.13(1H, d, J=8.8Hz), 8.20-8.2(1H, m), 8.31(1H, d, J=2.0Hz), 8.97(1H, dd, J=1.6, 4.4Hz).

Example E-18. Quinoline-6-carboxylic acid 3-(3-fluoro-phenoxy)-benzylamide

The title compound (70mg, 65%) was obtained as a colorless oil from 6-quinolinecarboxylic acid (50mg, 0.29mmol) and 3-(3-fluoro-phenoxy)-benzylamine described in Preparation Example 11 (63mg, 0.29mmol) according to an analogous method to Example L-4.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.71(2H, d, J=6.0Hz), 6.63(1H, brs), 6.69-6.73(1H, m), 6.78-6.83(2H, m), 6.96-6.98(1H, m), 7.07(1H, s), 7.18(1H, d, J=7.6Hz), 7.24-7.30(1H, m), 7.34-7.38(1H, m), 7.48(1H, dd, J=4.4, 8.4Hz), 8.06(1H, dd, J=2.0, 8.8Hz), 8.16(1H, d, J=8.8Hz), 8.23-8.25(1H, m), 8.33(1H, d, J=1.6Hz), 8.99(1H, dd, J=1.6, 4.4Hz).

Example E-19. 2-(3-Phenoxy-phenyl)-N-quinolin-6-yl-acetamide

The title compound (116mg, 94%) was obtained as a colorless oil from 6-aminoquinoline (50mg, 3.47mmol) and 3-phenoxyphenylacetic acid (79mg,

3.47mmol) according to an analogous method to Example L-4.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.77(2H, s), 6.98-7.06(4H, m), 7.10-7.16(2H, m), 7.34-7.44(6H, m), 8.01(1H, d, J=8.8Hz), 8.09-8.16(1H, m), 8.32(1H, d, J=2.0Hz), 8.83(1H, dd, J=1.6, 4.0Hz).

5

Example E-20. Quinoline-6-carboxylic acid 4-(furan-2-ylmethoxy)-benzylamide

The title compound (0.7mg, 4%) was obtained as a colorless oil from 6-quinolinecarboxylic acid (10mg, 0.0577mmol) and 4-(furan-2-ylmethoxy)-benzylamine described in Preparation Example 12 (12mg, 0.0577mmol) according to an analogous method to Example L-4.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.66(2H, d, J=5.6Hz), 5.02(2H, s), 6.48(1H, brs), 6.99-7.02(2H, m), 7.24-7.27(2H, m), 7.35-7.36(2H, m), 7.47-7.50(2H, m), 8.05-8.08(1H, m), 8.16-8.18(1H, m), 8.24-8.27(1H, m), 8.33(1H, s), 9.00(1H, s).

15 Example E-21. Quinoline-6-carboxylic acid 4-(thiophen-2-ylmethoxy)-benzylamide

The title compound (11mg, 53%) was obtained as a colorless solid from 6-quinolinecarboxylic acid (10mg, 0.0577mmol) and 4-(thiophen-2-ylmethoxy)-benzylamine described in Preparation Example 13 (13mg, 0.0577mmol) according to an analogous method to Example L-4.

20 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.65(2H, d, J=5.6Hz), 5.23(2H, s), 6.60(1H, brs), 6.97-7.03(3H, m), 7.11-7.12(1H, m), 7.32-7.34(3H, m), 7.45-7.48(1H, m), 8.05-8.07(1H, m), 8.13-8.16(1H, m), 8.21-8.23(1H, m), 8.32(1H, s), 8.97-8.99(1H, m).

Example E-22. Quinoline-6-carboxylic acid 4-(thiophen-3-ylmethoxy)-benzylamide

25 The title compound (78mg, 72%) was obtained as a white solid from

6-quinolinecarboxylic acid (50mg, 0.289mmol) and 4-(thiophen-3-ylmethyl)-benzylamine described in Preparation Example 14 (63mg, 0.289mmol) according to an analogous method to Example L-4.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.65(2H, d, J=5.6Hz), 5.08(2H, s), 6.51(1H, brs),  
 5 6.96-6.99(2H, m), 7.14-7.16(1H, m), 7.32-7.36(4H, m), 7.46-7.49(1H, m),  
 8.04-8.07(1H, m), 8.15(1H, d, J=8.8Hz), 8.22-8.24(1H, m), 8.32(1H, d, J=2.0Hz),  
 8.99(1H, dd, J=2.0, 4.4Hz).

Example E-23. Quinoline-6-carboxylic acid 4-((S)-1-phenyl-ethoxy)-benzylamide

10 The title compound (219mg, 80%) was obtained as a colorless oil from 6-quinolinecarboxylic acid (123mg, 0.712mmol) and 4-((S)-1-phenyl-ethoxy)-benzylamine described in Preparation Example 15 (172mg, 0.712mmol) according to an analogous method to Example L-4.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 1.64(3H, d, J=6.4Hz), 4.58(2H, d, J=5.2Hz),  
 15 5.31(1H, q, J=6.4Hz), 6.47(1H, brs), 6.86(2H, d, J=8.8Hz), 7.22(2H, d, J=8.8Hz),  
 7.26-7.27(2H, m), 7.32-7.39(3H, m), 7.46(1H, dd, J=4.0, 8.0Hz), 8.03(1H, dd, J=2.0,  
 8.8Hz), 8.13(1H, d, J=8.8Hz), 8.22(1H, d, J=8.0Hz), 8.30(1H, d, J=2.0Hz),  
 8.97-8.98(1H, m).

20 Example E-24. Quinoline-6-carboxylic acid 3-benzylamino-benzylamide

3-Benzylamino-benzonitrile described in Preparation Example 87 (57mg, 0.27mmol) was dissolved at 0°C in tetrahydrofuran (0.5mL), and lithium aluminum hydride (52mg, 1.35mmol) was added. After stirring overnight at room temperature, water (52μl), an aqueous solution of 5N sodium hydroxide (52μl) and water (156μl)  
 25 were sequentially added at 0°C. The reaction solution was filtered through Celite

pad, then, the solvent was evaporated *in vacuo*, and (3-aminomethyl-phenyl)-benzyl-amine (62mg, 0.29mmol) was obtained as an oil.

(3-Aminomethyl-phenyl)-benzylamine (62mg,0.29mmol),  
quinoline-6-carboxylic acid (52mg,0.30mmol),

5 benzotriazol-1-yl-tris(dimethylamino)phosphonium hexafluorophosphate (182mg,0.41mmol) and triethylamine (114 $\mu$ l,0.81mmol) were dissolved in N,N-dimethylformamide (0.5mL), and the solution was stirred at room temperature for 2 hours. Water was added to the reaction solution, which was then extracted with ethyl acetate, the organic layer was washed with water and brine, and dried over  
10 anhydrous magnesium sulfate. The solvent was evaporated *in vacuo*, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 1 : 1), and the title compound (73mg,0.198mmol,73%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>)  $\delta$ (ppm) : 4.22 (2H, d, J=6.0Hz), 4.39 (2H, d, J=6.0Hz), 6.25 (1H, t, J=6.0Hz), 6.40 (1H, dd, J=1.6, 7.7Hz), 6.50 (1H, d, J=7.7Hz), 6.60 (1H, s),  
15 6.97 (1H, t, J=7.8Hz), 7.14 (1H, t, J=7.3Hz), 7.23 (2H, t, J=7.8Hz), 7.30 (2H, d, J=7.1Hz), 7.60 (1H, dd, J=4.4, 8.4Hz), 8.07 (1H, d, J=8.8Hz), 8.19 (1H, dd, J=2.0, 8.8Hz), 8.45 (1H, d, J=7.6Hz), 8.52 (1H, d, J=1.6Hz), 8.97 (1H, dd, J=1.6, 4.4Hz), 9.14 (1H, t, J=6.0Hz).

20 Example E-25. Quinoline-6-carboxylic acid 4-phenylamino-benzylamide

(4-Aminomethyl-phenyl)-phenylamine (98mg, 0.494mmol) was obtained as an oil from 4-phenylamino-benzonitrile described in Preparation Example 88 (110mg, 0.566mmol) according to an analogous method to Example E-24.

Then, the title compound (52mg, 0.147mmol, 26%) was obtained from the  
25 resulting (4-aminomethyl-phenyl)-phenylamine and quinoline-6-carboxylic acid



(108mg,0.623mmol).

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.66 (2H, d, J=5.5Hz), 5.76 (1H, s), 6.50 (1H, brs), 6.96 (1H, t, J=7.4Hz), 7.07-7.10 (4H, m), 7.29-7.32 (4H, m), 7.49 (1H, dd, J=4.2, 8.4Hz), 8.08 (1H, dd, J=2.0, 8.6Hz), 8.17 (1H, d, J=8.8Hz), 8.25 (1H, d, J=7.5Hz),  
5 8.34 (1H, d, J=1.8Hz), 9.00 (1H, dd, J=1.7, 4.2Hz).

Example E-26. Quinoline-6-carboxylic acid 4-(benzyl-methyl-amino)-benzylamide

Quinoline-6-carboxylic acid 4-benzylamino-benzylamide described in Preparation Example E-3 (30mg, 82μmol), formalin (9μl, 115μmol), triacetoxysodium borohydride (25mg,115μmol) and acetic acid (several drops) were suspended  
10 in tetrahydrofuran (1mL), and the solution was stirred at room temperature for 4 hours. An aqueous solution of saturated sodium bicarbonate was added to the reaction suspension at 0°C, which was then extracted with ethyl acetate, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The  
15 solvent was evaporated *in vacuo*, purification was carried out using thin layer NH silica gel chromatography (hexane : ethyl acetate = 1 : 1), and the title compound (11mg,28μmol,35%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.05 (3H, s), 4.55 (2H, s), 4.59 (2H, d, J=5.1Hz), 6.38 (1H, brs), 6.74 (2H, d, J=8.8Hz), 7.21-7.26 (4H, m), 7.32 (2H, t, J=7.2Hz), 7.4  
20 (1H, d, J=7.0Hz), 7.47 (1H, dd, J=4.3, 8.2Hz), 8.05 (1H, dd, J=2.0, 8.8Hz), 8.14 (1H, d, J=8.8Hz), 8.23 (1H, d, J=7.1Hz), 8.30 (1H, d, J=1.7Hz), 8.98 (1H, dd, J=1.7, 4.2Hz).

Example E-27. Quinoline-6-carboxylic acid 3-phenylsulfanyl-benzylamide

25 The title compound (50mg, 0.14mmol, 75%) was obtained as a white solid

from 3-phenylsulfanyl-benzylamine described in Preparation Example 95 (38mg, 0.18mmol) and quinoline-6-carboxylic acid (31mg, 0.18mmol) according to an analogous method to Example A-26.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.67 (2H, d, J=5.7Hz), 6.52 (1H, brs), 7.23-7.26 (3H, m), 7.31 (4H, t, J=7.4Hz), 7.38 (2H, d, J=6.8Hz), 7.49 (1H, dd, J=4.2, 8.4Hz), 8.03 (1H, dd, J=1.8, 8.8Hz), 8.16 (1H, d, J=8.8Hz), 8.24 (1H, d, J=7.5Hz), 8.31 (1H, d, J=1.8Hz), 9.00 (1H, dd, J=1.7, 4.2Hz).

Example E-28. Quinoline-6-carboxylic acid 4-benzylsulfanyl-benzylamide

The title compound (54mg, 0.14mmol, 38%) was obtained as a white solid from 4-benzylsulfanyl-benzylamine described in Preparation Example 101 (84mg, 0.37mmol) and quinoline-6-carboxylic acid (70mg, 0.40mmol) according to an analogous method to Example A-26.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.20 (2H, s), 4.47 (2H, d, J=5.9Hz), 7.18-7.34 (9H, m), 7.59 (1H, dd, J=4.2, 8.4Hz), 8.07 (1H, d, J=9.0Hz), 8.18 (1H, dd, J=2.0, 8.8Hz), 8.45 (1H, d, J=7.9Hz), 8.52 (1H, d, J=2.0Hz), 8.97 (1H, dd, J=1.8, 4.2Hz), 9.24 (1H, t, J=5.9Hz).

Example E-29. Quinoline-6-carboxylic acid 3-(3-methylbutoxy)-benzylamide

To a mixture solution of quinoline-6-carboxylic acid 3-hydroxybenzylamide described in Preparation Example E+-4 (13mg, 0.048mmol) and N,N-dimethylformamide (0.5mL) were added potassium carbonate (13mg, 0.096mmol) and 1-iodine-3-methylbutane (0.013mL, 0.096mmol), and the solution was stirred overnight at room temperature. Water and dichloromethane was added to the reaction solution for extraction, which was then washed with brine,

then, the solvent was evaporated *in vacuo*. The residue was purified by silica gel column chromatography (ethyl acetate), and the title compound (12mg, 0.033mmol, 69%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.96 (6H, d, J=6.6Hz), 1.65-1.70 (2H, m),  
5 1.80-1.87 (1H, m), 3.99 (2H, t, J=6.7Hz), 4.68 (2H, d, J=5.7Hz), 6.56 (1H, brs),  
6.84-6.87 (1H, m), 6.93-6.98 (2H, m), 7.26-7.31 (1H, m), 7.48 (1H, dd, J=4.2, 8.2Hz),  
8.07 (1H, dd, J=2.0, 8.8Hz), 8.16 (1H, d, J=8.8Hz), 8.23-8.25 (1H, m), 8.33 (1H, d,  
J=2.0Hz), 8.99 (1H, dd, J=1.8, 4.2Hz).

10 Example E-30. Quinoline-6-carboxylic acid (Z)-4-styryl-benzylamide

To a mixture of quinoline-6-carboxylic acid 4-phenylethynyl-benzylamide described in Preparation Example E-5 (48mg, 0.13mmol), quinoline (26mg, 0.20mmol) and tetrahydrofuran (2mL) was added Lindlar catalyst (5.0mg), and the solution was stirred under hydrogen atmosphere at room temperature for 1 hour.

15 The interior of the reaction system was changed to nitrogen atmosphere, then, filtration was carried out through Celite pad. The filtrate was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 1 : 4), and the title compound (45mg, 0.12mmol, 92%) was obtained as a colorless oil.

20 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.57 (2H, d, J=6.0Hz), 6.58-6.60 (2H, m),  
7.16-7.27 (9H, m), 7.59 (1H, dd, J=4.2, 8.2Hz), 8.06 (1H, d, J=8.8Hz), 8.19 (1H, dd,  
J=2.0, 8.8Hz), 8.45-8.47 (1H, m), 8.54 (1H, d, J=1.7Hz), 8.97 (1H, dd, J=1.7, 4.2Hz),  
9.24 (1H, t, J=5.7Hz).

25 Example E-31. Quinoline-6-carboxylic acid 4-phenylaminomethyl-benzylamide

The title compound (13.8mg) was obtained according to an analogous method to Example E-26 using quinoline-6-carboxylic acid 4-formyl-benzylamide described in Preparation Example E+-6 (50mg, 0.172mmol) instead of formalin, and phenylamine (31μl, 0.34mmol) instead of quinoline-6-carboxylic acid 4-benzyl amino-benzylamide. Trifluoroacetic acid salt of the title compound was obtained by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used).

MS m/e (ESI) 368.5(MH<sup>+</sup>)

10 Example E-32. Quinoline-6-carboxylic acid  
4-((methyl-phenyl-amino)-methyl)-benzylamide

The title compound (4.25mg) was obtained from quinoline-6-carboxylic acid 4-phenylaminomethyl-benzylamide described in Example E-31 (30mg, 82μmol) according to an analogous method to Example E-26. Trifluoroacetic acid salt of the title compound was obtained by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used).

MS m/e (ESI) 382.3 (MH<sup>+</sup>)

20 Example E-33. Quinoline-6-carboxylic acid 3-(4-nitro phenoxy)-benzylamide

To a mixture solution of quinoline-6-carboxylic acid 3-hydroxybenzylamide described in Preparation Example E+-4 (3.0mg, 0.011mmol), copper(II) acetate (2.9mg, 0.016mmol), molecular sieves 4A (50mg) and dichloromethane (2mL) were added triethylamine (0.0077mL, 0.055mmol) and 4-nitrophenylboronic acid (1.8mg, 0.011mmol), and the solution was stirred in the presence of air at room temperature

for 10 days. The reaction solution was filtered, then, water, ethyl acetate and an aqueous solution of 29% ammonia were added for extraction, the solution was washed with brine, then, the solvent was evaporated *in vacuo*. The residue was purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), and trifluoroacetic acid salt of the title compound (1.14mg, 0.0022mmol, 20%) was obtained.

MS m/e (ESI) 400.2 (MH<sup>+</sup>)

Example E-34. Quinoline-6-carboxylic acid 3-(4-methanesulfonylphenoxy)-benzylamide

Trifluoroacetic acid salt of the title compound (0.21mg, 0.00038mmol, 3.5%) was obtained from quinoline-6-carboxylic acid 3-hydroxybenzylamide described in Preparation Example E+-4 (3.0mg, 0.011mmol) and 4-methanesulfonylphenylboronic acid (2.2mg, 0.011mmol) according to an analogous method to Example E-33.

MS m/e (ESI) 433.2 (MH<sup>+</sup>)

Example E-35. 4-(3-(((Quinoline-6-carbonyl)amino)methyl)phenoxy)benzoic acid methyl ester

Trifluoroacetic acid salt of the title compound (0.14mg, 0.00027mmol, 2.4%) was obtained from quinoline-6-carboxylic acid 3-hydroxybenzylamide described in Preparation Example E+-4 (3.0mg, 0.011mmol) and 4-methoxy-carbonyl phenylboronic acid (1.8mg, 0.011mmol) according to an analogous method to Example E-33.

MS m/e (ESI) 413.3 (MH<sup>+</sup>)

Example E-36. Quinoline-6-carboxylic acid 3-(3-cyanophenoxy)-benzylamide

Trifluoroacetic acid salt of the title compound (0.25mg, 0.00051mmol, 4.6%) was obtained from quinoline-6-carboxylic acid 3-hydroxybenzylamide described in Preparation Example E+-4 (5.0mg, 0.011mmol) and 3-cyanophenylboronic acid (2.6mg, 0.018mmol) according to an analogous method to Example E-33.  
MS m/e (ESI) 380.1 (MH<sup>+</sup>)

Example E-37. Quinoline-6-carboxylic acid 3-(3-acetylphenoxy)-benzylamide

Trifluoroacetic acid salt of the title compound (0.17mg, 0.00033mmol, 3.0%) was obtained from quinoline-6-carboxylic acid 3-hydroxybenzylamide described in Preparation Example E+-4 (5.0mg, 0.011mmol) and 3-acetylphenylboronic acid (3.0mg, 0.018mmol) according to an analogous method to Example E-33.  
MS m/e (ESI) 397.0 (MH<sup>+</sup>)

Example E-38. Quinoline-6-carboxylic acid 3-(3-trifluoromethoxyphenoxy)-benzylamide

Trifluoroacetic acid salt of the title compound (0.15mg, 0.00027mmol, 2.5%) was obtained from quinoline-6-carboxylic acid 3-hydroxybenzylamide described in Preparation Example E+-4 (5.0mg, 0.011mmol) and 3-trifluoromethoxyphenylboronic acid (3.7mg, 0.018mmol) according to an analogous method to Example E-330.  
MS m/e (ESI) 439.0 (MH<sup>+</sup>)

Example E-39. Quinoline-6-carboxylic acid (3'-fluorobiphenyl-3-ylmethyl)-amide

To a mixture of quinoline-6-carboxylic acid 3-bromobenzylamide described in Preparation Example E+-8 (4.0mg, 0.012mmol), toluene (1mL) and methanol

(0.25mL) were added an aqueous solution of 2M sodium carbonate (0.5mL), tetrakis(triphenylphosphine)palladium(0) (1.4mg, 0.0012mmol) and 3-fluorophenylboronic acid (1.7mg, 0.012mmol), and the solution was stirred at 70°C for 4 hours. After cooling, water, ethyl acetate and acetic acid were added for extraction, the solution was washed with brine, then, the solvent was evaporated *in vacuo*. The residue was purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), and trifluoroacetic acid salt of the title compound (0.47mg, 0.0010mmol, 8.3%) was obtained.

MS m/e (ESI) 357.2 (MH<sup>+</sup>)

Example E-40. Quinoline-6-carboxylic acid 3-benzyl-benzylamide

To a mixture of quinoline-6-carboxylic acid 3-bromobenzylamide described in Preparation Example E-8 (8.0mg, 0.023mmol), dichloro(1,1'-bis(diphenylphosphino)ferrocene) nickel(II) (3.2mg, 0.0047mmol) and tetrahydrofuran (1mL) was added benzylmagnesium chloride (1.1M tetrahydrofuran solution, 0.088mL, 0.094mmol) at room temperature, and the solution was stirred for 30 minutes at 50°C. After cooling, water and ethyl acetate were added for extraction, the solution was washed with brine, the solvent was then evaporated *in vacuo*. The residue was purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), and trifluoroacetic acid salt of the title compound (2.1mg, 0.0045mmol, 19%) was obtained.

MS m/e (ESI) 353.2 (MH<sup>+</sup>)

Example E-41. Quinoline-6-carboxylic acid 4-benzyl-benzylamide

Trifluoroacetic acid salt of the title compound (1.5mg, 0.0032mmol, 14%) was obtained from quinoline-6-carboxylic acid 4-bromobenzylamide described in Preparation Example E+9 (8.0mg, 0.023mmol) according to an analogous method to

5 Example E-40.

MS m/e (ESI) 353.3 (MH<sup>+</sup>)

Example E-42. Quinoline-6-carboxylic acid 4-phenethyl-benzylamide

Trifluoroacetic acid salt of the title compound (0.39mg, 0.00081mmol, 3.5%) was obtained from quinoline-6-carboxylic acid 4-bromobenzylamide described in Preparation Example E+9 (8.0mg, 0.023mmol) and phenethylmagnesium chloride (1.0M tetrahydrofuran solution, 0.094mL, 0.094mmol) according to an analogous method to Example E-40.

MS m/e (ESI) 367.3 (MH<sup>+</sup>)

15

Example E-43. Quinoline-6-carboxylic acid 3-cyclopropylmethoxy-benzylamide

To a mixture of quinoline-6-carboxylic acid 3-hydroxybenzylamide described in Preparation Example E+4 (87mg, 0.31mmol) and tetrahydrofuran (2mL) was added an aqueous solution of 1N sodium hydroxide (0.31mL, 0.31mmol), and the solvent was evaporated *in vacuo*. To a mixture of a portion (5.0mg) of the residue (93mg) and N,N-dimethylformamide (1mL) were added cyclopropylmethyl bromide (2.7mg, 0.020mmol) and catalytic amount of sodium iodide at room temperature, followed by stirring for 3 hours at the same temperature. The reaction solution was purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), and trifluoroacetic acid

25



salt of the title compound (1.50mg, 0.0034mmol, 20%) was obtained.

MS m/e (ESI) 333.0 (MH<sup>+</sup>)

Example E-44. N-(4-Benzyloxy-benzyl)-N'-methoxy-quinoline-6-carboxamidine

5           To a mixture of quinoline-6-carbothioic acid 4-benzyloxy-benzylamide described in Preparation Example E+-10 (57mg, 0.15mmol) and acetonitrile (3mL) was added 2-(bromomethyl)naphthalene (200mg, 0.94mmol), which was then refluxed for 2 hours. After cooling, the solvent was evaporated *in vacuo*, and the residue was washed with diethyl ether three times. A mixture of a portion (29mg) of  
10 the resulting crude product (57mg), methoxylamine hydrochloride (2.9mg, 0.035mmol), an aqueous solution of 1N sodium hydroxide (0.035mL, 0.035mmol) and N-methylpyrrolidinone (1mL) was stirred at room temperature for 25 minutes. The reaction solution was directly purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic  
15 acid) was used), and ditrifluoroacetic acid salt of the title compound (1.9mg, 0.0030mmol, 6.4%) was obtained.

MS m/e (ESI) 398.5 (MH<sup>+</sup>)

Example E-45. N-(4-Benzyloxy-benzyl)-N'-cyano-quinoline-6-carboxamidine

20           To a mixture of quinoline-6-carbothioic acid 4-benzyloxy-benzylamide described in Preparation Example E+-10 (57mg, 0.15mmol) and toluene (2mL) was added benzyl bromide (0.089mL, 0.74mmol), which was then stirred under reflux for 90 minutes. After cooling, the solvent was evaporated *in vacuo*, and the residue was washed with diethyl ether twice. A mixture of a portion (16mg) of the resulting  
25 crude product (72mg), cyanamide (20mg, 0.48mmol) and N-methylpyrrolidinone

(1mL) was stirred for 2.5 hours at 120°C. After cooling, the reaction solution was filtered through a membrane filter, the filtrate was directly purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), and trifluoroacetic acid salt of the title compound  
5 (0.76mg, 0.0015mmol, 4.5%) was obtained.

MS m/e (ESI) 393.5 (MH<sup>+</sup>)

Example E-46. Quinoline-6-carboxylic acid 4-(3-chloro-benzyloxy)-benzylamide

Trifluoroacetic acid salt of the title compound was obtained from  
10 quinoline-6-carboxylic acid 4-hydroxybenzylamine described in Preparation Example E+-1 and 3-chlorobenzyl chloride according to an analogous method to Example E-12.

MS m/e (ESI) 403 (MH<sup>+</sup>)

15 Example E-47. Quinoline-6-carboxylic acid 4-(3-fluoro-benzyloxy)-benzylamide

Trifluoroacetic acid salt of the title compound was obtained from quinoline-6-carboxylic acid 4-hydroxybenzylamine described in Preparation Example E+-1 and 3-fluorobenzyl bromide according to an analogous method to Example E-12.

20 MS m/e (ESI) 387 (MH<sup>+</sup>)

Example E-48. Quinoline-6-carboxylic acid 4-(benzo[1,3]dioxol-5-ylmethoxy)-benzylamide

Trifluoroacetic acid salt of the title compound was obtained from  
25 quinoline-6-carboxylic acid 4-hydroxybenzylamine described in Preparation Example

E+-1 and methanesulfonic acid benzo[1,3]dioxol-5-ylmethyl ester according to an analogous method to Example E-12.

MS m/e (ESI) 413 (MH<sup>+</sup>)

5 Example E-49. 6-Quinolinecarboxylic acid 3-(3-methyl-2-butenyloxy)-benzylamide

6-Quinolinecarboxylic acid (100mg, 0.577mmol), 3-(3-methyl-2-butenyloxy)-benzylamine described in Preparation Example 134 (112mg, 0.635mmol), benzotriazol-1-yl-tris(dimethylamino)phosphonium hexafluorophosphate (306mg, 0.693mmol) and triethylamine (0.12mL, 0.87mmol) were dissolved in tetrahydrofuran (5mL), and the solution was stirred at room temperature for 3 hours. The solvent was evaporated *in vacuo*, the obtained residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 4 : 1), and the title compound (153mg, 80.1%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 1.73(3H, s), 1.78(3H, s), 4.51(2H, d, J=6.8Hz), 4.68(2H, d, J=5.6Hz), 5.48(1H, t, J=6.8Hz), 6.62(1H, t, J=5.6Hz), 6.87(1H, dd, J=2.4, 8.4Hz), 6.95(1H, d, J=2.4Hz), 6.96(1H, d, J=8.4Hz), 7.28(1H, t, J=8.4Hz), 7.47(1H, dd, J=4.0, 8.0Hz), 8.05(1H, dd, J=2.0, 8.8Hz), 8.13(1H, d, J=8.8Hz), 8.23(1H, dd, J=1.2, 8.0Hz), 8.32(1H, d, J=2.0Hz), 8.97(1H, dd, J=1.2, 4.0Hz).

20 Example E-50. 6-Quinolinecarboxylic acid 3-(2-methylpropenyl)-benzylamide

The title compound (150mg, 0.475mmol, 82.2%) was obtained as a white solid from 6-quinolinecarboxylic acid (100mg, 0.577mmol) and 3-(2-methyl-propenyl)-benzylamine described in Preparation Example 137 (93mg, 0.577mmol) according to an analogous method to Example E-49.

25 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 1.87(3H, s), 1.90(3H, s), 4.70(2H, d, J=5.6Hz),

6.27(1H, s), 6.62(1H, t, J=5.6Hz), 7.20-7.46(4H, m), 7.47(1H, dd, J=4.0, 8.0Hz), 8.05(1H, dd, J=2.0, 8.8Hz), 8.13(1H, d, J=8.8Hz), 8.23(1H, dd, J=1.2, 8.0Hz), 8.32(1H, d, J=2.0Hz), 8.97(1H, dd, J=1.2, 4.0Hz).

5 Example E-51. 6-Quinolinecarboxylic acid 3-cyclopentylidenemethylbenzylamide

The title compound (150mg, 0.457mol, 79.3%) was obtained as a white solid from 6-quinolinecarboxylic acid (100mg, 0.577mmol) and 3-cyclopentylidenemethyl-benzylamine described in Preparation Example 139 (108mg, 0.577mmol) according to an analogous method to Example E-49.

10 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 1.65-1.79(4H, m), 2.47-2.58(4H, m), 4.70(2H, d, J=5.6Hz), 6.36(1H,s), 6.54(1H, t, J=5.6Hz), 7.20-7.35(4H, m), 7.47(1H, dd, J=4.0, 8.0Hz), 8.05(1H, dd, J=2.0, 8.8Hz), 8.13(1H, d, J=8.8Hz), 8.23(1H, dd, J=1.2, 8.0Hz), 8.32(1H, d, J=2.0Hz), 8.97(1H, dd, J=1.2, 4.0Hz).

15 Example E-52. 6-Quinolinecarboxylic acid 3-isobutylbenzylamide

The title compound (75mg, 0.236mol, 66.2%) was obtained as a white solid from 6-quinolinecarboxylic acid (60mg, 0.356mmol) and 3-isobutylbenzylamine described in Preparation Example 146 (58mg, 0.356mmol) according to an analogous method to Example E-49.

20 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.90(6H, d, J=6.8Hz), 1.87(1H, dq, J=7.6Hz, 6.8Hz), 2.48(2H, d, J=7.6Hz), 4.69(2H, d, J=6.0Hz), 6.52(1H, t, J=6.0Hz), 7.10-7.30(4H, m), 7.47(1H, dd, J=4.0, 8.0Hz), 8.05(1H, dd, J=2.0, 8.8Hz), 8.13(1H, d, J=8.8Hz), 8.23(1H, dd, J=1.2, 8.0Hz), 8.32(1H, d, J=2.0Hz), 8.97(1H, dd, J=1.2, 4.0Hz).

Example E-53. Quinoline-6-carboxylic acid 4-benzyloxy-2-fluoro-benzylamide

To a mixture of lithium aluminum hydride (84mg, 2.2mmol) and tetrahydrofuran (2mL) was added 4-benzyloxy-2-fluoro-benzonitrile described in Preparation Example 118 (100mg, 0.44mmol) on an ice bath, and the solution was stirred at room temperature for 1 hour. Water (0.084mL), an aqueous solution of 5N sodium hydroxide (0.084mL) and water (0.25mL) were added sequentially on an ice bath, and the solution was stirred at room temperature for 90 minutes. The reaction solution was filtered through Celite pad, then, the solvent was evaporated *in vacuo*. The title compound (140mg, 0.35mmol, 90%) was obtained as a white solid from the obtained residue (91mg) and quinoline-6-carboxylic acid (68mg, 0.39mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.68 (2H, d, J=5.7Hz), 5.05 (2H, s), 6.62 (1H, br s), 6.71-6.78 (2H, m), 7.34-7.43 (6H, m), 7.47 (1H, dd, J=4.3, 8.3Hz), 8.04 (1H, dd, J=2.0, 8.8Hz), 8.15 (1H, d, J=8.8Hz), 8.23-8.25 (1H, m), 8.31 (1H, d, J=2.0Hz), 8.98 (1H, dd, J=1.8, 4.3Hz).

Example E-54. Quinoline-6-carboxylic acid 4-benzyloxy-3-chloro-benzylamide

Trifluoroacetic acid salt of the title compound (29mg, 0.057mmol, 10%) was obtained from quinoline-6-carboxylic acid 4-benzyloxybenzylamide described in Example E-8 (200mg, 0.54mmol) according to an analogous method to Example A-171.

MS m/e (ESI) 403.1 (MH<sup>+</sup>)

Example E-55. Quinoline-6-carboxylic acid (4-phenoxy-pyridine-2-ylmethyl)-amide

The title compound (9mg, 8%) was obtained as a colorless oil from

6-quinolinecarboxylic acid (52mg, 0.30mmol) and C-(4-phenoxy-pyridine-2-yl)-methylamine described in Preparation Example 7 (60mg, 0.30mmol) according to an analogous method to Example L-4.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.74(2H, d, J=4.4Hz), 6.79(1H, dd, J=2.4Hz, 5.6Hz), 6.88(1H, d, J=2.4Hz), 7.10-7.12(2H, m), 7.43-7.49(4H, m), 7.81(1H, brs), 8.14-8.19(2H, m), 8.26-8.28(1H, m), 8.40-8.44(2H, m), 8.99-9.0(1H, m).

Example E-56. Quinoline-6-carboxylic acid (6-phenoxy-pyridin-2-ylmethyl)-benzylamide

10 The title compound (32mg, 28%) was obtained as a colorless oil from 6-quinolinecarboxylic acid (56mg, 0.325mmol) and C-(6-phenoxy-pyridin-2-yl)-methylamine described in Preparation Example 16 (65mg, 0.325mmol) according to an analogous method to Example L-4.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.73(2H, d, J=4.4Hz), 6.89-6.91(1H, m), 7.04-7.06(1H, m), 7.21-7.23(2H, m), 7.30-7.34(1H, m), 7.44-7.52(3H, m), 7.64(1H, 15 brs), 7.65-7.68(1H, m), 7.73-7.77(1H, m), 8.11(1H, d, J=8.4Hz), 8.26(1H, d, J=8.0Hz), 8.32(1H, d, J=1.6Hz), 9.01-9.03(1H, m).

Example E-57. Quinoline-6-carboxylic acid (1-benzyl-1H-pyrrol-3-ylmethyl)-amide

20 7N Ammonia/methanol (80mL) and Raney nickel (2g) were added to 1-benzyl-1H-pyrrole-3-carbaldehyde described in Preparation Example 57 (800mg, 4.3mmol), and the solution was stirred at room temperature for 22 hours under hydrogen atmosphere at ordinary pressure. The catalyst was removed by filtrating through Celite pad, then, the solvent was evaporated *in vacuo* and 25 C-(1-benzyl-1H-pyrrolo-3-yl)methylamine was quantitatively obtained as a brown oil.

The title compound (110mg, 0.32mmol, 24.8%) was obtained as a white solid from the resulting C-(1-benzyl-1*H*-pyrrolo-3-yl)methylamine (240mg, 1.3mmol) and 6-quinolinecarboxylic acid (180mg, 1.04mmol) according to an analogous method to Example Q-6.

- 5 <sup>1</sup>H-NMR Spectrum (DMSO-*d*<sub>6</sub>) δ(ppm) : 4.32 (2H, d, 5.6Hz), 5.02 (2H, s), 6.02 (1H, s), 6.74 (1H, s), 6.76 (1H, s), 6.18-7.34 (5H, m), 7.58 (1H, dd, *J*=4.0, 8.4Hz), 8.03 (1H, d, *J*=8.8Hz), 8.16 (1H, dd, *J*=2.0, 8.8Hz), 8.43 (1H, dd, *J*=1.6, 8.4Hz), 8.49 (1H, d, *J*=2.0Hz), 8.97 (1H, t, *J*=5.6Hz), 8.95 (1H, dd, *J*=1.6, 4.0Hz).

10 Example E-58. Quinoline-6-carboxylic acid (1-benzo[1,3]dioxol-5-ylmethyl-1*H*-pyrrol-3-ylmethyl)-amide

The title compound (30mg, 0.077mmol, 18.1%) was obtained as a white solid from C-(1-benzo[1,3]dioxol-5-ylmethyl-1*H*-pyrrol-3-yl)-methylamine described in Preparation Example 61 (100mg, 0.43mmol) and 6-quinolinecarboxylic acid (75mg,  
15 0.43mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-*d*<sub>6</sub>) δ(ppm) : 4.28-4.34 (2H, m), 4.90 (2H, s), 5.95 (2H, s), 6.00 (1H, s), 6.70-6.86 (5H, m), 7.55-7.60 (1H, m), 8.01-8.05 (1H, m), 8.14-8.20 (1H, m), 8.40-8.46 (1H, m), 8.48-8.51 (1H, m), 8.39-8.98 (2H, m).

20 Example E-59. Quinoline-6-carboxylic acid (1-phenethyl-1*H*-pyrrol-3-ylmethyl)-amide

The title compound (108mg, 0.304mmol, 30.4%) was obtained as a slightly yellow solid from 1-phenethyl-1*H*-pyrrole-3-carbaldehyde described in Preparation Example 62 and quinoline-6-carboxylic acid (173mg, 1.0mmol) according to an analogous method to Example E-57.

- 25 <sup>1</sup>H-NMR Spectrum (DMSO-*d*<sub>6</sub>) δ(ppm) : 2.96 (2H, t, *J*=7.6Hz), 4.03 (2H, t, *J*=7.6Hz),

4.31 (2H, d, J=5.6Hz), 5.96 (1H, s), 6.64 (1H, s), 6.73 (1H, s), 7.14-7.28 (5H, m), 7.58 (1H, dd, J=4.0, 8.0Hz), 8.04 (1H, d, J=8.8Hz), 8.17 (1H, d, J=8.8Hz), 8.43 (1H, d, J=8.0Hz), 8.50 (1H, s), 8.90 (1H, t, J=5.6Hz), 8.95-8.99 (1H, m).

5 Example E-60. Quinoline-6-carboxylic acid (1-benzyloxy-1*H*-pyrrol-3-ylmethyl)-amide

Diethyl azodicarboxylate (154mg, 0.869mmol) was added dropwise to a solution of (1-benzyloxy-1*H*-pyrrol-3-yl)-methanol described in Preparation Example 64 (168mg, 0.828mmol), phthalimide (130mg, 0.869mmol) and triphenylphosphine (230mg, 0.869mmol) in dichloromethane at 0°C, and then, the solution was stirred at  
 10 room temperature for 8 minutes. NH silica gel was added to the reaction solution, the solvent was evaporated *in vacuo* for adsorption, purification was carried out by NH silica gel column chromatography (hexane : ethyl acetate = 4 : 1), and a colorless oil (100mg) was obtained. Ethanol (5mL) and hydrazine monohydrate (0.1mL) were added to this oil (100mg), and the solution was stirred under reflux for 15 minutes.  
 15 The reaction solution was allowed to room temperature, solid was eliminated by filtration, then, the solvent was evaporated, and an oil containing C-(1-benzyloxy-1*H*-pyrrol-3-yl)methylamine was obtained (80mg). The title compound (31mg, 0.086mmol) was obtained as a colorless oil from this oil (80mg) and quinoline-6-carboxylic acid according to an analogous method to Example Q-6.  
 20 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.28 (2H, d, J=5.6Hz), 5.11 (2H, s), 5.86-5.89 (1H, m), 6.78-6.81 (1H, m), 6.88-6.91 (1H, m), 7.34-7.42 (5H, m), 7.59 (1H, dd, J=4.0, 8.0Hz), 8.04 (1H, d, J=8.8Hz), 8.16 (1H, d, J=8.8Hz), 8.44 (1H, d, J=8.0Hz), 8.49 (1H, s), 8.92-8.98 (2H, m).

25 Example E-61. Quinoline-6-carboxylic acid (1-phenyl-1*H*-pyrrol-3-ylmethyl)-amide



The title compound (136mg, 0.415mmol, 47.8%) was obtained as a white solid from C-(1-phenyl-1*H*-pyrrol-3-yl)-methylaniline described in Preparation Example 74 (150mg, 0.87mmol) and quinoline-6-carboxylic acid (150mg, 0.87mmol) according to an analogous method to Example Q-6.

5 <sup>1</sup>H-NMR Spectrum (DMSO-*d*<sub>6</sub>) δ(ppm) : 4.42 (2H, d, J=5.6Hz), 6.28-6.31 (1H, m), 7.18-7.24 (1H, m), 7.29-7.35 (2H, m), 7.40-7.45 (2H, m), 7.50-7.65 (2H, m), 7.59 (1H, dd, J=4.0, 8.0Hz), 8.05 (1H, d, J=8.8Hz), 8.20 (1H, dd, J=2.0, 8.8Hz), 8.43-8.48 (1H, m), 8.53 (1H, d, J=2.0Hz), 8.96 (1H, dd, J=1.6, 4.0Hz), 9.01 (1H, t, J=5.6Hz).

10 Example E-62. Quinoline-6-carboxylic acid (2-benzyl-2*H*-tetrazol-5-ylmethyl)-amide

Sodium azide (260mg, 4.0mmol) and ammonium chloride (210mg, 4.0mmol) were suspended in a solution of quinoline-6-carboxylic acid cyanomethyl-amide described in Preparation Example E-1 (420mg, 2.0mmol) in *N,N*-dimethylformamide (15mL), and the solution was stirred at 100°C for 12 hours. Benzyl bromide  
15 (0.12mL, 1.0mmol) and potassium carbonate (400mg, 3.0mmol) were added to a solution having half the amount of the solution containing quinoline-6-carboxylic acid (2*H*-tetrazol-5-ylmethyl)amide obtained, which was then stirred for 20 minutes at 50°C. Water and ethyl acetate were added to the reaction solution, which was then partitioned, NH silica gel was added to the organic layer, and the solvent was  
20 evaporated *in vacuo* for adsorption, purification was carried out by NH silica gel column chromatography, and the title compound (20mg, 0.058mmol) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-*d*<sub>6</sub>) δ(ppm) : 4.78 (2H, d, J=5.6Hz), 5.92 (2H, s), 7.34-7.43 (5H, m), 7.62 (1H, dd, J=4.0, 8.4Hz), 8.09 (1H, d, J=8.8Hz), 8.19 (1H, dd, J=2.0, 8.8Hz), 8.48 (1H, dd, J=1.2, 8.4Hz), 8.54 (1H, d, J=2.0Hz), 9.00 (1H, dd, J=1.2,  
25

4.0Hz), 9.44 (1H, t, J=5.6Hz).

Example E-63. Quinoline-6-carboxylic acid (2-phenoxy-thiazol-5-ylmethyl)-amide

The title compound (17mg, 46μmol, 91%) was obtained as a white solid from  
5 C-(2-phenoxy-thiazol-5-yl)-methylamine described in Preparation Example 117  
(10mg, 50μmol) and quinoline-6-carboxylic acid (11mg, 60μmol) according to an  
analogous method to Example A-26.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.55 (2H, d, J=5.7Hz), 7.23 (1H, s), 7.27-7.31  
(3H, m), 7.45 (2H, dd, J=7.2, 8.8Hz), 7.59 (1H, dd, J=4.2, 8.8Hz), 8.06 (1H, d,  
10 J=8.8Hz), 8.14 (1H, dd, J=1.8, 8.8Hz), 8.46 (1H, d, J=7.1Hz), 8.5 (1H, d, J=1.8Hz),  
8.97 (1H, dd, J=1.7, 4.2Hz), 9.39 (1H, t, J=5.7Hz).

Example E-64. Quinoline-6-carboxylic acid  
(5-(3-cyano-phenoxy)-thiophen-2-ylmethyl)-amide

To a solution of C-(5-(3-bromophenoxy)-thiophen-2-yl)-methylamine  
15 described in Preparation Example 17 (200mg, 0.703mmol) and 6-quinolinecarboxylic  
acid (123mg, 0.703mmol) in tetrahydrofuran (5mL) were added  
benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (373mg,  
0.844mmol) and triethylamine (0.2mL, 1.41mmol), and the solution was stirred at  
room temperature for 2 hours. Ethyl acetate and water were added to the reaction  
20 solution, which was then partitioned, the organic layer was washed with water, and  
then, dried over anhydrous magnesium sulfate. The solvent was evaporated, the  
residue was purified by NH silica gel column chromatography (hexane : ethyl acetate),  
and a mixture of quinoline-6-carboxylic acid  
(5-(3-bromophenoxy)-thiophene-2-ylmethyl)-amide and debrominated compound  
25 (170mg, 55%) was obtained as a colorless oil.

Next, to a solution of the mixture of quinoline-6-carboxylic acid (5-(3-bromophenoxy)-thiophene-2-ylmethyl)-amide and debrominated compound (130mg, 0.303mmol) in N,N-dimethylformamide (5.0mL) were added zinc cyanide (71mg, 0.605mmol) and tetrakis(triphenylphosphine)palladium(0) (70mg, 0.061mmol) under nitrogen atmosphere, the solution was stirred at 100°C for 1 hour, and the solution was stirred at 140°C for 3 hours. The reaction solution was allowed to room temperature, ethyl acetate and water were added to the reaction solution, which was then partitioned, the organic layer was washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated, then, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate), and title compound (25mg, 21%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.80(2H, d, J=5.2Hz), 6.49(1H, d, J=4.0Hz), 6.68(1H, brs), 6.85(1H, d, J=3.6Hz), 7.30-7.45(4H, m), 7.48(1H, dd, J=4.4, 8.4Hz), 8.07(1H, dd, J=2.0, 8.8Hz), 8.17(1H, d, J=8.8Hz), 8.24-8.27(1H, m), 8.35(1H, d, J=2.0Hz), 8.99-9.01(1H, m).

Example E-65. Quinoline-6-carboxylic acid (5-(3-fluorophenoxy)thiophen-2-ylmethyl) amide

The title compound (100mg, 0.265mmol, 29.4%) was obtained as a reddish brown oil from C-(5-(3-fluorophenoxy)thiophen-2-yl)methylamine described in Preparation Example 23 and 6-quinolinecarboxylic acid according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.59 (2H, d, J=5.6Hz), 6.59 (1H, d, J=3.6Hz), 6.84 (1H, d, J=3.6Hz), 6.90-7.7.02 (3H, m), 7.39 (1H, ddd, J=8.0, 8.0, 8.0Hz), 7.60(1H, dd, J=4.0, 8.0Hz), 8.06(1H, d, J=8.8Hz), 8.16 (1H, dd, J=1.6, 8.8Hz), 8.45

(1H, d, J=8.0Hz), 8.51 (1H, s), 8.97(1H, d, J=4.0Hz), 9.37(1H, t, J=5.6Hz).

Example E-66. Quinoline-6-carboxylic acid (5-phenoxythiophen-2-ylmethyl) amide

Sodium hydride (3g, 74mmol, 60% in oil) was added to a solution of phenol  
 5 (7g,74mmol) in dimethylsulfoxide (40mL), which was then stirred at room  
 temperature for 10 minutes, and 5-nitrothiophene-2-carbaldehyde (10g, 64mmol) was  
 further added, followed by stirring for 15 minutes. Water and ethyl acetate were  
 added to the reaction solution, which was then partitioned, the organic layer was  
 washed with an aqueous solution of 2N sodium hydroxide twice and with water three  
 10 times, then, passed through a glass filter lined with silica gel, and eluted with ethyl  
 acetate. The solvent was evaporated *in vacuo*, and a yellow oil containing  
 5-phenoxy thiophene-2-carbaldehyde was obtained (500mg). This oil (500mg) was  
 dissolved in 7N ammonia/methanol solution (30mL), Raney nickel (1.5g) was added  
 thereto, and the solution was stirred overnight under hydrogen atmosphere. Raney  
 15 nickel was removed by filtering through Celite pad, then, the filtrate was concentrated  
*in vacuo*, the residue was purified by silica gel chromatography (ethyl acetate, then  
 ethyl acetate : methanol = 4 : 1) and a brown oil containing  
 C-(5-phenoxythiophen-2-yl)methylamine was obtained (40mg).

Then, to a solution of the obtained oil (40mg, 0.195mmol) and  
 20 6-quinolinecarboxylic acid (41mg, 0.234mmol) in N,N-dimethylformamide (5mL) were  
 added benzotriazol-1-yl-tris(dimethylamino)phosphonium hexafluorophosphate  
 (100mg, 0.234mmol) and triethylamine (0.054mL, 0.39mmol), and the solution was  
 stirred for 30 minutes at 60°C. Water and ethyl acetate were added to the reaction  
 solution, which was then partitioned, the organic layer was washed with water twice,  
 25 NH silica gel was added to this organic layer, the solvent was evaporated *in vacuo* for

adsorption, and purification was carried out by NH silica gel column chromatography (hexane : ethyl acetate = 3 : 1, then 1:1, then ethyl acetate). The solvent was evaporated *in vacuo*, then, the solid generated by adding diethyl ether to the residue was collected by filtration, and the title compound (40mg, 0.111mmol, 56.9%) was obtained as a pale yellow solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.57 (2H, d, J=5.6Hz), 6.51 (1H, d, J=3.6Hz), 6.81 (1H, d, J=3.6Hz), 7.06-7.15 (3H, m), 7.23-7.40 (2H, m), 7.59 (1H, dd, J=4.0, 8.0Hz), 8.06 (1H, d, J=8.8Hz), 8.16 (1H, dd, J=2.0, 8.8Hz), 8.45 (1H, dd, J=1.6, J=8.0Hz), 8.51 (1H, d, J=2.0Hz), 8.97 (1H, dd, J=1.6, 4.0Hz), 9.36 (1H, t, J=5.6Hz).

Example E-67. Quinoline-6-carboxylic acid (5-(4-fluorophenoxy)thiophen-2-yl)methyl amide

The title compound (38mg, 0.100mmol, 27.8%) was obtained as a white solid from C-(5-(4-fluorophenoxy)thiophen-2-yl)methylamine described in Preparation Example 28 and 6-quinolinecarboxylic acid according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.57 (2H, d, J=6.0Hz), 6.49 (1H, d, J=3.6Hz), 6.80 (1H, d, J=3.6Hz), 7.10-7.17 (2H, m), 7.17-7.24 (2H, m), 7.59 (1H, dd, J=4.0, 8.0Hz), 8.06 (1H, d, J=8.8Hz), 8.16 (1H, dd, J=2.0, J=8.8Hz), 8.45 (1H, dd, J=1.6, 8.0Hz), 8.51 (1H, d, J=2.0Hz), 8.97 (1H, dd, J=1.6, 4.0Hz), 9.36 (1H, t, J=6.0Hz).

Example E-68. Quinoline-6-carboxylic acid (5-(4-chloro-phenoxy)-thiophen-2-ylmethyl)-amide

The title compound (87mg, 0.22mmol, 76.1%) was obtained as a brown oil from the resulting C-(5-(4-chlorophenoxy)thiophen-2-yl)methylamine (70mg,

0.29mmol) and 6-quinolinecarboxylic acid (51mg, 0.29mmol) according to an analogous method to Example E-66.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.58 (2H, d, 5.6Hz), 6.55 (1H, d, J=4.0Hz), 6.83 (1H, d, J=4.0Hz), 7.08-7.14 (2H, m), 7.38-7.45 (2H, m), 7.59 (1H, dd, J=4.4, 8.0Hz), 8.06 (1H, d, J=8.8Hz), 8.16 (1H, dd, J=2.0, 8.8Hz), 8.46 (1H, dd, J=1.6, 8.0Hz), 8.51 (1H, d, J=2.0Hz), 8.97 (1H, dd, J=1.6, 4.4Hz), 9.36 (1H, t, J=5.6Hz).

Example E-69. Quinoline-6-carboxylic acid (4-(3-fluoro-phenoxy)-thiophen-2-ylmethyl)-amide

The title compound (24mg, 0.063mmol, 39.7%) was obtained as a white solid from C-(4-(3-fluorophenoxy)thiophen-2-yl)methylamine described in Preparation Example 35 (35mg, 0.16mmol) and 6-quinolinecarboxylic acid (33mg, 0.19mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.63 (2H, d, J=6.0Hz), 6.83-6.97 (5H, m), 7.38 (1H, ddd, J=8.0, 8.0, 8.0Hz), 7.60 (1H, dd, J=4.4, 8.0Hz), 8.07 (1H, d, J=8.8Hz), 8.17 (1H, dd, J=2.0, 8.8Hz), 8.47 (1H, dd, J=2.0, 8.0Hz), 8.53 (1H, d, J=2.0Hz), 8.97 (1H, dd, J=2.0, 4.4Hz), 9.39 (1H, t, J=6.0Hz).

Example E-70. Quinoline-6-carboxylic acid (5-benzyl-thiophen-2-ylmethyl)-amide

The title compound (40mg, 0.111mmol, 41.3 %) was obtained as a white solid from quinoline-6-carboxylic acid (46mg, 0.27mmol) and C-(5-benzyl-thiophen-2-yl)-methylamine described in Preparation Example 42 (54mg, 0.27mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.04 (2H, s), 4.57 (2H, d, J=5.6Hz), 6.71 (1H, d, J=3.6Hz), 6.84 (1H, d, J=3.6Hz), 7.14-7.30 (5H, m), 7.58 (1H, dd, J=4.0, 8.4Hz),

8.04 (1H, d, J=8.8Hz), 8.14 (1H, dd, J=2.0, 8.8Hz), 8.44 (1H, dd, J=2.0, 8.4Hz), 8.49 (1H, d, J=2.0Hz), 8.96 (1H, dd, J=2.0, 4.0Hz), 9.29 (1H, t, J=5.6Hz).

Example E-71. Quinoline-6-carboxylic acid

5 (5-(3-chloro-benzyl)-thiophen-2-ylmethyl)-amide

The title compound (73mg, 0.18mmol, 85.7%) was obtained as a white solid from C-(5-(3-chloro-benzyl)-2-yl)-methylaniline described in Preparation Example 45 (50mg, 0.21mmol) and 6-quinolinecarboxylic acid (40mg, 0.23mmol) according to an analogous method to Example Q-6.

10 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.07 (2H, s), 4.58 (2H, d, J=5.2Hz), 6.74 (1H, d, J=2.4Hz), 6.84 (1H, d, J=2.4Hz), 7.17-7.34 (4H, m), 7.59 (1H, dd, J=4.0, 8.4Hz), 8.05 (1H, d, J=8.4Hz), 8.14 (1H, d, J=8.4Hz), 8.44 (1H, d, J=8.4Hz), 8.49 (1H, s), 8.95 (1H, d, J=4.0Hz), 9.30(1H, t, J=5.2Hz).

15 Example E-72. Quinoline-6-carboxylic acid

(5-(3-fluoro-benzyl)-thiophen-2-ylmethyl)-amide

The title compound (75mg, 0.199mmol, 80.2%) was obtained as a light brown solid from quinoline-6-carboxylic acid (43mg, 0.248mmol) and 5-(3-fluoro-benzyl)-thiophene-2-carbaldehyde described in Preparation Example 53

20 (50mg, 0.226mmol) according to an analogous method to Example A-146.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.08 (2H, s), 4.58 (2H, d, J=5.6Hz), 6.74 (1H, d, J=3.2Hz), 6.85 (1H, d, J=3.2Hz), 6.98-7.10 (3H, m), 7.28-7.35 (1H, m), 7.59 (1H, dd, J=4.0, 8.0Hz), 8.05 (1H, d, J=8.8Hz), 8.15 (1H, dd, J=1.6, 8.8Hz), 8.44 (1H, d, J=8.0Hz), 8.49 (1H, d, J=1.6Hz), 8.95-8.99 (1H, m), 9.30 (1H, t, J=5.6Hz).

Example                      E-73.                      Quinoline-6-carboxylic                      acid  
(5-(5-methyl-thiophen-2-ylmethyl)-thiophen-2-ylmethyl)-amide

To a solution of (5-(5-methyl-thiophen-2-ylmethyl)-thiophen-2-yl)-methanol described in Preparation Example 67 (640mg, 2.86mmol), phthalimide (420mg, 2.86mmol) and triphenylphosphine (750mg, 2.86mmol) in tetrahydrofuran (7mL) was added diethyl azodicarboxylate (500mg, 2.86mmol) dropwise at 0°C, which was then stirred at room temperature for 15 minutes. Water and ethyl acetate were added to the reaction solution, which was then partitioned, silica gel was added, the solvent was concentrated *in vacuo* for adsorption, and purification was carried out by silica gel column chromatography (hexane : ethyl acetate = 8 : 1). The solvent was evaporated *in vacuo*, a brown solid (360mg, 1.02mmol, 35.6%) was obtained.

Ethanol (5mL) and hydrazine monohydrate (180mg, 3.06mmol) were added to the resulting solid (360mg), and the solution was stirred for 20 minutes at 90°C. The solution was allowed to room temperature, the solid was eliminated by filtration, and a pale yellow oil containing C-5-(5-methyl-thiophen-2-ylmethyl)-thiophen-2-yl)-methylamine was obtained (200mg, 0.896mmol, 89.6%). The title compound (101mg, 0.267mmol, 29.8%) was obtained as a white solid from this oil (200mg, 0.896mmol) and quinoline-6-carboxylic acid (160mg, 0.896mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.35 (3H, s), 4.20 (2H, s), 4.61 (2H, d, J=5.6Hz), 6.58-6.62 (1H, m), 6.69 (1H, d, J=3.2Hz), 6.75 (1H, d, J=3.2Hz), 6.86 (1H, d, J=3.2Hz), 7.61 (1H, dd, J=4.0, 8.4Hz), 8.08 (1H, d, J=8.8Hz), 8.18 (1H, dd, J=2.0, 8.8Hz), 8.47 (1H, d, J=8.4Hz), 8.52 (1H, d, J=2.0Hz), 8.97-9.01 (1H, m), 9.34 (1H, t, J=5.6Hz).



Example E-74. Quinoline-6-carboxylic acid  
(5-(5-methyl-furan-2-ylmethyl)-thiophen-2-ylmethyl)-amide

The title compound (3.0mg, 0.008mmol) was obtained as a brown oil from (5-(5-methyl-furan-2-ylmethyl)-thiophen-2-yl)-methanol described in Preparation

5 Example 70 (210mg, 1.0mmol) according to an analogous method to Example E-73.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 2.18 (3H, s), 4.04 (2H, s), 4.61 (2H, d, J=5.6Hz), 5.94 (1H, d, J=3.2Hz), 6.03 (1H, d, J=3.2Hz), 6.74 (1H, d, J=3.2Hz), 6.87 (1H, d, J=3.2Hz), 7.61 (1H, dd, J=4.0, 8.0Hz), 8.07 (1H, d, J=8.4Hz), 8.17 (1H, dd, J=1.6, 8.4Hz), 8.47 (1H, dd, J=2.0, 4.0Hz), 8.52 (1H, d, J=1.6Hz), 8.98 (1H, dd, J=2.0, 4.0Hz), 9.33 (1H, t, J=5.6Hz).

Example E-75. Quinoline-6-carboxylic acid  
(5-benzofuran-2-ylmethyl-thiophen-2-ylmethyl)-amide

The title compound (55mg, 0.13mmol) was obtained as a white solid from (5-benzofuran-2-ylmethyl-thiophene)-methanol described in Preparation Example 72 according to an analogous method to Example E-73.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.31 (2H, s), 4.62 (2H, d, J=5.6Hz), 6.68 (1H, s), 6.85 (1H, d, J=3.6Hz), 6.91 (1H, d, J=3.6Hz), 7.16-7.26 (2H, m), 7.46-7.64 (3H, m), 8.07 (1H, d, J=8.8Hz), 8.17 (1H, dd, J=2.0, 8.8Hz), 8.44-8.48 (1H, m), 8.52 (1H, d, J=2.0Hz), 8.96-9.00 (1H, m), 9.34 (1H, t, J=5.6Hz).

Example E-76. Quinoline-6-carboxylic acid (5-benzyloxy-thiophen-2-ylmethyl)-amide

To a solution of 5-benzyloxy-thiophene-2-carbonitrile described in Preparation Example 81 (30mg, 0.14mmol) in tetrahydrofuran (3mL) was added lithium aluminum hydride (21mg, 0.557mmol), which was then stirred for 1.5 hours at room

temperature. Sodium fluoride (240mg, 5.72mmol) was added to the reaction mixture, which was stirred for 2 hours, then, on an ice bath, 10% hydrous tetrahydrofuran (2mL) was added. The reaction mixture was filtered through Celite pad, the filtrate was concentrated and C-(5-benzyloxy-thiophen-2-yl)methylamine (32mg, 0.147mmol) was obtained as a crude product. The title compound (3mg, 0.008mmol, 5.4%) was obtained from this and quinoline-6-carboxylic acid (26mg, 0.15mmol) according to an analogous method to Example H-1.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.74 (2H, d, J=4.4Hz), 5.09 (2H, s), 6.15 (1H, d, J=4.0Hz), 6.52-6.62 (1H, m), 6.71 (1H, d, J=4.0Hz), 7.32-7.47 (5H, m), 7.50 (1H, dd, J=4.0, 8.4Hz), 8.07 (1H, dd, J=2.0, 8.8Hz), 8.18 (1H, d, J=8.8Hz), 8.27 (1H, dd, J=1.6, 8.4Hz), 8.34 (1H, d, J=2.0Hz), 9.02 (1H, dd, J=1.6, 4.0Hz).

Example E-77. Quinoline-6-carboxylic acid (5-(3-chloro-phenoxy)-thiophen-2-ylmethyl)-amide

The title compound (9.53mg) was obtained from C-(5-(3-chloro-phenoxy)-thiophen-2-yl)-methylamine described in Example A-73 (30mg, 0.13mmol) and quinoline-6-carboxylic acid (22mg, 0.13mmol) according to an analogous method to Example E-24. Trifluoroacetic acid salt of the title compound was obtained by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used).

MS m/e (ESI) 395.35(MH<sup>+</sup>)

Example E-78. 6-Quinolinecarboxylic acid (5-(2-methylpropenyl)thiophen-2-ylmethyl)-amide

The title compound (15mg, 0.0466mmol, 56.1%) was obtained as a white

solid from 6-quinolinecarboxylic acid (15mg, 0.083mmol) and C-(5-(2-methylpropenyl)-thiophen-2-yl)-methylaniline described in Preparation Example 145 (14mg, 0.083mmol) according to an analogous method to Example E-49.

5 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 1.91(3H, s), 1.96(3H, s), 4.83(2H, d, J=5.2Hz), 6.33(1H, s), 6.62(1H, t, J=5.2Hz), 6.75(1H, d, J=3.6Hz), 6.96(1H, d, J=3.6Hz), 7.47(1H, dd, J=4.0, 8.0Hz), 8.05(1H, dd, J=2.0, 8.8Hz), 8.13(1H, d, J=8.8Hz), 8.23(1H, dd, J=1.2, 8.0Hz), 8.32(1H, d, J=2.0Hz), 8.97(1H, dd, J=1.2, 4.0Hz).

10 Example E-79. Quinoline-6-carboxylic acid  
(5-(2-fluoro-phenoxy)-thiophen-2-ylmethyl)-amide

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.55 (2H, d, J=6.0Hz), 6.47 (1H, d, J=4.0Hz), 6.78 (1H, d, J=3.6Hz), 7.16-7.26 (3H, m), 7.32-7.39 (1H, m), 7.59 (1H, dd, J=4.0, J=8.0Hz), 8.06 (1H, d, J=8.8Hz), 8.15 (1H, dd, J=2.0, J=8.8Hz), 8.45 (1H, d, J=8.0Hz), 8.51 (1H, d, J=2.0Hz), 8.95-8.98 (1H, m), 9.35 (1H, t, J=6.0Hz).

Example E-80. Quinoline-6-carboxylic acid  
(5-pyridin-2-ylmethyl-thiophen-2-ylmethyl)-amide

20 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.20 (2H, s), 4.60 (2H, d, J=5.6Hz), 6.77 (1H, d, J=3.2Hz), 6.86 (1H, d, J=3.2Hz), 7.22 (1H, dd, J=5.2, J=7.6Hz), 7.31 (1H, d, J=7.6Hz), 7.61 (1H, dd, J=4.0, 8.0Hz), 7.71 (1H, ddd, J=1.6, J=7.6, J=7.6Hz), 8.07 (1H, d, J=8.0Hz), 8.17 (1H, dd, J=2.0, J=8.0Hz), 8.42-8.51 (2H, m), 8.52 (1H, d, J=2.0Hz), 8.98 (1H, dd, J=1.2, J=4.0Hz), 9.32 (1H, t, J=5.6Hz).

25 Example E-81. Quinoline-6-carboxylic acid

(5-benzo[1,3]dioxol-5-ylmethyl-thiophen-2-ylmethyl)-amide

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.97 (2H, s), 4.59 (2H, d, J=6.0Hz), 5.95 (2H, s), 6.69-6.75 (2H, m), 6.78-6.87 (3H, m), 7.61 (1H, dd, J=4.4, J=8.4Hz), 8.07 (1H, d, J=8.8Hz), 8.17 (1H, dd, J=2.0, J=8.8Hz), 8.46 (1H, dd, J=1.6, J=8.4Hz), 8.51 (1H, d, J=2.0Hz), 8.98 (1H, dd, J=1.6, J=4.4Hz), 9.31 (1H, t, J=6.0Hz).

Example E-82. Quinoline-6-carboxylic acid  
(5-(3-hydroxy-phenoxy)-thiophen-2-ylmethyl)-amide

To a solution of C-(5-(3-benzyloxy-phenoxy)-thiophen-2-yl)-methylaniline described in Preparation Example 18 (180mg, 0.578mmol) and 6-quinolinecarboxylic acid (100mg, 0.578mmol) in tetrahydrofuran (5mL) were added benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (307mg, 0.694mmol) and triethylamine (0.16mL, 1.16mmol), and the solution was stirred overnight at room temperature. Ethyl acetate and water were added to the reaction solution, which was then partitioned, the organic layer was washed with water, and then, dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate), and quinoline-6-carboxylic acid (5-(3-benzyloxy-phenoxy)-thiophen-2-ylmethyl)-amide (73mg, 27%) was obtained as a pale yellow solid.

Trifluoroacetic acid (1.0mL) and thioanisole (100μl) were added to the resulting quinoline-6-carboxylic acid (5-(3-benzyloxy-phenoxy)-thiophene-2-ylmethyl)-amide (73mg, 0.156mmol), and the solution was stirred for 30 minutes at room temperature. The reaction solution was neutralized with an aqueous solution of saturated sodium bicarbonate, then, extracted with ethyl acetate and dried over anhydrous magnesium sulfate. The

solvent was evaporated, the residue was purified by NH silica gel column chromatography (ethyl acetate : methanol), and the title compound (47mg, 80%) was obtained as a colorless solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.73(2H, d, J=5.6Hz), 6.39(1H, d, J=3.6Hz),  
5 6.59-6.62(2H, m), 6.64-6.67(1H, m), 6.74(1H, d, J=3.6Hz), 6.83(1H, brs), 7.17(1H, t, J=8.4Hz), 7.49(1H, dd, J=4.4, 8.4Hz), 8.05(1H, dd, J=2.0, 8.4Hz), 8.14(1H, d, J=8.4Hz), 8.24-8.26(1H, m), 8.33(1H, d, J=2.0Hz), 8.98-8.99(1H, m).

Example F-1. Cinnoline-6-carboxylic acid 3-phenoxy-benzylamide

10 To a solution of cinnoline-6-carboxylic acid methyl ester described in Preparation Example F-4 (16mg, 0.085mmol) in ethanol (1mL) was added an aqueous solution of 1N sodium hydroxide (0.7mL), and the solution was stirred at room temperature for 2 hours. 1N Hydrochloric acid was added to the reaction mixture to adjust the pH to 4, toluene was added, and the solution was concentrated  
15 *in vacuo*. To a solution of the obtained residue in N,N-dimethylformamide (2mL) were added 4-phenoxybenzylamine described in Preparation Example 3 (17mg, 0.085mmol), benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (48mg, 0.108mmol) and triethylamine (24μl, 0.172mmol), and the solution was stirred at room temperature for 14 hours. Water was added to the  
20 reaction mixture, which was extracted with ethyl acetate, the organic layer was washed with an aqueous solution of saturated sodium bicarbonate, and then concentrated. The residue was purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), and the title compound (4.4mg, 0.0093mmol, 11%) was obtained as  
25 a trifluoroacetic acid salt.

MS m/e(ESI) 356.44 (MH<sup>+</sup>)

Example G-1. Isoquinoline-6-carboxylic acid 3-phenoxybenzylamide

The title compound (3.4mg, 33%) was obtained as a colorless oil from  
5 isoquinoline-6-carboxylic acid described in Preparation Example G-1 (5mg,  
0.0289mmol) and 4-phenoxybenzylamine described in Preparation Example 3 (6mg,  
0.0289mmol) according to an analogous method to Example L-4.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.70(2H, d, J=5.6Hz), 6.57(1H, brs),  
6.94-6.96(1H, m), 7.02-7.04(3H, m), 7.11-7.15(2H, m), 7.32-7.37(3H, m),  
10 7.72-7.74(1H, m), 7.96-7.98(1H, m), 8.05-8.07(1H, m), 8.26(1H, s), 8.61-8.62(1H, m),  
9.32-9.33(1H, m).

Example H-1. Quinazoline-6-carboxylic acid 3-phenoxybenzylamide

To a solution of quinazoline-6-carboxylic acid obtained in Preparation  
15 Example H-3 (9mg, 0.052mmol) in N,N-dimethylformamide (3mL) were added  
3-phenoxybenzylamine described in Preparation Example 4 (11mg, 0.052mmol),  
benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (28mg,  
0.062mmol) and triethylamine (17μl, 0.125mmol), and the solution was stirred for 2  
days at room temperature. Water was added to the reaction mixture, which was  
20 extracted with ethyl acetate, and concentrated. The residue was purified by NH  
silica gel column chromatography (hexane-ethyl acetate), and the title compound  
(11mg, 0.031mmol, 50%) was obtained.

<sup>1</sup>H-NMR Spectrum (CD<sub>3</sub>OD) δ(ppm) : 4.62 (2H, s), 6.88 (1H, dd, J=8.0, 1.2Hz), 6.98  
(2H, dd, J=1.2, 8.0Hz), 7.09 (1H, s), 7.07 (1H, dd, J=7.6, 8.0Hz), 7.15 (1H, d,  
25 J=7.6Hz), 7.29-7.35 (3H, m), 8.10 (1H, d, J=8.8Hz), 8.40 (1H, dd, J=2.0, 8.8Hz), 8.57

(1H, d, J=2.0Hz), 9.32 (1H, s), 9.61 (1H, s).

Example I-1. Quinoxaline-6-carboxylic acid 3-phenoxybenzylamide

To a solution of quinoxaline-6-carboxylic acid described in Preparation  
5 Example I-1 (15mg, 0.063mmol) and 4-phenoxybenzylamine described in  
Preparation Example 3 (13mg, 0.063mmol) in N,N-dimethylformamide (2mL) were  
added benzotriazol-1-yloxytris(pyrrolidino)phosphonium hexafluorophosphate (36mg,  
0.069mmol) and triethylamine (19μl, 0.14mmol), which was then stirred at room  
temperature for 24 hours. The reaction mixture was concentrated, the residue was  
10 purified by reverse phase high performance liquid chromatography (acetonitrile-water  
mobile phase (containing 0.1% trifluoroacetic acid) was used), and the title  
compound (12mg, 0.025mmol, 40%) was obtained as a trifluoroacetic acid salt.

MS m/e(ESI) 356.37(MH<sup>+</sup>)

15 Example J-1. [1,8]Naphthylidene-3-carboxylic acid 3-phenoxybenzylamide

Tetrahydrofuran (1mL), methanol (0.1mL) and water (0.1mL) were added to  
[1,8]naphthylidene-3-carboxylic acid ethyl ester described in Preparation Example J-7  
(8.1mg, 0.040mmol) and lithium hydroxide monohydrate (3.4mg, 0.080mmol), and  
the solution was stirred at 50°C for 1 hour. The solvent was evaporated *in vacuo*,  
20 then, the obtained residue and 3-phenoxybenzylamine (5.0mg, 0.025mmol) were  
reacted according to an analogous method to Example Q-6, and trifluoroacetic acid  
salt of the title compound (3.7mg, 0.0079mmol, 20%) was obtained.

MS m/e (ESI) 356.3 (MH<sup>+</sup>)

25 Example K-1. 2-Methyl-benzoxazole-6-carboxylic acid 3-phenoxybenzylamide

The title compound (22mg, 0.061mmol, 72%) was obtained from 2-methyl-benzoxazole-6-carboxylic acid described in Preparation Example K-2 (15mg, 0.085mmol) and 4-phenoxybenzylamine described in Preparation Example 3 (17mg, 0.085mmol) according to an analogous method to Example H-1.

5 <sup>1</sup>H-NMR Spectrum (CD<sub>3</sub>OD) δ(ppm) : 2.66 (3H, s), 4.56 (2H, s), 6.80-7.15 (6H, m), 7.22-7.78 (3H, m), 7.62-7.68 (1H, m), 7.81-7.86 (1H, m), 8.00-8.04 (1H, m), 9.08(1H, brs).

Example L-1. Benzothiazole-6-carboxylic acid

10 (5-(3-fluoro-benzyl)-furan-2-ylmethyl)-amide

The title compound (290mg, 0.791mmol, 76%) was obtained from benzothiazole-6-carboxylic acid (188mg, 1.05mmol) and C-(5-(3-fluoro-benzyl)-furan-2-yl)-methylamine described in Preparation Example 84 (236mg, 1.15mmol) according to an analogous method to Example H-1 (with the  
15 proviso that only the reaction temperature was changed to 60°C).

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.96 (2H, s), 4.43 (2H, d, J=5.2Hz), 6.06 (1H, d, J=3.2Hz), 6.20 (1H, d, J=3.2Hz), 7.00-7.09 (3H, m), 7.29-7.36 (1H, m), 7.99 (1H, dd, J=1.6, 8.4Hz), 8.12 (1H, d, J=8.4Hz), 8.64 (1H, d, J=1.6Hz), 9.07 (1H, t, J=5.2Hz), 9.51 (1H, s).

20

Example L-2. Benzothiazole-6-carboxylic acid 4-benzyloxybenzylamide

The title compound (41mg, 47%) was obtained as a white solid from benzothiazole-6-carboxylic acid (42mg, 0.234mmol) and 4-benzyloxybenzylamine described in Preparation Example 1 (50mg, 0.234mmol) according to an analogous  
25 method to Example E-8.



<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.62(2H, d, J=5.2Hz), 5.08(2H, s), 6.40(1H, brs), 6.98(2H, d, J=8.8Hz), 7.30-7.35(3H, m), 7.37-7.45(4H, m), 7.85-7.88(1H, m), 8.16(1H, d, J=8.4Hz), 8.49(1H, d, J=1.6Hz), 9.11(1H, s).

5 Example L-3. Benzothiazole-6-carboxylic acid 3-phenoxybenzylamide

To a solution of 3-phenoxybenzylamine described in Preparation Example 4 (33mg, 0.167mmol) and benzothiazole-6-carboxylic acid (30mg, 0.167mmol) in tetrahydrofuran (1mL) were added benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (89mg, 0.20mmol) and triethylamine (28μl, 0.20mmol), and the solution was stirred at room temperature for 17 hours. The solvent was evaporated, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate), and the title compound (37mg, 62%) was obtained as a colorless oil.

10 <sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.68(2H, d, J=6.0Hz), 6.50(1H, brs), 6.94(1H, dd, J=2.0, 8.0Hz), 7.02-7.04(3H, m), 7.11-7.15(2H, m), 7.31-7.37(3H, m), 7.88(1H, dd, J=1.6, 8.8Hz), 8.18(1H, d, J=8.8Hz), 8.49(1H, d, J=1.6Hz), 9.13(1H, s).

Example L-4. Benzothiazole-6-carboxylic acid 4-(3-fluoro-benzyloxy)-benzylamide

To a solution of 4-(3-fluorobenzyloxy)-benzylamine described in Preparation Example 6 (129mg, 0.558mmol) and benzothiazole-6-carboxylic acid (100mg, 0.558mmol) in tetrahydrofuran (5mL) were added benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (296mg, 0.670mmol) and triethylamine (93μl, 0.670mmol), and the solution was stirred overnight at room temperature. Ethyl acetate and water were added to the reaction solution, which was then partitioned, the organic layer was washed with water, and

then, dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate), and the title compound (148mg, 68%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.64(2H, d, J=5.6Hz), 5.08(2H, s), 6.42(1H, brs),  
 5 6.97(2H, d, J=8.8Hz), 7.02(1H, td, J=2.8, 8.4Hz), 7.15-7.21(2H, m), 7.31-7.38(3H, m),  
 7.88(1H, dd, J=1.6Hz, 8.4Hz), 8.17(1H, d, J=8.8Hz), 8.50(1H, d, J=1.6Hz), 9.12(1H, s).

Example L-5. N-Benzothiazol-6-yl-2-(3-phenoxy-phenyl)-acetamide

10 The title compound (118mg, 95%) was obtained as a colorless oil from 6-aminobenzothiazole (50mg, 3.33mmol) and 3-phenoxyphenylacetic acid (76mg, 3.33mmol) according to an analogous method to Example L-4.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.75(2H, s), 6.97-7.22(6H, m), 7.31(1H, brs),  
 7.34-7.40(4H, m), 8.01(1H, d, J=8.8Hz), 8.50(1H, d, J=2.0Hz), 8.91(1H, s).

15

Example L-6. Benzothiazole-6-carboxylic acid  
(5-(3-fluorophenoxy)thiophene-2-ylmethyl) amide

The title compound (100mg, 0.26mmol, 48.2%) was obtained as a white solid from benzotriazole-6-carboxylic acid (96mg, 0.54mmol) and  
 20 C-(5-(3-fluorophenoxy)thiophen-2-yl)methylamine described in Preparation Example 23 (120mg, 0.54mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.57 (2H, d, 5.6Hz), 6.58 (1H, d, J=3.6Hz),  
 6.83 (1H, d, J=3.6Hz), 6.89-7.00 (3H, m), 7.39 (1H, ddd, J=8.0, 8.0, 8.0Hz), 8.00 (1H, dd, J=1.6, 8.8Hz), 8.14 (1H, d, J=8.8Hz), 8.66 (1H, d, J=1.6Hz), 9.27 (1H, t, J=5.6Hz),  
 25 9.51 (1H, s).

Example L-7. Benzothiazole-6-carboxylic acid (5-phenoxythiophen-2-ylmethyl)-amide

The title compound (97mg, 0.265mmol, 54.0 %) was obtained as a light brown solid from benzothiazole-6-carboxylic acid (87mg, 0.49mmol) and C-(5-phenoxy-thiophen-2-yl)-methylamine described in Preparation Example 26 (100mg, 0.49mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.55 (2H, d, J=5.6Hz), 6.49 (1H, d, J=3.6Hz), 6.79 (1H, d, J=3.6Hz), 7.05-7.15 (3H, m), 7.30-7.40 (2H, m), 7.99 (1H, dd, J=1.6, 8.8Hz), 8.13 (1H, d, J=8.8Hz), 8.66 (1H, d, J=1.6Hz), 9.25 (1H, t, J=5.6Hz), 9.51 (1H, s).

Example L-8. Benzothiazole-6-carboxylic acid (5-(3-chloro-benzyl)-thiophen-2-ylmethyl)-amide

The title compound (64mg, 0.16mmol, 47.2%) was obtained as a white solid from C-(5-(3-chloro-benzyl-2-yl)-methylamine described in Preparation Example 45 (80mg, 0.34mmol) and benzothiazole-6-carboxylic acid (66mg, 0.37mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.06 (2H, s), 4.56 (2H, d, J=5.6Hz), 6.74 (1H, d, J=3.6Hz), 6.84 (1H, d, J=3.6Hz), 7.18-7.34 (4H, m), 7.98 (1H, dd, J=2.0, 8.8Hz), 8.12 (1H, d, J=8.8Hz), 8.64 (1H, d, J=2.0Hz), 9.21 (1H, t, J=5.6Hz), 9.51 (1H, s).

Example L-9. Benzothiazole-6-carboxylic acid (5-(3-chloro-phenoxy)-thiophen-2-ylmethyl) amide

The title compound (7.28mg) was obtained from C-(5-(3-chloro-phenoxy)-thiophen-2-yl)-methylamine described in Example A-73

(30mg, 0.13mmol) and benzothiazole-6-carboxylic acid (22mg, 0.13mmol) according to an analogous method to Example E-24. Trifluoroacetic acid salt of the title compound (7.28mg) was obtained by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used).

MS m/e (ESI) 401.32(MH<sup>+</sup>)

Example L-10. Benzothiazole-6-carboxylic acid  
(5-(2-fluoro-phenoxy)-thiophen-2-ylmethyl)-amide

Trifluoroacetic acid salt of the title compound (10.7mg, 0.021mmol, 14.3%) was obtained as a brown oil from benzothiazole-6-carboxylic acid (27.4mg, 0.15mmol) and C-(5-(2-fluoro-phenoxy)-thiophen-2-yl)-methylamine described in Preparation Example 161 (33.5mg, 0.15mmol) according to an analogous method to Example A-75.

MS m/e (ESI) 385(MH<sup>+</sup>)

Example L-11. Benzothiazole-6-carboxylic acid  
((5-(3-cyano-phenoxy)-thiophen-2-ylmethyl)-amide

To a solution of C-(5-(3-bromophenoxy)-thiophen-2-yl)-methylamine described in Preparation Example 17 (141mg, 0.496mmol) and benzothiazole-6-carboxylic acid (89mg, 0.496mmol) in tetrahydrofuran (5mL) were added benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (263mg, 0.595mmol) and triethylamine (0.14mL, 0.992mmol), and the solution was stirred at room temperature for 3 hours. Ethyl acetate and water were added to the reaction solution, which was then partitioned, the organic layer was washed with

water, and then, dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate), and a mixture of benzothiazole-6-carboxylic acid (5-(3-bromophenoxy)-thiophene-2-ylmethyl)-amide and debrominated compound (120mg, 53%) was obtained as a yellow oil.

Next, to a solution of the mixture of benzothiazole-6-carboxylic acid (5-(3-bromophenoxy)-thiophene-2-ylmethyl)-amide and debrominated compound (120mg, 0.269mmol) in N,N-dimethylformamide (3.0mL) were added zinc cyanide (63mg, 0.538mmol) and tetrakis(triphenylphosphine)palladium(0) (62mg, 0.054mmol) under nitrogen atmosphere, which was then stirred at 140°C for 14 hours, and the solution was stirred at 140°C for 3 hours. The reaction solution was allowed to room temperature, ethyl acetate and water were added to the reaction solution, which was then partitioned, the organic layer was washed with water and dried over anhydrous magnesium sulfate. After removing the solvent, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate), and the title compound (6.2mg, 6%) was obtained as a colorless oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.78(2H, d, J=6.0Hz), 6.48(1H, d, J=4.0Hz), 6.57(1H, brs), 6.83(1H, d, J=4.0Hz), 7.30(1H, s), 7.32-7.35(1H, m), 7.39-7.43(2H, m), 7.90(1H, dd, J=1.6, 8.8Hz), 8.19(1H, d, J=8.8Hz), 8.51(1H, d, J=1.6Hz), 9.13(1H, s).

#### Example M-1. Benzof[1,2 5]thiadiazole-5-carboxylic acid 3-phenoxybenzylamide

The title compound (43mg, 71%) was obtained as a colorless solid from 2,1,3-benzothiadiazole-5-carboxylic acid (30mg, 0.167mmol) obtained by hydrolysis of benzo-2,1,3-thiadiazole-5-carboxylic acid methyl ester with sodium hydroxide and 4-phenoxybenzylamine described in Preparation Example 3 (33mg, 0.167mmol)

according to an analogous method to Example L-4.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.67(2H, d, J=6.0Hz), 6.65(1H, brs), 6.94(1H, dd, J=2.4Hz, 8.4Hz), 7.01-7.03(3H, m), 7.10-7.14(2H, m), 7.30-7.36(3H, m), 8.01-8.07(2H, m), 8.37(1H, s).

5

Example O-1. 2,3-Dihydro-1*H*-pyrrolo[2,3-*b*]pyridine-5-carboxylic acid (5-phenoxy-thiophen-2-ylmethyl)-amide

The title compound (15mg, 43μmol, 47%) was obtained as a white solid from 2,3-dihydro-1*H*-pyrrolo[2,3-*b*]pyridine-5-carboxylic acid described in Preparation

10 Example O-2 (15mg, 91μmol) and C-(5-phenoxy-thiophen-2-yl)-methylamine described in Preparation Example 26 (19mg, 91μmol) according to an analogous method to Example A-26.

<sup>1</sup>H-NMR Spectrum (DMSO-*d*<sub>6</sub>) δ(ppm) : 2.99 (2H, t, J=9.0Hz), 3.53 (2H, t, J=8.6Hz), 4.46 (2H, d, J=6.0Hz), 6.49 (1H, d, J=3.8Hz), 6.74 (1H, d, J=3.7Hz), 7.00 (1H, s),  
15 7.09 (2H, d, J=8.6Hz), 7.14 (1H, t, J=7.5Hz), 7.38 (2H, t, J=7.5Hz), 7.63 (1H, s), 8.25 (1H, s), 8.71-8.77 (1H, m).

Example P-1. Furo[2,3-*b*]pyridine-5-carboxylic acid (5-phenoxy-thiophen-2-ylmethyl)-amide

20 Furo[2,3-*b*]pyridine-5-carboxylic acid (31mg) was obtained as a lithium salt from furo[2,3-*b*]pyridine-5-carboxylic acid ethyl ester described in Preparation Example P-4 (33mg, 0.17mmol) according to an analogous method to Example T-2.

The title compound (28mg, 80μmol, 79%) was obtained as a white solid from the lithium salt (17mg) of the resulting furo[2,3-*b*]pyridine-5-carboxylic acid  
25 (31mg) and C-(5-phenoxy-thiophen-2-yl)-methylamine described in Preparation

Example 26 (23mg, 0.11mmol).

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.57 (2H, d, J=5.9Hz), 6.52 (1H, d, J=3.8Hz), 6.82 (1H, d, J=3.7Hz), 7.09-7.17 (4H, m), 7.39 (2H, t, J=8.4Hz), 8.21 (1H, d, J=2.6Hz), 8.58 (1H, d, J=2.2Hz), 8.79 (1H, d, J=2.2Hz), 9.34 (1H, t, J=6.0Hz).

5

Example Q-1. Imidazo[1,2-a]pyridine-6-carboxylic acid  
(5-benzyl-furan-2-ylmethyl)-amide

7N Ammonia/methanol(40mL) and Raney nickel(3g) were added to 5-benzyl-furan-2-carbaldehyde described in Preparation Example 39 (2.5g, 13mmol), and the solution was stirred for 22 hours under hydrogen atmosphere at room temperature. After removing the catalyst by filtering through Celite pad, the solvent was evaporated *in vacuo* and C-(5-benzyl-furan-2-yl)methylamine (1.6g, 8.6mmol, 65.8%) was obtained.

The title compound (150mg, 0.45mmol, 45.3%) was obtained as a white solid from the resulting C-(5-benzylfuran-2-yl)methylamine (200mg, 1.07mmol) and imidazo[1,2-a]pyridine-6-carboxylic acid (170mg, 1.07mmol) according to an analogous method to Example Q-6.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.92 (2H, s), 4.41 (2H, d, J=5.2Hz), 6.01 (1H, d, J=2.4Hz), 6.19 (1H, d, J=2.4Hz), 7.16-7.30 (5H, m), 7.56-7.66 (3H, m), 8.03 (1H, s), 9.00 (1H, t, J=5.2Hz), 9.10 (1H, s).

20

Example Q-2. Imidazo[1,2-a]pyridine-6-carboxylic acid  
(5-(3-fluoro-benzyl)-furan-2-ylmethyl)-amide

The title compound (363mg, 1.04mmol, 90%) was obtained from imidazo[1,2-a]pyridine-6-carboxylic acid (188mg, 1.16mmol) and

25

C-(5-(3-fluoro-benzyl)-furan-2-yl)-methylaniline described in Preparation Example 84 (286mg, 1.39mmol) according to an analogous technique to Example H-1 (with the proviso that only the reaction temperature was changed to 60°C).

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 3.97 (2H, s), 4.42 (2H, d, J=5.2Hz), 6.07 (1H, d, J=3.2Hz), 6.21 (1H, d, J=3.2Hz), 7.00-7.08 (3H, m), 7.29-7.37 (1H, m), 7.56-7.66 (3H, m), 8.04 (1H, s), 9.01 (1H, t, J=5.2Hz), 9.10 (1H, dd, J=1.2, 1.6Hz).

Example Q-3. Imidazo[1,2-a]pyridine-6-carboxylic acid 4-benzyloxybenzylamide

The title compound (121mg, 55%) was obtained as a white solid from imidazo[1,2-a]pyridine-6-carboxylic acid (100mg, 0.617mmol) and 4-benzyloxybenzylamine described in Preparation Example 1 (132mg, 0.617mmol) according to an analogous method to Example E-8.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.59(2H, d, J=5.6Hz), 5.07(2H, s), 6.44(1H, brs), 6.96-6.98(2H, m), 7.27-7.30(3H, m), 7.33-7.44(5H, m), 7.59-7.61(1H, m), 7.65-7.66(1H, m), 7.69-7.70(1H, m), 8.83-8.84(1H, m).

Example Q-4. Imidazo[1,2-a]pyridine-6-carboxylic acid 3-phenoxybenzylamide

The title compound (22mg, 35%) was obtained as a colorless oil from imidazo[1,2-a]pyridine-6-carboxylic acid (30mg, 0.185mmol) and 3-phenoxybenzylamine described in Preparation Example 3 (37mg, 0.185mmol) according to an analogous method to Example L-3.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.64(2H, d, J=5.6Hz), 6.60(1H, brs), 6.92-6.94(1H, m), 7.00-7.02(3H, m), 7.08-7.14(2H, m), 7.30-7.41(4H, m), 7.59-7.61(1H, m), 7.69(1H, s), 7.69-7.70(1H, m), 8.83(1H, s).



Example                      Q-5.                      Imidazo[1,2-a]pyridine-6-carboxylic                      acid  
4-(3-fluoro-benzyloxy)-benzylamide

The title compound (64mg, 45%) was obtained as a white solid from 4-(3-fluorobenzyloxy)-benzylamine described in Preparation Example 6 (87mg, 0.376mmol) and imidazo[1,2-a]pyridine-6-carboxylic acid (61mg, 0.376mmol) according to an analogous method to Example L-4.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.60(2H, d, J=5.6Hz), 5.07(2H, s), 6.44(1H, brs), 6.96(2H, d, J=8.8Hz), 7.02(1H, dt, J=2.4, 8.4Hz), 7.15(1H, d, J=9.6Hz), 7.19(1H, d, J=8.4Hz), 7.30(2H, d, J=8.4Hz), 7.32-7.40(2H, m), 7.61(1H, d, J=9.6Hz), 7.67(1H, s), 7.70(1H, s), 8.84(1H, s).

Example                      Q-6.                      Imidazo[1,2-a]pyridine-6-carboxylic                      acid  
(5-(3-fluorophenoxy)thiophen-2-ylmethyl) amide

To a solution of imidazo[1,2-a]pyridine-6-carboxylic acid (87mg, 0.54mmol) and C-(5-(3-fluorophenoxy)thiophen-2-yl)methylamine described in Preparation Example 23 (120mg, 0.54mmol) in N,N-dimethylformamide (5mL) were added benzotriazol-1-yl-tris(dimethylamino)phosphonium hexafluorophosphate (240mg, 0.54mmol) and triethylamine (0.15mL, 1.08mmol), and the solution was stirred for 40 minutes at 80°C. Water and ethyl acetate were added to the reaction solution, which was then partitioned, and the organic layer was washed with water twice. Silica gel was added to the organic layer, the solvent was evaporated *in vacuo* for adsorption, purification was carried out by silica gel column chromatography (hexane : ethyl acetate = 1 : 1, then ethyl acetate), and the title compound (90mg, 0.25mmol, 45.4%) was obtained as a light brown oil.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.55 (2H, d, J=5.6Hz), 6.58 (1H, d, J=4.0Hz),

6.83 (1H, d, J=4.0Hz), 6.90-7.00 (3H, m), 7.40 (1H, ddd, J=8.0, 8.0, 8.0Hz), 7.57-7.66 (3H, m), 8.04 (1H, s), 9.12 (1H, d, J=0.8Hz), 9.20 (1H, t, J=5.6Hz).

Example Q-7. Imidazo[1,2-a]pyridine-6-carboxylic acid

5 (5-phenoxy-thiophen-2-ylmethyl)-amide

The title compound (160mg, 0.458mmol, 93.5 %) was obtained as a light brown oil from imidazo[1,2-a]pyridine-6-carboxylic acid (80mg, 0.49mmol) and C-(5-phenoxythiophen-2-yl)methylamine described in Preparation Example 26 (100mg,0.49mmol) according to an analogous method to Example Q-6.

10 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.53 (2H, d, J=5.6Hz), 6.50 (1H, d, J=4.0Hz), 6.79 (1H, d, J=4.0Hz), 7.04-7.15 (3H, m), 7.32-7.40 (2H, m), 7.56-7.66 (3H, m), 8.03 (1H, s), 9.08-9.13 (1H, m), 9.19 (1H, t, J=5.6Hz).

Example Q-8. Imidazo[1,2-a]pyridine-6-carboxylic acid

15 (5-(3-chloro-phenoxy)-thiophen-2-ylmethyl)-amide

To a solution of C-(5-(3-chloro-phenoxy)-thiophen-2-yl)-methylamine described in Preparation Example 167 (104mg, 0.434mmol) and imidazo[1,2-a]pyridine-6-carboxylic acid (77mg, 0.477mmol) in N,N-dimethylformamide (3mL) were added benzotriazol-1-yloxy  
20 tris(dimethylamino)phosphonium hexafluorophosphate (250mg, 0.564mmol) and triethylamine (181μL) at room temperature, and the solution was stirred at room temperature for 4 hours. Water was added to the reaction solution, which was then extracted with ethyl acetate, and the organic layer was washed with water and brine. Anhydrous magnesium sulfate was added to the organic layer for drying, filtration  
25 was carried out, then, the solvent was evaporated *in vacuo*, the residue was purified

by NH silica gel column chromatography (ethyl acetate / methanol = 40 / 1), and the title compound (149mg, 89%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.55(2H, d, J=5.3Hz), 6.59(1H, dd, J=1.1, 3.7Hz), 6.83(1H, d, J=3.7Hz), 7.06(1H, dd, J=1.1, 8.2Hz), 7.13(1H, s), 7.20(1H, d, J=8.1Hz), 7.40(1H, t, 8.1Hz), 7.58-7.64(3H, m), 8.05(1H, s), 9.12(1H, s), 9.23(1H, t, J=5.5Hz).

Example R-1. 1H-Pyrrolo[2,3-b]pyridine-5-carboxylic acid  
(5-phenoxy-thiophen-2-ylmethyl)-amide

The title compound (22mg, 63μmol, 68%) was obtained from 1H-pyrrolo[2,3-b]pyridine-5-carboxylic acid described in Preparation Example R-7 (15mg, 93μmol) and C-(5-phenoxy-thiophen-2-yl)-methanamine described in Preparation Example 26 (19mg, 93μmol) according to an analogous method to reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used).

MS m/e (ESI) 350.26(MH<sup>+</sup>)

Example R-2. 6-Amino-1H-pyrrolo[2,3-b]pyridine-5-carboxylic acid  
(5-phenoxy-thiophen-2-ylmethyl)-amide

To a solution of 6-amino-1H-pyrrolo[2,3-b]pyridine-5-carboxylic acid ethyl ester described in Preparation Example R-2 (95mg, 0.46mmol) in ethanol (10mL) was added an aqueous solution of 1N sodium hydroxide (5mL, 5mmol), which was then heated for 3 hours in an oil bath at 98°C. After cooling the reaction solution, the reaction solution was concentrated until it reached 1/3 of its volume, neutralized

with 1N hydrochloric acid, and further concentrated. The resulting crude product was suspended in N,N-dimethylformamide (5mL), triethylamine (0.096mL, 0.69mmol), benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (153mg, 0.35mmol) and C-(5-phenoxy-thiophen-2-yl)-methylamine described in Preparation Example 26 (71mg, 0.35mmol) were added thereto, followed by stirring at room temperature for 15 hours. After the reaction was completed, the reaction solution was poured into brine, and extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, then concentrated, the obtained residue was purified by silica gel column chromatography (ethyl acetate : hexane = 2 : 1), and the title compound (31mg, 0.085mmol, 18.5%) was obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.49 (2H, d, J=5.6Hz), 6.24 (1H, dd, J=2.0, 3.6Hz), 6.50 (1H, d, J=4.0Hz), 6.78 (1H, d, J=4.0Hz), 6.80 (2H, brs), 7.04 (1H, dd, J=2.0, 3.2Hz), 7.07-7.16 (3H, m), 7.35-7.41 (2H, m), 8.14 (1H, s), 8.91-8.95 (1H, m), 11.0(1H, brs).

#### Example S-1. Pyrrolo[3,2-b]pyridine-1-carboxylic acid 3-phenoxy-benzylamide

1H-Pyrrolo[3,2-b]pyridine described in Preparation Example S-4 (44mg, 0.37mmol) was dissolved in N,N-dimethylformamide (3mL), sodium hydride (18mg, 0.45mmol, 60% in oil) was added thereto, followed by stirring at room temperature for 30 minutes. Next, (4-phenoxy-benzyl)-carbamic acid phenyl ester described in Preparation Example 75 (143mg, 0.45mmol) was added, and the solution was stirred at room temperature for 2 hours. After the reaction was completed, the reaction solution was poured into brine, which was then extracted with ethyl acetate, and concentrated. The obtained residue was purified by silica gel column chromatography (ethyl acetate : hexane = 1 : 1), and the title compound (11mg,

0.032mmol, 8.7%) was obtained.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.16 (2H, d, J=6.0 Hz), 6.49 (1H, d, J=3.6 Hz), 6.56 (1H, dd, J=2.0, 8.0 Hz), 6.65-6.73 (3H, m), 6.77-6.85 (2H, m), 6.93 (1H, dd, J=4.4, 8.0 Hz), 7.01-7.06 (3H, m), 7.85 (1H, d, J=3.6 Hz), 8.10-8.17 (2H, m), 8.58 (1H, t, J=6.0 Hz).

Example T-1. 6-Amino-thieno[2,3-b]pyridine-5-carboxylic acid (5-phenoxy-thiophen-2-ylmethyl)-amide

The title compound (30mg, 78μmol, 76%) was obtained as a white solid from 6-amino-thieno[2,3-b]pyridine-5-carboxylic acid described in Preparation Example T-6 (20mg, 0.10mmol) and C-(5-phenoxy-thiophen-2-yl)-methylaniline described in Preparation Example 26 (21mg, 0.10mmol) according to an analogous method to Example A-26.

MS m/e (ESI) 382.35(MH<sup>+</sup>)

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.52 (2H, d, J=6.0Hz), 6.52 (1H, d, J=3.7Hz), 6.81 (1H, d, J=3.7Hz), 7.09-7.19 (6H, m), 7.33 (1H, dd, J=1.1, 5.9Hz), 7.39 (2H, t, J=8.6Hz), 8.34 (1H, s), 9.21 (1H, m).

Example T-2. Thieno[2,3-b]pyridine-5-carboxylic acid (5-phenoxy-thiophen-2-ylmethyl)-amide

Thieno[2,3-b]pyridine-5-carboxylic acid methyl ester described in Preparation Example T-10 (4.0mg, 21μmol) and lithium hydroxide monohydrate (0.9mg, 21μmol) were dissolved in a mixture solvent of tetrahydrofuran (0.5mL), methanol (50μl) and water (50μl), and the solution was heated under reflux for 1 hour. The reaction solution was cooled to room temperature, then, removal *in vacuo* was carried out,

and thieno[2,3-b]pyridine-5-carboxylic acid was obtained as a lithium salt.

Then, the lithium salt of the resulting thieno[2,3-b]pyridine-5-carboxylic acid, C-(5-phenoxy-thiophen-2-yl)-methylamine described in Preparation Example 26 (4.7mg, 23 $\mu$ mol), benzotriazol-1-yl-tris(dimethylamino)phosphonium hexafluorophosphate (14mg, 32 $\mu$ mol) and triethylamine (9 $\mu$ l, 63 $\mu$ mol) were dissolved in N,N-dimethylformamide (0.5mL), and the solution was stirred at room temperature for 2 hours. Water and ethyl acetate were added to the reaction solution, the organic layer was partitioned, and washed with brine. The solvent was evaporated *in vacuo*, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 1 : 1), and the title compound (3.0mg, 8.2 $\mu$ mol, 40%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>)  $\delta$ (ppm) : 4.75 (2H, d, J=5.2Hz), 6.40 (1H, d, J=3.6Hz), 6.77 (1H, s), 6.93 (1H, d, J=3.2Hz), 7.08-7.12 (3H, m), 7.30-7.34 (3H, m), 7.62 (1H, d, J=6.4Hz), 8.51 (1H, d, J=2.0Hz), 8.92 (1H, d, J=2.0Hz).

Example U-1. 5-Amino-thieno[3,2-b]pyridine-6-carboxylic acid (5-phenoxy-thiophen-2-ylmethyl)-amide

The title compound (85mg, 0.22mmol, 87%) was obtained as a white solid from 5-amino-thieno[3,2-b]pyridine-6-carboxylic acid described in Preparation Example U-4 (50mg, 0.26mmol) and C-(5-phenoxy-thiophen-2-yl)-methylamine described in Preparation Example 26 (53mg, 0.26mmol) according to an analogous method to Example A-26.

<sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>)  $\delta$ (ppm) : 4.52 (2H, d, J=5.7Hz), 6.52 (1H, d, J=3.7Hz), 6.82 (1H, d, J=3.8Hz), 6.99 (2H, s), 7.10 (2H, d, J=7.5Hz), 7.15 (1H, t, J=7.3Hz), 7.18 (1H, d, J=5.5Hz), 7.39 (2H, dd, J=7.3, 8.8Hz), 8.08 (1H, d, J=5.5Hz), 8.49 (1H, s),

9.17 (1H, t, J=5.5Hz).

Example U-2. Thieno[3,2-b]pyridine-6-carboxylic acid  
(5-phenoxy-thiophen-2-ylmethyl)-amide

5 The title compound (7mg, 19μmol, 89%) was obtained as a white solid from trifluoromethanesulfonic acid 6-((5-phenoxy-thiophen-2-ylmethyl)-carbamoyl)-thieno[3,2-b]pyridin-5-yl ester described in Preparation Example U+-2 (11mg, 21μmol) according to an analogous method to Preparation Example T-10.

10 <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 4.57 (2H, d, J=5.7Hz), 6.50 (1H, d, J=3.7Hz), 6.81 (1H, d, J=3.7Hz), 7.08 (2H, d, J=8.1Hz), 7.12 (1H, t, J=7.5Hz), 7.36 (2H, t, J=7.7Hz), 7.62 (1H, d, J=5.7Hz), 8.31 (1H, d, J=5.5Hz), 8.90 (1H, s), 9.08 (1H, d, J=2.0Hz), 9.32-9.38 (1H, m).

15 Example V-1. 1H-Indole-5-carboxylic acid 3-phenoxybenzylamide

The title compound was obtained from 1H-indole-5-carboxylic acid and 4-phenoxybenzylamine described in Preparation Example 3 according to an analogous method to Example H-1.

MS m/e(ESI) 343.15(MH<sup>+</sup>)

20

Example W-1. (4-Benzyloxy-phenyl)-quinolin-6-ylmethyl-amine

To a solution of lithium aluminum hydride (52mg, 1.37mmol) in tetrahydrofuran (10mL) was added a solution of quinoline-6-carboxylic acid (4-benzyloxyphenyl)-amide described in Preparation Example Z+-2 (194mg, 19%) in  
25 tetrahydrofuran, and the solution was stirred under reflux for 3 hours. The reaction

solution was allowed to room temperature, an aqueous solution of saturated ammonium chloride was added thereto, the solution was extracted with ethyl acetate and dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by NH silica gel column chromatography (hexane:ethyl acetate), and the title compound (62mg, 33%) was obtained as a pale yellow solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 3.97(1H, brs), 4.49(2H, s), 4.98(2H, s), 6.61-6.64(2H, m), 6.84-6.86(2H, m), 7.30-7.42(6H, m), 7.73(1H, dd, J=1.6, 8.4Hz), 7.80(1H, s), 8.09(1H, d, J=8.8Hz), 8.12(1H, d, J=8.4Hz), 8.89-8.90(1H, m).

10 Example W-2. (4-Benzyloxy-benzyl)-quinolin-6-yl-amine

To a solution of lithium aluminum hydride (58mg, 1.54mmol) in tetrahydrofuran (10mL) was added a solution of 4-benzyloxy-N-quinolin-6-yl-benzamide described in Preparation Example Z+-3 (218mg, 0.615mmol) in tetrahydrofuran (1mL), and the solution was stirred under reflux for 7 hours. The reaction solution was allowed to room temperature, an aqueous solution of saturated ammonium chloride was added, the solution was extracted with ethyl acetate and dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate), and the title compound (147mg, 70%) was obtained as a yellow solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 4.27(1H, m), 4.36(2H, d, J=4.8Hz), 5.07(2H, s), 6.73(1H, d, J=2.4Hz), 6.98(2H, d, J=8.8Hz), 7.12(1H, dd, J=2.8, 9.2Hz), 7.24-7.27(1H, m), 7.31-7.45(7H, m), 7.87-7.91(2H, m), 8.61-8.62(1H, m).

25 Example W-3. 2-(3-Benzyloxy-phenyl)-1-quinolin-6-yl-ethanone



A mixture of 1-quinolin-6-yl-ethanone described in Preparation Example 131 (171 mg, 1mmol), 1-benzyloxy-3-bromo-benzene (289mg, 1mmol), bis(dibenzylideneacetone)palladium (3mg, 0.0052mmol), 1,1'-bis(diphenylphosphino)ferrocene (5mg, 0.009mmol), potassium *tert*-butoxide (236 mg, 2.1mmol) and tetrahydrofuran (15mL) were stirred under nitrogen atmosphere for 7 hours at 70 °C. Silica gel (80mL) was added to the reaction solution at room temperature, filtration was carried out by silica gel column chromatography (hexane : ethyl acetate = 4 : 6), the filtrate was concentrated *in vacuo*, and a residue of yellow oil (0.191g) was obtained. This residue was purified by thin layer silica gel chromatography (hexane : ethyl acetate = 1 : 10), and a yellow oily residue (98mg) was obtained. In addition, the residue was purified by NH silica gel column chromatography (hexane : ethyl acetate = 85 : 15), and the title compound (50mg, 0.14mmol, 14%) was obtained as a pale yellow oil.

<sup>1</sup>H-NMR Spectrum (Acetone-d<sub>6</sub>) δ(ppm) : 4.51 (2H, s), 5.10 (2H, s), 6.89-6.91(1H, m), 6.96-6.98(1H, m), 7.07(1H, m), 7.25(1H, t, J=4.4 Hz), 7.29-7.33(1H, m), 7.35-7.39(2H, m), 7.44-7.47(2H, m), 7.61 (1H, dd, J=8.4, 4.4 Hz), 8.11 (1H, d, J=8.8 Hz), 8.31 (1H, dd, J=8.8, 2.0 Hz), 8.50-8.53 (1H, m), 8.81 (1H, d, J=2.0 Hz), 9.02 (1H, m).

#### Example W-4. 2-(3-Phenoxy-phenyl)-1-quinolin-6-yl-ethanone

According to an analogous method to Example W-3, 1-quinolin-6-yl-ethanone described in Preparation Example 131 (171mg, 1mmol), 1-phenoxy-3-bromo-benzene (274mg, 1.1mmol), bis(dibenzylideneacetone)palladium (10mg, 0.0187mmol), 1,1'-bis(diphenylphosphino)ferrocene (12.5mg, 0.0225mmol), potassium *tert*-butoxide (236mg, 2.1mmol) and tetrahydrofuran (15mL) were stirred at 70°C for 6 hours under nitrogen atmosphere, and the title compound (64mg,

0.189mmol, 19%) was obtained as a pale yellow oily substance.

<sup>1</sup>H-NMR Spectrum (Acetone-d<sub>6</sub>) δ(ppm) : 4.56 (2H, s), 6.88-6.91(1H, m),  
6.98-7.00(2H, m), 7.06-7.16(3H, m), 7.32-7.37(3H, m), 7.62 (1H, dd, J=8.4, 4.4 Hz),  
8.12 (1H, d, J=8.8 Hz), 8.31 (1H, dd, J=8.8, 2.0 Hz), 8.49-8.52 (1H, m), 8.81 (1H, d,  
5 J=2.0 Hz), 9.02 (1H, m).

Example W-5. 6-(4-Benzyloxy-benzyloxy)-quinoline

To a solution of quinoline-6-ol (37mg, 0.25mmol),  
1-benzyloxy-4-chloromethyl-benzene (70mg, 0.30mmol) in dimethylsulfoxide (2.5mL)  
10 was added potassium *tert*-butoxide (43mg, 0.38mmol) under nitrogen atmosphere,  
and the solution was stirred for 13 hours at room temperature. This reaction  
solution was purified by thin layer silica gel chromatography (hexane : ethyl acetate =  
1 : 1), and the title compound (74mg, 0.217mmol, 86%) was obtained as a pale  
yellow solid.

15 <sup>1</sup>H-NMR Spectrum (Acetone-d<sub>6</sub>) δ(ppm) : 5.16 (2H, s), 5.19 (2H, s), 7.05-7.09(2H, m),  
7.31-7.35(1H, m), 7.38-7.46(5H, m), 7.46-7.51(4H, m), 7.95(1H, d, J=8.8 Hz),  
8.19(1H, dd, J=8.4, 0.8 Hz), 8.74(1H, dd, J=4.0, 1.6 Hz).

Example W-6. 6-(3-Phenoxy-benzylsulfanyl)-quinoline

20 To a solution of (3-phenoxy-phenyl)-methanol (2.0g, 10mmol) in  
tetrahydrofuran (50ml) was added potassium *tert*-butoxide (1.35g, 12mmol) under  
nitrogen atmosphere and an ice-cold stirring, then, 4-methyl-benzenesulfonyl chloride  
(2.48g, 13mmol) was added, and the solution was stirred at room temperature for 21  
hours. An aqueous solution of saturated ammonium chloride was added, the  
25 solution was extracted with ethyl acetate (200ml), washed with brine (150ml), dried

over anhydrous magnesium sulfate, then, the solvent was evaporated, and a residue of pale yellow oil (3.97g) was obtained. This residue was purified by silica gel column chromatography (hexane : ethyl acetate = 4 : 1), and toluene-4-sulfonic acid 3-phenoxybenzyl ester (2.93g, 8.27mmol, 82%) was obtained as a white solid.

5           To a solution of dithiocarbonic acid O-ethyl ester S-quinolin-6-yl ester described in Preparation Example W-1 (50mg, 0.201mmol), toluene-4-sulfonic acid 3-phenoxybenzyl ester (110mg, 0.313mmol) and methanol (1ml) in tetrahydrofuran (5ml) was added potassium *tert*-butoxide (135mg, 1.20mmol) under nitrogen atmosphere, which was then stirred at room temperature for 24 hours. This reaction  
10 solution was purified by thin layer silica gel chromatography (hexane : ethyl acetate = 1 : 1), and the title compound (39mg, 0.114mmol, 57%) was obtained as a yellow oil.  
<sup>1</sup>H-NMR Spectrum (Acetone-d<sub>6</sub>) δ(ppm) :4.36(2H, s), 6.85-6.90(3H, m), 7.02-7.03(1H, m), 7.07-7.11(1H, m), 7.20-7.22(1H, m), 7.27-7.34(3H, m), 7.49(1H, dd, J=8.4, 4.0 Hz), 7.69(1H, dd, J=8.8, 2.0 Hz), 7.84-7.85(1H, m), 7.94(1H, d, J=8.8 Hz),  
15 8.18-8.20(1H, m), 8.84-8.85(1H, m).

Example X-1. (4-*tert*-Butyl-benzyl)-quinazolin-4-yl-amine

To a solution of 4-chloro-quinazoline described in Preparation Example X-1 (8mg, 0.049mmol) in 1-methyl-2-pyrrolidone (0.5mL) were added  
20 4-*tert*-butyl-benzylamine (10μl, 0.059mmol) and N,N-diisopropylethylamine (17μl, 0.098mmol), and the solution was stirred for 5 hours at 160°C. The reaction mixture was directly purified by reverse phase high performance liquid chromatography (acetonitrile-water mobile phase (containing 0.1% trifluoroacetic acid) was used), and the title compound (16.5mg, 0.041mmol, 72%) was obtained as a trifluoroacetic acid  
25 salt.

MS m/e(ESI) 292.20(MH<sup>+</sup>)

Example X-2. (4-Benzyloxy-benzyl)-quinazolin-4-yl-amine

The title compound (9.63mg, 0.021mmol, 38%) was obtained as a  
5 trifluoroacetic acid salt from 4-chloro-quinazoline described in Preparation Example  
X-1 (9mg, 0.055mmol) and 4-benzyloxy-benzylamine described in Preparation  
Example 1 (12mg, 0.055mmol) according to an analogous method to Example X-1.

MS m/e(ESI) 342.27(MH<sup>+</sup>)

10 Example Y-1. N<sup>4</sup>-(4-Benzyloxy-benzyl)-pyrido[2,3-d]pyrimidine-2,4-diamine

To a mixture of 2-amino-N-(4-benzyloxy-benzyl)-thionicotinamide described in  
Example A-22 (30mg, 0.083mmol) and toluene (1mL) was added benzyl bromide  
(0.044mL, 0.37mmol), and the solution was refluxed for 3 hours. After cooling, the  
solvent was evaporated *in vacuo*, and the residue was washed with diethyl ether  
15 twice. A mixture of a portion (13mg) of the resulting crude product (33mg),  
cyanamide (20mg, 0.48mmol) and N-methylpyrrolidinone (1mL) was stirred for 2  
hours at 120°C. After cooling, the reaction solution was directly purified by reverse  
phase high performance liquid chromatography (acetonitrile-water mobile phase  
(containing 0.1% trifluoroacetic acid) was used), and ditrifluoroacetic acid salt of the  
20 title compound (0.75mg, 0.0013mmol, 4.2%) was obtained.

MS m/e (ESI) 358.2 (MH<sup>+</sup>)

Example Z-1.(4-Butyl-3-methyl-phenyl)-quinolin-6-yl-metanone

To a solution of (4-butyl-3-methyl-phenyl)-quinoline-6-yl-methanol described  
25 in Preparation Example Z+-1 (152mg, 0.50mmol) in chloroform (1.0mL) was added

active manganese dioxide (510mg, 5.0mmol), and the solution was stirred at room temperature for 5 hours. Manganese dioxide was removed by filtering through Celite pad, the filtrate was concentrated, the residue was purified by silica gel column chromatography (hexane : ethyl acetate), and the title compound (56mg, 37%) was obtained as a pale yellow oil.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 0.98(3H, t, J=7.2Hz), 1.40-1.49(2H, m), 1.58-1.66(2H, m), 2.39(3H, s), 2.70(2H, t, J=8.0Hz), 7.26-7.28(1H, m), 7.48-7.51(1H, m), 7.60-7.62(1H, m), 7.67(1H, s), 8.13-8.16(1H, m), 8.20-8.22(1H, m), 8.25-8.27(2H, m), 9.03-9.04(1H, m).

#### Example Z-2. Quinoline-6-carboxylic acid (3-benzyloxyphenyl)-amide

Thionyl chloride (2mL) was added to 6-quinolinecarboxylic acid (100mg, 0.577mmol), and the solution was stirred for 2 hours under reflux. Then, the solution was allowed to cool to room temperature, and excess thionyl chloride was evaporated *in vacuo*. The obtained residue was dissolved in N,N-dimethylformamide (2mL), 3-benzyloxyaniline (115mg, 0.577mmol), triethylamine (0.12mL, 0.866mmol) and dimethylaminopyridine (1mg) were added, and the solution was stirred overnight at room temperature. Ethyl acetate and water were added to the reaction solution, which was then partitioned, the organic layer was washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated, the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 1 : 1), and the title compound (53mg, 26%) was obtained as a white solid.

<sup>1</sup>H-NMR Spectrum (CDCl<sub>3</sub>) δ(ppm) : 5.12 (2H, s), 6.80-6.83(1H, m), 7.16-7.18(1H, m), 7.27-7.35(2H, m), 7.38-7.42(2H, m), 7.45-7.47(2H, m), 7.49-7.52(1H, m),

7.58-7.59(1H, m), 8.03(1H, s), 8.13(1H, dd, J=2.0, 8.8Hz), 8.21(1H, d, J=8.8Hz), 8.25-8.28(1H, m), 8.38(1H, d, J=2.0Hz), 9.01-9.03(1H, m).

Example Z-3. 3-Phenoxy-N-quinolin-6-yl-benzamide

5           To a solution of 6-aminoquinoline (250mg, 1.04mmol) in N,N-dimethylformamide (15mL) were added 3-phenoxybenzoic acid (371mg, 1.73mmol), O-benzotriazol-1-yl-N,N,N',N'-tetramethyluronium hexafluorophosphate (722mg, 1.90mmol) and triethylamine (603μl, 4.33mmol), and the solution was stirred for 2 days at room temperature. Water was added to the reaction mixture, which  
10 was extracted with ethyl acetate, the organic layer was washed with water, an aqueous solution of saturated sodium bicarbonate and brine, dried over anhydrous magnesium sulfate, and then, concentrated *in vacuo*. The obtained residue was purified by silica gel column chromatography (hexane-ethyl acetate), and the title compound (328mg, 0.964mmol, 93%) was obtained.

15       <sup>1</sup>H-NMR Spectrum (DMSO-d<sub>6</sub>) δ(ppm) : 7.08 (2H, d, J=8.4Hz), 7.18 (1H, t, J=7.6Hz), 7.24 (1H, d, J=8.0Hz), 7.43 (2H, dd, J=7.6, 8.4Hz), 7.48 (1H, dd, J=4.0, 8.4Hz), 7.56 (1H, dd, J=7.6, 8.0Hz), 7.63 (1H, s), 7.79 (1H, d, J=7.6Hz), 7.94-8.15 (2H, m), 8.29 (1H, d, J=8.0Hz), 8.50 (1H, s), 8.80 (1H, dd, J=2.0, 4.0Hz), 10.6 (1H, s).

20       Example Z-4. 1H-Indole-5-carboxylic acid (4-benzyloxy-phenyl) amide

The title compound was obtained from 1H-indole-5-carboxylic acid and 4-benzyloxy-phenylamine hydrochloride according to an analogous method to Example H-1

MS m/e(ESI) 343.30(MH<sup>+</sup>)

The structural formulae of the compounds obtained in the above preparation examples and examples are shown in the following Table 1 to Table 51.

5

10

15


20

25


[0095]

[Table 1]


Preparation Example A-1




Preparation Example A-2




Preparation Example A-3




Preparation Example A-4




Preparation Example A-5




Preparation Example A-6




Preparation Example A-7




Preparation Example A-8




Preparation Example A-9




Preparation Example A-10




Preparation Example A-11




Preparation Example A-12




Preparation Example A-13




Preparation Example A-14




Preparation Example A-15




Preparation Example A-16



Preparation Example A-17



Preparation Example A-18







[0099]

[Table 2]


Preparation Example A-19




Preparation Example A-20




Preparation Example A-21




Preparation Example A-22




Preparation Example A-23




Preparation Example A-24




Preparation Example A-25




Preparation Example A-26




Preparation Example A-27




Preparation Example A-28



Preparation Example AA-1



Preparation Example AA-2




Preparation Example B-1




[0097]

[Table 3]


Preparation Example A+-1




Preparation Example A+-7




Preparation Example A+-2




Preparation Example A+-8




Preparation Example A+-3




Preparation Example A+-9




Preparation Example A+-4




Preparation Example A+-10




Preparation Example A+-5




Preparation Example A+-11



Preparation Example A+-6




Preparation Example A+-12




[0098]

[Table 4]


Preparation Example A+-13




Preparation Example A+-19




Preparation Example A+-14




Preparation Example A+-20




Preparation Example A+-21




Preparation Example A+-22




Preparation Example A+-15




Preparation Example A+-23




Preparation Example A+-16




Preparation Example A+-24




Preparation Example A+-17




Preparation Example A+-25



Preparation Example A+-18




Preparation Example A+-26




[0099]

[Table 5]


Preparation Example C-1




Preparation Example C-2




Preparation Example C-3




Preparation Example C-4




Preparation Example C-5




Preparation Example D-1




Preparation Example E-1




Preparation Example F-1




Preparation Example F-2




Preparation Example F-3




Preparation Example F-4




Preparation Example G-1




Preparation Example H-1




Preparation Example H-2



Preparation Example H-3




Preparation Example I-1




[0100]

[Table 6]


Preparation Example E+-1




Preparation Example E+-2




Preparation Example E+-3




Preparation Example E+-4




Preparation Example E+-5




Preparation Example E+-6




Preparation Example E+-7




Preparation Example E+-8




Preparation Example E+-9



Preparation Example E+-10




Preparation Example Q+-1




[0101]

[Table 7]


Preparation Example J-1




Preparation Example J-2




Preparation Example J-3




Preparation Example J-4




Preparation Example J-5




Preparation Example J-6




Preparation Example J-7




Preparation Example K-1




Preparation Example K-2




Preparation Example O-1




Preparation Example O-2




Preparation Example P-1




Preparation Example P-2




Preparation Example P-3




Preparation Example P-4




Preparation Example S-1




Preparation Example S-2



Preparation Example S-3




Preparation Example S-4




[0102]

[Table 8]


Preparation Example R-1




Preparation Example R-2




Preparation Example R-3




Preparation Example R-4




Preparation Example R-5




Preparation Example R-6




Preparation Example R-7





Preparation Example T-1




Preparation Example T-2




Preparation Example T-3

Preparation Example  
T-4


Preparation Example T-5




Preparation Example T-6





Preparation Example T-7



Preparation Example T-8




Preparation Example T-9

Preparation Example  
T-10


[0103]

[Table 9]


Preparation Example U-1




Preparation Example U-2




Preparation Example U-3




Preparation Example U-4




Preparation Example W-1




Preparation Example X-1




Preparation Example Z-1




Preparation Example Z-2




Preparation Example U+-1




Preparation Example U+-2




Preparation Example Z+-1



Preparation Example Z+-2



Preparation Example Z+-3







[0104]

[Table 10]


Preparation Example 1




Preparation Example 2




Preparation Example 3




Preparation Example 4




Preparation Example 5




Preparation Example 6




Preparation Example 7




Preparation Example 8




Preparation Example 9




Preparation Example 10




Preparation Example 11




Preparation Example 12




Preparation Example 13




Preparation Example 14




Preparation Example 15




Preparation Example 16



Preparation Example 17




Preparation Example 18




[0105]

[Table 11]


Preparation Example 19




Preparation Example 20




Preparation Example 21




Preparation Example 22




Preparation Example 23




Preparation Example 24




Preparation Example 25




Preparation Example 26




Preparation Example 27




Preparation Example 28




Preparation Example 29




Preparation Example 30




Preparation Example 31




Preparation Example 32




Preparation Example 33




Preparation Example 34



Preparation Example 35





















Preparation Example 36



[0106]

[Table 12]

Preparation Example 37	Preparation Example 43	Preparation Example 49
		
Preparation Example 38	Preparation Example 44	Preparation Example 50
		
Preparation Example 39	Preparation Example 45	Preparation Example 51
		
Preparation Example 40	Preparation Example 46	Preparation Example 52
		
Preparation Example 41	Preparation Example 47	Preparation Example 53
		
Preparation Example 42	Preparation Example 48	Preparation Example 54
		



















[0107]

[Table 13]

Preparation Example 55	Preparation Example 61	Preparation Example 67
Preparation Example 56	Preparation Example 62	Preparation Example 68
Preparation Example 57	Preparation Example 63	Preparation Example 69
Preparation Example 58	Preparation Example 64	Preparation Example 70
Preparation Example 59	Preparation Example 65	Preparation Example 71
Preparation Example 60	Preparation Example 66	Preparation Example 72


[0108]

[Table 14]

<p>Preparation Example 73</p> 	<p>Preparation Example 79</p> 	<p>Preparation Example 85</p> 
<p>Preparation Example 74</p> 	<p>Preparation Example 80</p> 	<p>Preparation Example 86</p> 
<p>Preparation Example 75</p> 	<p>Preparation Example 81</p> 	<p>Preparation Example 87</p> 
<p>Preparation Example 76</p> 	<p>Preparation Example 82</p> 	<p>Preparation Example 88</p> 
<p>Preparation Example 77</p> 	<p>Preparation Example 83</p> 	<p>Preparation Example 89</p> 
<p>Preparation Example 78</p> 	<p>Preparation Example 84</p> 	<p>Preparation Example 90</p> 



















[0109]

[Table 15]





















[0110]

[Table 16]

<p>Preparation Example 109</p> 	<p>Preparation Example 115</p> 	<p>Preparation Example 121</p> 
<p>Preparation Example 110</p> 	<p>Preparation Example 116</p> 	<p>Preparation Example 122</p> 
<p>Preparation Example 111</p> 	<p>Preparation Example 117</p> 	<p>Preparation Example 123</p> 
<p>Preparation Example 112</p> 	<p>Preparation Example 118</p> 	<p>Preparation Example 124</p> 
<p>Preparation Example 113</p> 	<p>Preparation Example 119</p> 	<p>Preparation Example 125</p> 
<p>Preparation Example 114</p> 	<p>Preparation Example 120</p> 	<p>Preparation Example 126</p> 

[0111]



















[Table 17]

Preparation Example 127 	Preparation Example 133 	Preparation Example 139 
Preparation Example 128 	Preparation Example 134 	Preparation Example 140 
Preparation Example 128 	Preparation Example 135 	Preparation Example 141 
Preparation Example 130 	Preparation Example 136 	Preparation Example 142 
Preparation Example 131 	Preparation Example 137 	Preparation Example 143 
Preparation Example 132 	Preparation Example 138 	Preparation Example 144 




[0112]

[Table 18]



<p>Preparation Example 145</p> 	<p>Preparation Example 146</p> 	<p>Preparation Example 157</p> 
<p>Preparation Example 146</p> 	<p>Preparation Example 152</p> 	<p>Preparation Example 158</p> 
<p>Preparation Example 147</p> 	<p>Preparation Example 153</p> 	<p>Preparation Example 159</p> 
<p>Preparation Example 148</p> 	<p>Preparation Example 154</p> 	<p>Preparation Example 160</p> 
<p>Preparation Example 149</p> 	<p>Preparation Example 155</p> 	<p>Preparation Example 161</p> 
<p>Preparation Example 150</p> 	<p>Preparation Example 156</p> 	<p>Preparation Example 162</p> 

[0113]


[Table 19]

Preparation  
Example 163


Preparation Example 170

Preparation  
Example 175


Preparation Example 164




Preparation Example 171




Preparation Example 177




Preparation Example 165






Preparation Example 172







Preparation Example 178





Preparation Example 166

Preparation  
Example 173Preparation  
Example 179


Preparation Example 167

Preparation  
Example 174Preparation  
Example 180Preparation  
Example 168

Preparation Example 175


Preparation  
Example 181

Preparation Example 169



[0114]


[Table 20]




[0115]

[Table 21]


Example A-1




Example A-7




Example A-2




Example A-8




Example A-3




Example A-9




Example A-4




Example A-10




Example A-5




Example A-11



Example A-6




Example A-12




[0116]

[Table 22]


Example A-13




Example A-14




Example A-15




Example A-16




Example A-17




Example A-18




Example A-19




Example A-20




Example A-21




Example A-22



Example A-23




Example A-24




[0117]

[Table 23]


Example A-25




Example A-26




Example A-27




Example A-28




Example A-29




Example A-30




Example A-31




Example A-32




Example A-33




Example A-34



Example A-35




Example A-36




[0118]

[Table 24]


Example A-37




Example A-38




Example A-39




Example A-40




Example A-41




Example A-42




Example A-43




Example A-44




Example A-45




Example A-46



Example A-47




Example A-48




[0119]

[Table 25]


Example A-49




Example A-50




Example A-51




Example A-52




Example A-53




Example A-54




Example A-55




Example A-56




Example A-57




Example A-58



Example A-59



Example A-60







[0120]

[Table 26]


Example A-61




Example A-62




Example A-67




Example A-68




Example A-69




Example A-64




Example A-70




Example A-65




Example A-71



Example A-66




Example A-72




[0121]

[Table 27]


Example A-73




Example A-79




Example A-74




Example A-80




Example A-75




Example A-81




Example A-76




Example A-82




Example A-77




Example A-83



Example A-78




Example A-84




[0122]

[Table 28]


Example A-85




Example A-91




Example A-86




Example A-92




Example A-87




Example A-93




Example A-88




Example A-94




Example A-89




Example A-95



Example A-90




Example A-96




[0123]

[Table 29]


Example A-97




Example A-103




Example A-98




Example A-104




Example A-99




Example A-105



Example A-100




Example A-106




Example A-101




Example A-107



Example A-102




Example A-108




[0124]

[Table 30]


Example A-109




Example A-110




Example A-111




Example A-112




Example A-113




Example A-114




Example A-115




Example A-116




Example A-117




Example A-118



Example A-119




Example A-120




[0125]

[Table 31]


Example A-121




Example A-127




Example A-122




Example A-128




Example A-123




Example A-129




Example A-124




Example A-130




Example A-125




Example A-131



Example A-126




Example A-132




[0126]

[Table 32]


Example A-133




Example A-134




Example A-135




Example A-136




Example A-137




Example A-138




Example A-139




Example A-140




Example A-141




Example A-142



Example A-143




Example A-144




[0127]

[Table 33]


Example A-145




Example A-151




Example A-146




Example A-152




Example A-147




Example A-153




Example A-148




Example A-154




Example A-149




Example A-155



Example A-150



Example A-156







[0128]

[Table 34]


Example A-157




Example A-158




Example A-159




Example A-160




Example A-161




Example A-162




Example A-163




Example A-164




Example A-165




Example A-166



Example A-167




Example A-168




[0129]

[Table 35]


Example A-169




Example A-170




Example A-171




Example A-172




Example A-173




Example A-174




Example A-175




Example A-176




Example A-177




Example A-178



Example A-179




Example A-180




[0130]

[Table 36]


Example A-181




Example A-187




Example A-182




Example A-188




Example A-183




Example A-189




Example A-184




Example A-190




Example A-185




Example A-191



Example A-186




Example A-192




[0131]

[Table 37]


Example A-193




Example C-1




Example A-194




Example AA-1




Example AA-2



Example B-1




Example B-2




[0132]

[Table 38]


Example D-1




Example D-2




Example D-3




Example D-4




Example D-5




Example D-6




Example D-7




Example D-8



Example D-9




Example D-10




[0133]

[Table 39]


Example E-1




Example E-2




Example E-3




Example E-4




Example E-5




Example E-6




Example E-7




Example E-8




Example E-9




Example E-10



Example E-11




Example E-12




[0134]

[Table 40]


Example E-13




Example E-19




Example E-14




Example E-20




Example E-15




Example E-21




Example E-16




Example E-22




Example E-17




Example E-23



Example E-18




Example E-24




[0135]

[Table 41]


Example E-25




Example E-31




Example E-26




Example E-32




Example E-27




Example E-33




Example E-28




Example E-34




Example E-29




Example E-35



Example E-30



Example E-36







[0136]

[Table 42]


Example E-37




Example E-43




Example E-38




Example E-44




Example E-39




Example E-45




Example E-40




Example E-46




Example E-41




Example E-47



Example E-42




Example E-48




[0137]

[Table 43]


Example E-49




Example E-55




Example E-50




Example E-56




Example E-51




Example E-57




Example E-52




Example E-58




Example E-53




Example E-59



Example E-54




Example E-60




[0138]

[Table 44]


Example E-61




Example E-62




Example E-63




Example E-64




Example E-65




Example E-66




Example E-67




Example E-68




Example E-69




Example E-70



Example E-71




Example E-72




[0139]

[Table 45]


Example E-73




Example E-79




Example E-74




Example E-80




Example E-75




Example E-81




Example E-76




Example E-82



Example E-77




Example E-78




[0140]

[Table 46]


Example F-1




Example G-1




Example H-1



Example I-1




Example J-1




[0141]

[Table 47]


Example K-1




Example L-6




Example L-1




Example L-7




Example L-2




Example L-8




Example L-3




Example L-9




Example L-4




Example L-10



Example L-5




Example L-11




[0142]

[Table 48]


Example M-1




Example Q-4




Example O-1




Example Q-5




Example P-1




Example Q-6




Example Q-1




Example Q-7




Example Q-2



Example Q-8




Example Q-3




[0143]

[Table 49]


Example R-1




Example R-2




Example S-1




Example T-1




Example T-2




Example U-1



Example U-2



Example V-1







[0144]

[Table 50]


Example W-1




Example W-2




Example W-3




Example W-4




Example W-5




Example W-6




Example X-1



Example X-2




Example Y-1




[0145]

[Table 51]


Example Z-1




Example Z-2



Example Z-3



Example Z-4



5

[0146]

The heterocyclic compounds (I) according to the present invention, or salts, or hydrates thereof, demonstrate excellent inhibitory activity on the GPI-anchored protein transport process, anti-*Candida* activity, anti-*Aspergillus* activity. In addition, it is also excellent in terms of property, safety and metabolic stability, and is extremely

useful as a preventive agent or therapeutic agent against fungal infection.

[Pharmacological Test Example]

In order to demonstrate the usefulness of the compounds (I) according to the present invention, 1. the inhibitory activity on the GPI-anchored protein transport process, 2. the anti-*Candida* activity and the anti-*Aspergillus* activity, 3. the activity in the experimental systemic *candida* infection model in mice, and 4. the activity in the experimental respiratory *aspergillus* infection model in mice, were measured for the antifungal activity of the compounds (I) according to the present invention.

10

1. Inhibitory activity on the GPI-anchored protein transport process

A reporter system that reflects the GPI-anchored protein transport process was constructed using cephalosporinase to which the CWP2 carboxyl-terminal sequence, which is known as the cell wall transport signal of a GPI-anchored protein (Van Der Vaat JM et al, J. Bacteriol., 177:3104-3110,1995), was added. Then, using the constructed reporter system, the activity of the present invention compound that inhibits the transport process of the GPI-anchored protein was measured.

(1). Construction of the reporter gene

With pESH plasmid containing the ENO1 promoter, a secretion signal, and a lysozyme gene (Ichikawa K et al, Biosci. Biotech. Biochem., 57(10), 1686-1690, 1993) as template and the oligonucleotides listed in SEQ ID NO. 1 and SEQ ID NO. 2 as primers, the DNA containing a promoter sequence-secretion signal portion was amplified by PCR, and this was subcloned into the BamHI-NotI site of pUC19 (a). In addition, with the *C. freundii* chromosomal DNA as template and the oligonucleotides listed in SEQ ID NO. 3 and SEQ ID NO. 4 as primers, the cephalosporinase gene

25

was PCR-amplified, and this was subcloned into the NspV-XbaI site of pUC19 (b). Similarly, with the *S. cerevisiae* chromosomal DNA as template and the oligonucleotides listed in SEQ ID NO. 5 and SEQ ID NO. 6 as primers, the CWP2 gene was PCR-amplified, and this was subcloned into the XbaI-HindIII site of pUC19 (c). Furthermore, with pYES2 (INVITROGEN) as template and the oligonucleotides listed in SEQ ID NO. 7 and SEQ ID NO. 8 as primers, the CYC1 terminator was PCR-amplified, and this was subcloned into the NotI-KpnI site of pUC19 (d).

After producing the full length ENO1 promoter and secretion signal portion by inserting the BamHI-Sall fragment of pESH into the BamHI-Sall cleavage site of the plasmid into which (a) had been inserted, the cephalosporinase gene (b) excised with NspV-XbaI and the CWP2 gene (c) excised with XbaI-HindIII were inserted into the NspV-HindIII cleavage site. Next, pRCW63T was produced by excising with EcoRI-HindIII, inserting this fragment into the integration vector pRS306 (Sikorski RS et al, Genetics. 122(1):19-27, 1989), and then inserting the CYC1 terminator (d) into the HindIII-KpnI cleavage site

#### (2). Introduction of the reporter gene into *S. cerevisiae*

The *S. cerevisiae* G2-10 strain was cultured by shaking in 10ml of YPD culture medium at 30°C, and the cells were collected at the late logarithmic growth phase ( $2$  to  $5 \times 10^7$  cells/ml). After washing with sterilized water, pRCW63T produced in (1) was introduced by the lithium acetate method (described in the YEASTMAKER™ Yeast Transformation System User Manual) using the YEASTMAKER™ Yeast Transformation System (Clontech). pRCW63T in which the URA3 gene was cleaved with ApaI was used. After culturing in SD(Ura-) culture medium for 3 days at 30°C, the grown colonies were cultured in YPD culture medium.

When the localization of the cephalosporinase activity was confirmed; the

activity was mainly localized in the cell wall, the C-terminal sequence of CWP2 was confirmed to function as a transport signal to the cell wall.

(3). Measurement by a reporter system of the inhibitory activity on the GPI-anchored protein transport process

5        A screening of the compounds was performed using *S. cerevisiae* in which pRCW63T was introduced (*S. cerevisiae* CW63 strain).

After a standing culture in a YPD liquid culture medium for 48 hours at 30°C, 75µl/well of a fungal suspension diluted 10-fold with YPD liquid culture medium (3 to 5 x 10<sup>5</sup> cells/ml) was used to inoculate a V-bottomed 96 well plate containing  
10    25µl/well of test compound dilute solution, and a standing culture was carried out for 48 hours at 30°C. The plate was centrifuged, then, 25µl of supernatant was sampled into a 96 well flat-bottomed plate, the resulting solution serving as the culture medium supernatant fraction.

The precipitated cells were suspended, 75µl/well of zymolyase (Seikagaku  
15    Kogyo) solution prepared with 2.4M sorbitol was added, and allowed to act for 1 hour at 30°C. The plate was centrifuged, then, 10µl of supernatant was sampled in a 96 well flat-bottomed plate, 15µl of phosphate buffer was added, the resulting solution serving as the cell wall fraction.

The pooled sample was added 200µM of nitrocefin solution, after a  
20    predetermined time, the reaction was stopped with a citric acid buffer, then, the optical density at 490 nm was measured, to measure the cephalosporinase activity in the culture medium and the cell wall fraction. A compound that increases the cephalosporinase activity in the culture medium supernatant fraction, or decreases the cephalosporinase activity in the cell wall fraction, was considered as a compound  
25    that inhibits the transport process of GPI-anchored protein to the cell wall.

In addition, fungal growth in the presence of the test compound was visually determined.

## 2. Anti-*Candida* activity and anti-*Aspergillus* activity

### 5 (1). Preparation of fungal suspension

For the *C. albicans* E81022 strain, a fungal suspension from a standing culture for 48 hours at 30°C in a Sabouraud dextrose liquid culture medium (SDB) was diluted 10,000-fold with a 1.3-fold concentrated SDB to adjust to a fungal suspension of 1 to 2 x 10<sup>4</sup> cells/ml. For the *A. fumigatus* Tsukuba strain, -80°C  
10 stored strain was diluted 1,000-fold with a 1.3-fold concentrated SDB to adjust to a fungal suspension of 2 to 3 x 10<sup>3</sup> cells/ml.

### (2). Preparation of an agent dilution plate

Using a U-bottomed 96 well plate, 8 samples/plate (A to H) of sample dilution solutions were prepared. On the 3<sup>rd</sup> row of each plate was dispensed 240µl of  
15 sterilized and distilled water, and on the 2<sup>nd</sup> and 4<sup>th</sup> to 12<sup>th</sup> rows were dispensed 125µl of 4% dimethylsulfoxide solution. Weighed sample was dissolved in dimethylsulfoxide to prepare a 2.5 to 20 mg/ml solution, 10µl of this solution was then added to the 3<sup>rd</sup> row of the prepared plate, and 10 steps of two-fold step dilutions (125µl of solution + 125µl of 4% dimethylsulfoxide) were performed on the plate.  
20 This sample dilution solution was dispensed in the amount of 25µl to a V-bottomed or flat-bottomed 96 well plate for MIC measurement to prepare a sample dilution plate.

### (3). Inoculation of fungal suspension and culture

The fungal suspension prepared in (1) was used in the amount of 75µl/well to inoculate a V-bottomed or flat-bottomed 96 well plate containing 25µl/well of test  
25 compound dilution solution prepared in (2), and a standing culture was carried out

aerobically for 48 hours at 30 to 35°C.

(4). MIC measurement

The minimum concentration that clearly inhibited fungal growth as compared to the control by visual inspection was determined as the minimum inhibitory  
5 concentration (MIC).

The following representative compounds prepared in the examples were measured for their inhibitory activity on the GPI-anchored protein transport process, anti-*candida* activity and anti-*aspergillus* activity by the measurement methods described in 1 and 2. As a result, as shown in Table 52 to Table 54, it was found  
10 that the compounds according to the present invention inhibited the GPI-anchored protein transport process, and have anti-*candida* activity and anti-*aspergillus* activity.

15

20

25

[0147]

[Table 52]

**TABLE 52**

Example Nos.	Reporter System Inhibitory Activity (µg/ml)	Example Nos.	Reporter System Inhibitory Activity (µg/ml)	Example Nos.	Reporter System Inhibitory Activity (µg/ml)
A-5	0.2	A-136	0.78	E-65	0.1
A-8	3.13	A-137	0.78	E-67	0.025
A-10	0.2	A-139	0.78	E-69	0.2
A-22	0.2	A-170	0.39	E-71	0.1
A-37	0.1	A-171	0.1	F-1	1.56
A-39	0.78	B-1	3.13	G-1	1.56
A-53	0.1	C-1	0.1	H-1	1.56
A-54	0.2	D-2	0.39	I-1	0.2
A-68	0.05	E-8	0.1	K-1	0.39
A-70	0.025	E-11	0.2	L-4	0.2
A-81	1.56	E-13	0.2	L-7	0.1
A-82	0.2	E-19	0.2	M-1	0.39
A-92	3.13	E-25	0.2	Q-7	0.39
A-98	0.2	E-43	0.1	R-1	0.2
A-99	0.78	E-44	0.78	U-2	0.2
A-101	0.1	E-45	0.78	V-1	0.78
A-112	3.13	E-51	0.2	W-1	0.78
A-116	0.39	E-53	0.2	W-2	0.39
A-124	1.56	E-57	0.1	Y-1	0.2
A-126	1.56	E-64	0.39		
A-131	0.39	E-73	0.2		



[0148]

[Table 53]

**TABLE 53**

Example Nos.	Anti-Candida Activity (µg/ml)	Example Nos.	Anti-Candida Activity (µg/ml)	Example Nos.	Anti-Candida Activity (µg/ml)
A-5	0.2	A-131	1.56	E-64	0.2
A-8	3.13	A-136	1.56	E-65	0.1
A-10	0.78	A-137	1.56	E-67	0.39
A-22	0.78	A-139	1.56	E-69	0.1
A-37	0.39	A-170	0.78	E-71	0.2
A-39	1.56	A-171	0.2	E-73	0.2
A-53	0.39	B-1	3.13	F-1	3.13
A-54	0.78	C-1	0.1	G-1	6.25
A-68	0.05	D-2	0.78	I-1	0.78
A-70	0.05	E-8	0.2	K-1	0.78
A-81	3.13	E-11	0.39	L-4	0.2
A-82	0.2	E-13	0.78	L-7	0.1
A-92	3.13	E-19	0.78	M-1	1.56
A-98	0.39	E-25	1.56	Q-7	0.39
A-99	3.13	E-43	0.39	R-1	0.39
A-101	0.1	E-44	3.13	U-2	0.78
A-112	3.13	E-45	3.13	W-1	0.39
A-116	0.39	E-51	0.78	W-2	1.56
A-124	1.56	E-53	1.56	Y-1	0.78
A-126	0.78	E-57	0.2		

[0149]

[Table 54]

**TABLE 54**

Example Nos.	Anti-Aspergillus Activity (µg/ml)	Example Nos.	Anti-Aspergillus Activity (µg/ml)	Example Nos.	Anti-Aspergillus Activity (µg/ml)
A-5	0.78	A-126	6.25	E-69	1.56
A-8	6.25	A-131	6.25	E-71	1.56
A-10	1.56	A-139	1.56	E-73	0.78
A-22	3.13	A-171	1.56	I-1	1.56
A-37	3.13	B-1	6.25	K-1	1.56
A-39	1.56	D-2	0.78	L-4	0.2
A-53	0.78	E-8	0.2	L-7	0.78
A-54	1.56	E-11	1.56	M-1	1.56
A-68	0.78	E-13	1.56	Q-7	6.25
A-70	0.78	E-19	1.56	R-1	3.13
A-81	6.25	E-45	3.13	U-2	3.13
A-82	1.56	E-51	3.13	V-1	6.25
A-98	1.56	E-53	3.13	W-1	0.78
A-99	6.25	E-57	6.25	W-2	0.78
A-101	0.39	E-64	6.25	Y-1	6.25
A-112	6.25	E-65	1.56		
A-116	3.13	E-67	0.78		

5

[0150]

3. Experimental systemic *candida* infection model in mice

## (1). Preparation of fungal inoculum

A standing culture of *C. albicans* E81022 strain was carried out for 48 hours at 30°C in Sabouraud dextrose agar medium (SDA), the recovered fungal cells were suspended in sterilized physiological saline. By counting the fungal number on a cytometry plate, the suspension was diluted to  $2 \times 10^7$  cells/ml with sterilized

physiological saline to serve as fungal inoculum.

(2). Infection

The fungal inoculum was used in the amounts of 0.2 ml to inoculate 4.5 to 5.5 week-old female ICR mice in the tail vein ( $4 \times 10^6$  cells/mouse).

5 (3). Treatment

From 0.5 to 1 hour after fungal inoculation, 0.2 ml of agent solution (dissolved or suspended in sterilized physiological saline containing 6.5% dimethylsulfoxide and 3.5% Tween 80) was administered into the stomach using a peroral probe, 3 times every 4 hours. The agent concentration was 2.5 mg/kg, and the number of animals

10 in one group was 5 animals.

(4). Determination of the effect

The protective effect was determined by observing life/death until 14 days after infection and calculating the mean survival days.

As a result, as shown in Table 55, mice administered with the compounds  
15 according to the present invention survived for a long time as compared to the untreated group, and the compounds according to the present invention have been also found to demonstrate anti-*candida* activity *in vivo*.

[0151]

[Table 55]

**TABLE 55**

Example Nos.	Mean Survival Days (Days)
A-10	12.5
A-53	13.4
A-54	10.0
A-68	10.6
A-70	12.6
A-137	12.8
E-65	11.6
R-1	10.4
Non-Administered Group	2.2~4.0

[0152]

**4. *Aspergillus* respiratory infection**

## 5       (1). Increasing susceptibility of mice to infection

In order to increase susceptibility of the mice to infection, 200 mg/kg of 5-fluorouracil was administered subcutaneously 6 days prior to infection.

## (2). Test strain

*A. fumigatus* Tsukuba strain was used.

## 10       (3). Preparation of fungal inoculum

A spore suspension of the infection strain was used to coat a potato dextrose agar medium (PDA), a standing culture was carried out for 4 to 7 days at 35°C, spore was then suspended in a sterilized physiological saline containing 0.05% Tween 80. By counting the pore count on a cytometry plate, the suspension was diluted to 6 x 10<sup>5</sup> cells/ml with sterilized physiological saline containing 0.05% Tween 80 to serve

as fungal inoculum.

(4). Infection

Under ketalar anesthesia, 50 $\mu$ l of fungal inoculum was used to inoculate 8 to 9 week-old female DBA/2N mice in the nose ( $3 \times 10^4$  cells/mouse) .

5 (5). Treatment

From 1 hour after fungal inoculation, 0.2 ml of agent solution (dissolved or suspended in sterilized physiological saline containing 6.5%DMSO and 3.5% Tween 80), was administered into the stomach using a peroral probe 3 times per day for 3 days. The agent dosage was from 20 or 40 mg/kg, and the number of animals in  
10 one group was 5 animals.

(6). Determination of the effect

The protective effect was determined by observing life/death until 14 days after infection and calculating the mean survival days.

As a result, as shown in Table 56, mice administered with the compounds  
15 according to the present invention survived for a long time as compared to the untreated group, and the compounds according to the present invention have been also found to demonstrate anti-*aspergillus* activity *in vivo*.

[0153]

[Table 56]

**TABLE 56**

<b>Example Nos.</b>	<b>Mean Survival Days (Days)</b>
<b>A-11</b>	<b>10.4</b>
<b>A-31</b>	<b>11.2</b>
<b>A-32</b>	<b>11.8</b>
<b>A-55</b>	<b>12.6</b>
<b>A-56</b>	<b>10.8</b>
<b>D-4</b>	<b>10.8</b>
<b>D-7</b>	<b>10.6</b>
<b>Non-Administered Group</b>	<b>4.0~4.2</b>

[Industrial Applicability]

[0154]

- 5           The heterocyclic compounds (I) according to the present invention, or a salt or a hydrate thereof, 1) demonstrate effects against occurrence, development and persistence of infectious disease by inhibiting the expression of cell wall surface layer protein and inhibiting cell wall assembly, while at the same time, inhibiting adhesion of fungus onto the cells, preventing the pathogen from showing pathogenicity, in
- 10 addition, 2) are also excellent in terms of property, safety and metabolic stability, and extremely useful as a preventive or therapeutic agent of fungal infection.